

City University of New York (CUNY)

CUNY Academic Works

Publications and Research

Medgar Evers College

2007

Maintaining phase of the tri-phasic crab pyloric rhythm

Christina Mouser

CUNY Medgar Evers College

Farzan Nadim

New Jersey Institute of Technology

Amitabha Bose

New Jersey Institute of Technology

[How does access to this work benefit you? Let us know!](#)

More information about this work at: https://academicworks.cuny.edu/me_pubs/7

Discover additional works at: <https://academicworks.cuny.edu>

This work is made publicly available by the City University of New York (CUNY).

Contact: AcademicWorks@cuny.edu

Poster presentation

Open Access

Maintaining phase of the tri-phasic crab pyloric rhythm

Christina Mouser*¹, Farzan Nadim² and Amitabha Bose²

Address: ¹Department of Mathematical Sciences, Medgar Evers College of CUNY, Brooklyn, NY 11225, USA and ²Department of Mathematical Sciences, New Jersey Institute of Technology, Newark, NJ 07102, USA

Email: Christina Mouser* - cmouser@mec.cuny.edu

* Corresponding author

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007
Toronto, Canada. 7–12 July 2007

Published: 6 July 2007

BMC Neuroscience 2007, 8(Suppl 2):P97 doi:10.1186/1471-2202-8-S2-P97

© 2007 Mouser et al; licensee BioMed Central Ltd.

Introduction

Synaptic depression is a type of short-term plasticity that is observed in many rhythmically active networks [1]. We examine the role that synaptic depression plays in determining the phase of a group of neurons of the crab stomatogastric nervous system. We mathematically construct and analyze a model network consisting of an oscillator neuron that inhibits two follower neurons. We show that constant phase maintenance can be achieved solely through the interplay of the two follower neurons due to the depressive nature of their synaptic connectivity.

Model

The network we are studying is comprised of three neurons, AB, LP, and PY. AB is the pacemaker neuron of the pyloric network while LP and PY are the follower neurons. The activity of these neurons is modeled using Morris-Lecar type first order differential equations. LP and PY receive synaptic inputs from one another and from AB; see figure 1. The synapses are depressing and inhibitory meaning that the strength of the synapse weakens while the pre-synaptic neuron is active. The set of equations used to model this network is similar in form to that of [2].

Results and conclusion

To understand the effect of the depressing synapses between LP and PY, we derived a set of equations that allows us to determine the time at which LP and PY enter their active states relative to when AB enters its active state. This then allowed us to determine which parameters most

significantly contribute to the phase of LP, Φ_{LP} , and the phase of PY, Φ_{PY} . We found that reciprocal inhibition between LP and PY leads to better phase maintenance than when AB is the sole input to these neurons. This occurs because when the reciprocal inhibition is present, Φ_{LP} and Φ_{PY} are determined mostly by the synaptic properties rather than by their membrane kinetics. In addition, we found that when the input from the oscillator neuron




Figure 1
Crab pyloric network.

AB is not depressing, the connectivity between LP and PY is in fact able to produce phase maintenance. The essential property necessary to produce this phase maintenance is for the synapses to increasingly recover from the synaptic depression as the period increases. However, phase maintenance is optimal when the synapse from AB is depressing. Furthermore, these analytic results can be compared to experimental data and can be used to make predictions about the biological network in the absence of synaptic depression from the group pacemaker.

Acknowledgements

This work was supported in part by PSC-CUNY PSCREG-37-74, the National Science Foundation DMS-0315862 (CA), the National Science Foundation DMS-0615168 (AB) and the National Institutes of Health MH-60605 (FN).

References

1. Marder E: **From biophysics to models of network function.** *Annu Rev Neurosci* 1998, **21**:25-45.
2. Manor Y, Bose A, Booth V, Nadim F: **Contribution of synaptic depression to phase maintenance in a model rhythmic network.** *J Neurophysiol* 2003, **90**:3513-3528.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

