

City University of New York (CUNY)

CUNY Academic Works

Publications and Research

Lehman College

2013

Diffraction Effects and General Boundary Conditions in Casimir Energy

Dimitra Karabali
CUNY Lehman College

V P. Nair
CUNY City College of New York

[How does access to this work benefit you? Let us know!](#)

More information about this work at: https://academicworks.cuny.edu/le_pubs/33

Discover additional works at: <https://academicworks.cuny.edu>

This work is made publicly available by the City University of New York (CUNY).
Contact: AcademicWorks@cuny.edu

Diffractive effects and general boundary conditions in Casimir energyDimitra Karabali^{1,*} and V. P. Nair^{2,†}¹*Department of Physics and Astronomy, Lehman College of the CUNY, Bronx, New York 10468, USA*²*Physics Department, City College of the CUNY, New York, New York 10031, USA*

(Received 7 April 2013; published 31 May 2013)

The effect of edges and apertures on the Casimir energy of an arrangement of plates and boundaries can be calculated in terms of an effective nonlocal lower-dimensional field theory that lives on the boundary. This formalism has been developed in a number of previous papers and applied to specific examples with Dirichlet boundary conditions. Here we generalize the formalism to arbitrary boundary conditions. As a specific example, the geometry of a flat plate and a half-plate placed parallel to it is considered for a number of different boundary conditions, and the area-dependent and edge-dependent contributions to the Casimir energy are evaluated. While our results agree with known results for those special cases (such as the Dirichlet and Neumann limits) for which other methods of calculation have been used, our formalism is suitable for general boundary conditions, especially for the diffractive effects.

DOI: [10.1103/PhysRevD.87.105021](https://doi.org/10.1103/PhysRevD.87.105021)

PACS numbers: 03.70.+k, 42.25.Fx

I. INTRODUCTION

The Casimir effect [1], the classic example of the influence of boundary conditions in quantum field theory, has been of considerable interest over the last several years. A vast body of literature has emerged with a number of new geometries being explored, both analytically and numerically and both at zero and nonzero temperatures [2]. Recent developments in nanomachinery have also provided further impetus to these efforts. Geometries with edges and apertures are a particularly interesting set because of the possible interplay of diffractive effects and boundary conditions on the fields. The analytic calculation of the propagators in the given geometry with the given boundary conditions is rarely feasible and numerical and analytical approximations have to be used; diffractive effects are then a formidable computational task. We have recently developed a formalism which focuses on the diffractive effects [3–6]. In our approach, an effective nonlocal lower-dimensional field theory defined on the boundaries, with a boundary action S_B , is first extracted. In the subsequent analysis of this lower-dimensional theory, apertures and edges can be easily incorporated, leading to a systematic calculational scheme for diffractive effects. The results, both for zero and nonzero temperatures, were in good agreement with alternate methods of calculation, such as the numerical worldline method [7,8], and the scattering matrix method [9,10] and in those geometries where such calculations had been done. We also obtained a universal low-energy theorem for the Casimir force between holes on a plate [4].

In this paper, we extend this formalism, addressing two important and related issues. The boundary action S_B is a functional of the field on the boundary, and hence it is

naturally defined for Dirichlet boundary conditions for the propagator or the fields. So the first question is: How do we apply our formalism to the case of Neumann or even more general boundary conditions? Although the case of two full parallel plates with no edge or diffractive effects for a scalar field with Robin boundary conditions has been studied before in Refs. [11,12], our focus here is on the edge and diffractive effects. While Dirichlet, Neumann, or Robin are the easiest to work with, the general boundary conditions, say on a Laplace-type kinetic energy operator, are given by the von Neumann theory of self-adjoint extensions characterized by a self-adjoint operator on the space of boundary values [13,14]. However, there is a problem with the simple and straightforward generalization to arbitrary boundary conditions because, generically, the (minus) Laplace operator has negative eigenvalues, suggesting an instability for the quantum theory [14]. On the other hand, physically, the boundary of interest is produced by the insertion of a material plate or some such object, and it would be rather strange if this process leads to a global instability. So the next question would be: How do we square these concerns? Further, it is not clear that, in a full fledged interacting quantum field theory, mimicking the material plate by boundary conditions is an adequate characterization. These are some of the issues discussed in this paper.

We start with a quick résumé of the boundary action method. We then generalize the method to arbitrary boundary conditions in Sec. III. In Sec. IV, we apply this to the case of two parallel plates with Robin boundary conditions, one of them being a semi-infinite plate so that edge effects and diffraction at the edge become important. The analytical part of the calculation is explained in some detail, followed by the numerical evaluation of some of the integrals needed. A variety of values of the Robin parameter are considered. The case of two full plates (i.e., with no edge effects) with Robin boundary conditions [11,12] and

*dimitra.karabali@lehman.cuny.edu

†vpn@sci.ccny.cuny.edu

the geometry of a plate and half-plate but restricted to Neumann-Neumann conditions [9,10] have been analyzed by other methods previously. We give a comparison of our results with these in the discussion section. Some other interesting features of our calculation are also pointed out there. The paper concludes with an Appendix where some of the mathematical calculations we have done are outlined.

II. RÉSUMÉ OF BOUNDARY ACTION METHOD

We start with a brief summary of the approach developed in Refs. [3–5]. We will consider a scalar field theory with action $S(\phi)$ in a cubic box of volume V , with $V \rightarrow \infty$ eventually. For simplicity, we start off with a free field theory with a kinetic energy operator given by the Laplacian, i.e., $S(\phi) = \frac{1}{2} \int (\partial\phi)^2$. The quantity of interest for the Casimir energy is the functional integral (or partition function),

$$Z = \int [d\phi] \exp[-S(\phi)]. \quad (1)$$

Our basic strategy is to consider the box as divided into a left region V_L and a right region V_R as shown in Fig. 1, with an interface (shown as the dashed line) at, say, $x_1 = b$. Then the functional integral can be done in two stages. We integrate over all field configurations in V_L with a fixed value of the field (say, $\phi = \varphi$) on the interface and similarly for the right region V_R . Each integration leads to a functional of φ , which we will denote by Ψ_L and $\tilde{\Psi}_R$, respectively, up to constant factors Z_L, Z_R , so that

$$Z = Z_L Z_R \int [d\varphi] \exp[-S_B(\varphi)] \quad (2)$$

$$\Psi_L \tilde{\Psi}_R = \exp[-S_B(\varphi)].$$

The final integration over the values of the field φ at the interface completes the functional integration in Eq. (1). If there is a plate placed at $x_1 = b$ with the field vanishing on it, then this last integration is trivial; we just set $\varphi = 0$. If we have a plate with an opening, then $\varphi = 0$ everywhere on the interface except at the opening. The integration of $\exp(-S_B(\varphi))$ with φ restricted to being nonzero only at




FIG. 1. Schematics of splitting the functional integration into that over fields in two regions V_L and V_R .

the opening gives a boundary contribution to the partition function, which will include all the diffractive effects due to the opening. This is our approach in a nutshell. It is clear that the formalism can be extended to boundaries with many components with different configurations of apertures, edges, etc. There will be many contributory terms to $S_B(\varphi)$, and these would capture the various boundary effects.

This stage-wise functional integration is easily implemented by writing $\phi_L = \phi_{0L} + \eta_L$, where ϕ_{0L} is a specific field configuration in V_L with the boundary value $\phi_{0L} \rightarrow \varphi$ as $x_1 \rightarrow b$. Explicitly, we may take it to be

$$\phi_{0L}(x) = \int_{\partial V_L} \varphi(x') n \cdot \partial_{x'} G_L(x, x'), \quad (3)$$

where $G_L(x, x')$ is the Green's function for \square with Dirichlet boundary conditions, i.e., $\square G_L(x, x') = \delta^{(4)}(x - x')$, $G_L = 0$ if x or $x' \in \partial V_L$. In this equation, $n \cdot \partial$ denotes the derivative normal to the boundary ∂V_L . The functional freedom for the fields in V_L is in η_L , which is taken to vanish at all boundaries (including the apertures), so that the full field ϕ_L has the required boundary behavior, $\phi_L \rightarrow \varphi$. This is consistent with the Dirichlet boundary condition for the Green's function G_L , which is also the propagator for η_L . The functional integration over η_L gives the partition function for the fields in V_L , namely, Z_L , evaluated with the Dirichlet boundary condition, and also Ψ_L , which carries the φ dependence. A similar splitting $\phi_R = \phi_{0R} + \eta_R$ can be used for the right region V_R . The boundary action is then found to be

$$S_B(\varphi) = \frac{1}{2} \int \varphi(x) n \cdot \partial_x n \cdot \partial_{x'} [G_L(x, x') + G_R(x, x')] \varphi(x'), \quad (4)$$

where the integration is over the common interface and where G_R also satisfies Dirichlet boundary conditions.

We see that the value of the field at the interface, namely, φ is what is left free until the last integration. It is easy to impose a vanishing condition on this field wherever there is a material plate at the interface. However, imposing a Neumann condition (or anything more general) is not straightforward. We may use a mode expansion of the fields in V_L in terms of eigenmodes of \square , obeying more general boundary conditions, but then extracting a boundary action (which can easily accommodate apertures) becomes awkward. For example, if we want to use the Neumann condition, we have to use bulk modes which obey this condition for the plate parts of the interface but must have an arbitrary value φ on the openings, left free until the final integration. Finding such modes is a rather involved mathematical problem. We would like a formalism which has the facility of dealing with apertures and diffractive effects as easily as we can do for the Dirichlet case yet accommodates general boundary conditions directly in terms of the boundary action, so that the bulk

modes are easy to construct. In the next section, we will address this question.

Two remarks are useful at this point. Although we used a free scalar field theory to illustrate the setup, it is clear that we can use a similar formalism for an interacting field theory as well. Secondly, if we think of the x_1 direction as time (with a Minkowski signature), the result of doing the functional integral with the specified value of the field, namely, φ at $x_1 = b$, would be a wave functional (for some state) of the field theory. We have emphasized this by using the notation $\Psi_L(\varphi)$. (If we take the left end of the box to be at $x_1 = -\infty$, then we get the ground state wave functional). So what we have is really a Euclidean version of this wave functional setup. As shown in Ref. [4], this wave functional can be obtained directly from the quantum effective action $\Gamma[\chi]$. We solve $(\delta\Gamma_L/\delta\chi) = 0$ subject to the boundary value $\chi \rightarrow \varphi$ as $x_1 \rightarrow b$. Then $\Psi_L = \exp(-\Gamma_L^*)$ where Γ_L^* is given by $\Gamma[\chi]$ evaluated on this solution. (Obviously, similar statements hold for $\tilde{\Psi}_R$ as well.) This gives another way to think of our procedure in a fully interacting quantum field theory.

III. GENERAL BOUNDARY CONDITIONS

We start again with the simple case of a free scalar field and consider the von Neumann theory of self-adjoint extensions [13]. This has recently been rephrased nicely by Asorey, Ibort, and Marmo, so we can use their approach [14]. For the Laplace operator in some region V , we consider the field φ and its normal derivative $\partial_n\varphi$. On the boundary, we consider square integrable functions, i.e., they form a Hilbert space. Thus, the combination $\varphi + i\partial_n\varphi$ (where $\partial_n = n \cdot \partial$ is the normal derivative) may be viewed as an element of this Hilbert space. The most general boundary condition, according to the von Neumann theory, is

$$\begin{aligned} \varphi + i\partial_n\varphi &= U(\varphi - i\partial_n\varphi) \\ (\varphi + i\partial_n\varphi)(x) &= \oint_y U(x,y)(\varphi - i\partial_n\varphi)(y), \end{aligned} \quad (5)$$

where U is a unitary operator on the boundary Hilbert space, namely, on the space of \mathbb{L}^2 -functions on the boundary, of which φ is an element. In the second line of the equation above, we emphasize this by writing $U(x,y)$. Using the operator notation of the first line of Eq. (5), we find

$$\partial_n\varphi = -i\left(\frac{U-1}{U+1}\right)\varphi \equiv -\mathcal{K}\varphi. \quad (6)$$

\mathcal{K} is a Hermitian operator, and the transformation from U to \mathcal{K} is the so-called Cayley transform. There are two interesting limits, $\mathcal{K} \rightarrow 0$ and $\mathcal{K} \rightarrow \infty$. The first one, equivalent to $U = 1$, gives the Neumann boundary condition, and the second one ($U \rightarrow -1$) corresponds to the Dirichlet condition. (For the latter, it is better to divide

Eq. (6) by \mathcal{K} and then take the limit.) These are special points. The case of \mathcal{K} being a constant (proportional to the identity on the Hilbert space) is the Robin condition. The general theorem is that, in the space of \mathcal{K} 's, infinitesimally close to the two limits of Neumann and Dirichlet, the Laplace operator can have negative eigenvalues [14].

The obvious question is whether this can affect our evaluation of the functional integral. To ensure consistency of the variational problem with the boundary conditions, extra surface terms may have to be added to the action, and one can ask if they are sufficient to avoid any pathologies. The answer, in general, is that the negative eigenvalues can lead to pathologies; but for certain types of \mathcal{K} , or range of eigenvalues for the same, we can have a stable situation. This is further commented upon in the discussion section.

There is a different way to look at this problem, which also suggests the solution. Let us go back again to Eq. (2), writing the partition function as

$$\begin{aligned} Z &= [\det(-\square_L)]^{-\frac{1}{2}}[\det(-\square_R)]^{-\frac{1}{2}} \int [d\varphi] \Psi_L(\varphi) \tilde{\Psi}_R(\varphi) \\ &\equiv Z_L Z_R Z_B. \end{aligned} \quad (7)$$

If the partition between V_L and V_R is real, like a plate, maybe with openings, we express that as the vanishing of the field $\phi = \varphi$ in the plate region of the partition but free on the open region. Equivalently, we may say that the plate is represented by $\delta(\varphi)$ where the delta function is only on the plate region. Explicitly, we can expand φ in terms of modes on the open regions of the interface and on the plate parts. The coefficients of the latter vanish by the delta function. The result then agrees with what we did in Ref. [3]. We may interpret this delta function as the operator representing the insertion of the plate, the plate operator. So the final integral, apart from the determinants, looks like

$$Z_B = \int [d\varphi] \Psi_L(\varphi) [\delta_{\text{plate}}(\varphi)] \tilde{\Psi}_R(\varphi). \quad (8)$$

Now consider what happens when we have a normal derivative. The normal derivative acts as a functional derivative on the Ψ 's. This can be seen easily from slicing up the path integral along the direction normal to the interface, the x_1 direction. The Euclidean action then looks like

$$\begin{aligned} S &= S(\{\varphi_i\}) \\ &= \frac{1}{2} \int d^3x^T \left[\frac{(\varphi_N - \varphi_{N-1})^2}{x_N - x_{N-1}} + \frac{(\varphi_{N-1} - \varphi_{N-2})^2}{x_{N-1} - x_{N-2}} + \dots \right. \\ &\quad \left. + (\nabla^T \varphi)^2 \right]. \end{aligned} \quad (9)$$

φ_N is the boundary value φ . The boundary action is given by

$$\Psi(\varphi) \equiv e^{-S_B(\varphi)} = \int \prod_1^{N-1} d\varphi_i \exp(-S(\{\varphi_i\})). \quad (10)$$

Differentiating e^{-S} with respect to φ_N , we find

$$\frac{\delta}{\delta\varphi_N} e^{-S} = -\frac{(\varphi_N - \varphi_{N-1})}{x_N - x_{N-1}} e^{-S} \rightarrow -\partial_n \varphi e^{-S}. \quad (11)$$

So, going back to Eqs. (7) and (8), we see that imposing a boundary condition $\varphi' + \mathcal{K}\varphi = 0$ on the plate is equivalent to imposing

$$\left[-\frac{\delta}{\delta\varphi} + \mathcal{K}\varphi \right] \Psi(\varphi) = 0. \quad (12)$$

To see how this requirement can be obtained, consider the integral

$$I = \int [d\varphi] \exp\left(-\frac{1}{2} \int \varphi \mathcal{K}\varphi\right) \Psi[\varphi]. \quad (13)$$

We can then write

$$\begin{aligned} & \int [d\varphi] \exp\left(-\frac{1}{2} \int \varphi \mathcal{K}\varphi\right) (\partial_n \varphi) \Psi[\varphi] \\ &= \int [d\varphi] \exp\left(-\frac{1}{2} \int \varphi \mathcal{K}\varphi\right) \left(-\frac{\delta}{\delta\varphi}\right) \Psi[\varphi] \\ &= \int [d\varphi] \exp\left(-\frac{1}{2} \int \varphi \mathcal{K}\varphi\right) (-\mathcal{K}\varphi) \Psi[\varphi], \end{aligned} \quad (14)$$

where, in the second line, we have done a partial integration. This equation can be rewritten as

$$\int [d\varphi] \exp\left(-\frac{1}{2} \int \varphi \mathcal{K}\varphi\right) (\partial_n \varphi + \mathcal{K}\varphi) \Psi[\varphi] = 0. \quad (15)$$

Thus, we do obtain the required vanishing of $\partial_n \varphi + \mathcal{K}\varphi$, showing that the operator representing the plate must be taken as $\exp(-\frac{1}{2} \int \varphi \mathcal{K}\varphi)$. The boundary condition applies only for the fields on the plate part of the boundary, so we must restrict the fields in $\int \varphi \mathcal{K}\varphi$ to be only on the plate part. Further, we can choose the boundary conditions independently on the left and right sides of the same plate, which have independent boundary fields φ_L and φ_R , so that the insertion of $\exp(-\frac{1}{2} \int \varphi \mathcal{K}\varphi)$ is to be done separately for each. (This issue did not arise in previous calculations because we used Dirichlet conditions setting $\varphi = 0$ for both sides of the plate.) The fields on the aperture part are the same on both sides of the boundary.

So the calculational algorithm is: Use Dirichlet conditions with left and right regions and fields ϕ_L and ϕ_R . These go to the boundary values φ_L and φ_R . Then calculate

$$\begin{aligned} Z_B &= \int [d\varphi_L d\varphi_R] \delta_{\text{aperture}}(\varphi_L - \varphi_R) \\ &\times \exp\left(-\frac{1}{2} \int_{\text{plate},L} \varphi_L \mathcal{K}_L \varphi_L - \frac{1}{2} \int_{\text{plate},R} \varphi_R \mathcal{K}_R \varphi_R\right) \\ &\times \Psi[\varphi_L] \Psi[\varphi_R]. \end{aligned} \quad (16)$$

We integrate over all fields φ_L and φ_R , but the δ function ensures that the fields are the same on both sides of the aperture and the integrals in the exponents are restricted to

the fields on the plate part of the boundary on the left and right sides. This formulation, at least partially, avoids the problem of potential negative eigenvalues since the bulk determinants leading to Z_L and Z_R are always calculated with Dirichlet conditions.

For the free theory, $\Psi(\varphi)$ is of the form

$$\Psi(\varphi) = \exp\left(-\frac{1}{2} \int \varphi(x) M(x, x') \varphi(x')\right), \quad (17)$$

where $M(x, x') = n \cdot \partial_x n \cdot \partial_{x'} (G_L(x, x') + G_R(x, x'))$. Thus, in this case, the expression (16) for Z_B becomes

$$\begin{aligned} Z_B &= \int [d\varphi_L d\varphi_R] \delta_{\text{aperture}}(\varphi_L - \varphi_R) e^{-S_B} \\ S_B &= \frac{1}{2} \int_{\text{plate},L} \varphi_L \mathcal{K}_L \varphi_L + \frac{1}{2} \int_{\text{plate},R} \varphi_R \mathcal{K}_R \varphi_R \\ &+ \frac{1}{2} \int_{\text{boundary}} \varphi M \varphi. \end{aligned} \quad (18)$$

Once again, the integrals involving \mathcal{K} 's are only over the plate regions of the boundary, while the last integral is over the entire boundary, including apertures. If the boundary has disconnected components, as would be the case for, say, parallel plates, then all such components must be included in Eq. (18). We are now in a position to apply this method of calculation to a specific example.

IV. PLATE AND HALF-PLATE

A. Modes on the boundary and the boundary action

We will consider the arrangement shown in Fig. 2, where we have an infinite plate and a semi-infinite plate parallel to it, separated in the x_1 direction by a distance b . The case when the field has Dirichlet boundary conditions with $\phi = 0$ on the plates was analyzed previously [3]. Here we will consider more general boundary conditions. The bulk contribution from the left and right regions, namely, $Z_L Z_R$, which, in the present case, is also calculated with Dirichlet boundary conditions, is the same as before. For the boundary contributions, first of all, we need the boundary action. The Dirichlet propagator in the left region, between the two plates, is given by




FIG. 2. The arrangement of a Robin plate and a Robin half-plate with values of fields indicated.

$$\begin{aligned}
 G(x, x') &= \int \frac{d^3 k}{(2\pi)^3} e^{ik \cdot (x^T - x'^T)} G_D(\omega, x_1, x'_1) \\
 G_D(\omega, x_1, x'_1) &= -\frac{1}{2\omega} N(\omega) [\theta(x_1 - x'_1) \\
 &\quad \times (e^{\omega(x_1 - x'_1)} + e^{2b\omega} e^{-\omega(x_1 - x'_1)}) \\
 &\quad + \theta(x'_1 - x_1) (e^{-\omega(x_1 - x'_1)} + e^{2b\omega} e^{\omega(x_1 - x'_1)}) \\
 &\quad - (e^{\omega(x_1 + x'_1)} + e^{2b\omega} e^{-\omega(x_1 + x'_1)})], \quad (19)
 \end{aligned}$$

where $\theta(x_1 - x'_1)$ is the step function and

$$N(\omega) = \frac{1}{e^{2b\omega} - 1}. \quad (20)$$

Further, $\omega = \sqrt{k^2} = \sqrt{k_0^2 + k_2^2 + k_3^2}$, $x^T = (x_0, x_2, x_3)$. We will take the open part of the second plate to be in the range $x_2 \geq 0$. Calculating the normal derivatives, we find

$$\begin{aligned}
 S_B &= \frac{1}{2} \int \varphi_I(x) (\omega \coth b\omega + \mathcal{K}_I)_{x,x'} \varphi_I(x') \\
 &\quad + \int \varphi_I(x) (-\omega \operatorname{csch} b\omega)_{x,x'} \varphi_{II}(x') \\
 &\quad + \frac{1}{2} \int \varphi_{II}(x) (\omega \coth b\omega + \mathcal{K}_L)_{x,x'} \varphi_{II}(x') \\
 &\quad + \frac{1}{2} \int \varphi_R(x) (\omega + \mathcal{K}_R)_{x,x'} \varphi_R(x'), \quad (21)
 \end{aligned}$$

where φ_I refers to the field on plate I, at $x_1 = 0$, φ_{II} refers to that on the left side (V_L side) of the interface at $x_1 = b$ (which includes the half-plate and aperture), and φ_R refers to the field again at the same interface, but on the right side (the V_R side).

We will now need to specify the operators \mathcal{K}_I , \mathcal{K}_L , and \mathcal{K}_R . For illustrating the techniques outlined, we will choose a simple Robin boundary condition $\mathcal{K}_I = \kappa_I = \text{constant}$ for the left plate (plate I). The special cases $\kappa_I = 0$ and $\kappa_I \rightarrow \infty$ will correspond to the Neumann and Dirichlet limits for the left plate. To specify \mathcal{K}_L and \mathcal{K}_R , it is useful to separate φ_{II} into modes which have support on the aperture, modes with support on the plate, and the value of the field at the edge of plate II, $x_2 = 0$. This is done by writing

$$\begin{aligned}
 \varphi_{II} &= \int \frac{d^3 k}{(2\pi)^3} e^{ik \cdot x} \left[\int_0^\infty \frac{dp}{\pi} \frac{2p}{p^2 - (k_2 - i\epsilon)^2} \xi(\vec{k}, p) \right. \\
 &\quad \left. - \int_0^\infty \frac{dp}{\pi} \frac{2p}{p^2 - (k_2 + i\epsilon)^2} \phi_{2L}(\vec{k}, p) + \frac{2|\vec{k}|}{k_2^2 + \vec{k}^2} \rho(\vec{k}) \right]. \quad (22)
 \end{aligned}$$

Here $\vec{k} = (k_0, k_3)$. We define the fields

$$\begin{aligned}
 \phi_{2L}(x) &= \int \frac{d^2 k}{(2\pi)^2} \int_0^\infty 2 \frac{dp}{\pi} e^{ik \cdot \vec{x}} \sin(px_2) \phi_{2L}(\vec{k}, p) \\
 \xi(x) &= \int \frac{d^2 k}{(2\pi)^2} \int_0^\infty 2 \frac{dp}{\pi} e^{ik \cdot \vec{x}} \sin(px_2) \xi(\vec{k}, p) \quad (23) \\
 \rho(x) &= \int \frac{d^2 k}{(2\pi)^2} e^{ik \cdot \vec{x}} \rho(\vec{k}) \begin{cases} e^{-|\vec{k}|x_2} & x_2 > 0 \\ e^{|\vec{k}|x_2} & x_2 < 0. \end{cases}
 \end{aligned}$$

The field ξ is the field at the aperture ($x_2 > 0$) but vanishing at the edge $x_2 = 0$; ϕ_{2L} is the field on the plate ($x_2 < 0$), vanishing at $x_2 = 0$; and ρ is essentially the value of the field at $x_2 = 0$, but continued in a very specific way, with no additional functional degrees of freedom, to $x_2 > 0$ and $x_2 < 0$. (While the value of the field at $x_2 = \pm\infty$ will not be relevant for us, the value of the field vanishing at $x_2 = \pm\infty$ will make some of the integrals easier and better defined. The continuation of the field value at the edge, namely, ρ , to other values of x_2 has been done in one particular way which ensures this.)

Carrying out the p integration in Eq. (22) shows that

$$\varphi_{II}(x) = \begin{cases} \xi(x) + \rho(x) & x_2 > 0 \\ \phi_{2L}(x) + \rho(x) & x_2 < 0 \\ \rho(x_0, x_3, x_2 = 0) & x_2 = 0. \end{cases} \quad (24)$$

The amplitudes of the modes in Eq. (22), namely, $\xi(\vec{k}, p)$, $\rho(\vec{k})$, and $\phi_{2L}(\vec{k}, p)$ constitute the functional freedom in the value of the field at the boundary. In a similar way, we can write

$$\varphi_R(x) = \begin{cases} \xi(x) + \rho(x) & x_2 > 0 \\ \phi_{2R}(x) + \rho(x) & x_2 < 0 \\ \rho(x_0, x_3, x_2 = 0) & x_2 = 0, \end{cases} \quad (25)$$

where

$$\phi_{2R} = \int \frac{d^2 k}{(2\pi)^2} \int_0^\infty 2 \frac{dp}{\pi} e^{ik \cdot \vec{x}} \sin(px_2) \phi_{2R}(k). \quad (26)$$

For φ_I , since there is a single plate for all x_2 , we can use a simple mode expansion,

$$\varphi_I = \int \frac{d^3 k}{(2\pi)^3} e^{ik \cdot x} \phi_1(k). \quad (27)$$

The normalization for $\sin(px_2)$ is

$$\int_0^\infty dx_2 \sin(px_2) \sin(qx_2) = \frac{\pi}{2} \delta(p - q). \quad (28)$$

One advantage in parametrizing the fields as in Eq. (22) is that this already takes care of the δ function $\delta_{\text{aperture}}(\varphi_{II} - \varphi_R)$, enforcing equality of fields on the aperture $x_2 \geq 0$.

We may now regard \mathcal{K}_L as an operator on the fields (ϕ_{2L}, ρ) and likewise \mathcal{K}_R as an operator on (ϕ_{2R}, ρ) . While ξ and ϕ_{2L} (and ξ and ϕ_{2R}) are orthogonal to each other, ρ is not orthogonal to ξ or ϕ_{2L} (or ϕ_{2R}). Nevertheless, (ξ, ϕ_{2L}, ρ) and (ξ, ϕ_{2R}, ρ) form a complete

basis for fields at $x_1 = b$. There is clearly an infinity of choices possible for \mathcal{K}_L and \mathcal{K}_R , but, once again, for our analysis, we will make a simple choice. We will take $\mathcal{K}_L = \kappa_L = \text{constant}$ for all the modes ϕ_{2L} and $\mathcal{K}_R = \kappa_R = \text{constant}$ for all the modes ϕ_{2R} . In other words, the operators \mathcal{K}_L and \mathcal{K}_R are diagonal with the same eigenvalue κ_L (respectively, κ_R) for all ϕ_{2L} (respectively, ϕ_{2R}). This is almost like a Robin condition for the plate on the right as well. We say ‘‘almost’’ because the situation with ρ is a bit tricky. It corresponds to the field just at the edge

$x_2 = 0$. In principle, the value of \mathcal{K}_L for ρ can be different from the values of the same operator for ϕ_{2L} , even after we have chosen the latter to be the same for all modes ϕ_{2L} . A similar statement applies to \mathcal{K}_R for ρ versus \mathcal{K}_R for ϕ_{2R} . In addition, we can have nondiagonal terms mixing ρ with ϕ_{2L} and ϕ_{2R} . Thus, even with the simplifying choices we have made, there are many parameters specifying the boundary behavior. Summarizing, the boundary action (18) for this geometry, with the choices we have made, is

$$S_B = \frac{1}{2} \int \varphi_I(x)(\omega \coth b\omega + \kappa_I)_{,x,x'} \varphi_I(x') + \int \varphi_I(x)(-\omega \operatorname{csch} b\omega)_{,x,x'} \varphi_{II}(x') + \frac{1}{2} \int \varphi_{II}(x)(\omega \coth b\omega)_{,x,x'} \varphi_{II}(x') \\ + \frac{1}{2} \int \varphi_R(x)(\omega)_{,x,x'} \varphi_R(x') + \frac{1}{2} \kappa_L \int \phi_{2L} \phi_{2L} + \frac{1}{2} \kappa_R \int \phi_{2R} \phi_{2R} + \frac{1}{2} c_0 \int \rho \rho + c_{2L} \int \phi_{2L} \rho + c_{2R} \int \phi_{2R} \rho. \quad (29)$$

B. Simplification of the boundary action

We are now in a position to evaluate the action in terms of the mode expansions we have given and then carry out the functional integrals. We will only give the final form of the boundary action here, relegating the details to the Appendix.

We will separate the terms into two categories. Because $\rho(\vec{k})$ depends only on two dimensions, it is easier to integrate out the other fields first and leave the ρ integral to the end. Toward this, we shall first simplify the terms involving φ_I , ϕ_{2L} , ϕ_{2R} , and ξ . In this way, we separate out the terms in Eq. (29) as

$$S_B = S_B^{(1)} + S_B^{(2)} \\ S_B^{(1)} = \frac{1}{2} \int \phi_1 \mathcal{M}_1 \phi_1 + \int \phi_1 N_{12L} \phi_{2L} + \int \phi_1 N_{1\xi} \xi + \frac{1}{2} \int \xi \mathcal{M}_\xi \xi + \frac{1}{2} \int \phi_{2L} \mathcal{M}_{2L} \phi_{2L} + \frac{1}{2} \int \phi_{2R} \mathcal{M}_{2R} \phi_{2R} \\ + \int \phi_{2L} Q_{2L\xi} \xi + \int \phi_{2R} N_{2R\xi} \xi \quad (30)$$

$$S_B^{(2)} = \frac{1}{2} \int \rho \mathcal{M}_\rho \rho + \int \phi_1 N_{1\rho} \rho + \int \phi_{2L} Q_{2L\rho} \rho + \int \phi_{2R} Q_{2R\rho} \rho + \int \rho Q_{\rho\xi} \xi. \quad (31)$$

The various coefficient functions, such as \mathcal{M}_1 , N_{12L} , etc., arise naturally from the restriction of the coefficient functions in Eq. (29) to the appropriate modes. We do not give their expressions here; they are given in the Appendix. Integrating over ϕ_1 , ϕ_{2L} , and ϕ_{2R} yields

$$-\log Z_B = \frac{1}{2} \operatorname{Tr} \log (\omega \coth b\omega + \kappa_I) + \frac{1}{2} \operatorname{Tr} \log (\mathbb{M}_L) + \frac{1}{2} \operatorname{Tr} \log (\mathbb{M}_R) - \log Z'_B \quad (32)$$

$$Z'_B = \int [d\xi d\rho] e^{-S'_B}$$

$$S'_B = \frac{1}{2} \int_{\vec{k}, p, q} \xi(-\vec{k}, p) \mathbb{M}_\xi(p, q) \xi(\vec{k}, q) + \int_{\vec{k}, p} \rho(-\vec{k}) N_{\rho\xi}(\vec{k}, p) \xi(\vec{k}, p) + \frac{1}{2} \int_{\vec{k}} \rho(-\vec{k}) \mathbb{M}_\rho \rho(\vec{k}) \\ - \frac{1}{2} \int_{\vec{k}, p, q} \left\{ \left[\int_{p'} N_{2L\xi}(\vec{k}, p, p') \xi(-\vec{k}, p') - \rho(-\vec{k}) N_{2L\rho}(\vec{k}, p) \right] G_{2L}(\vec{k}, p, q) \left[\int_{q'} N_{2L\xi}(\vec{k}, q, q') \xi(\vec{k}, q') - \rho(\vec{k}) N_{2L\rho}(\vec{k}, q) \right] \right\} \\ - \frac{1}{2} \int_{\vec{k}, p, q} \left\{ \left[\int_{p'} N_{2R\xi}(\vec{k}, p, p') \xi(-\vec{k}, p') + \rho(-\vec{k}) N_{2R\rho}(\vec{k}, p) \right] G_{2R}(\vec{k}, p, q) \left[\int_{q'} N_{2R\xi}(\vec{k}, q, q') \xi(\vec{k}, q') + \rho(\vec{k}) N_{2R\rho}(\vec{k}, q) \right] \right\}. \quad (33)$$

The indicated integrations are done with the measures

$$\int_{\vec{k}, p, q} = \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{dq}{\pi}, \quad \int_{p'} = \int_0^\infty \frac{dp'}{\pi}, \quad \text{etc.} \quad (34)$$

The other quantities in Eqs. (32) and (33) are

$$\begin{aligned} \mathbb{M}_L &= 2\pi(H_p + \kappa_L)\delta(p - q) + (\Delta\mathbb{M}_L)_{pq} \\ (\Delta\mathbb{M}_L)_{pq} &= 4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{H_s}{(p^2 - s^2)(q^2 - s^2)} \right. \\ &\quad \left. + \frac{H_p}{(s^2 - p^2)(q^2 - p^2)} + \frac{H_q}{(s^2 - q^2)(p^2 - q^2)} \right] \end{aligned} \quad (35)$$

$$\begin{aligned} \mathbb{M}_R &= 2\pi(\omega_p + \kappa_R)\delta(p - q) + (\Delta\mathbb{M}_R)_{pq} \\ (\Delta\mathbb{M}_R)_{pq} &= 4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{\omega_s}{(p^2 - s^2)(q^2 - s^2)} \right. \\ &\quad \left. + \frac{\omega_p}{(s^2 - p^2)(q^2 - p^2)} + \frac{\omega_q}{(s^2 - q^2)(p^2 - q^2)} \right] \end{aligned} \quad (36)$$

$$\begin{aligned} \mathbb{M}_\xi &= 2\pi(H_p + \omega_p)\delta(p - q) + (\Delta\mathbb{M}_\xi)_{pq} \\ (\Delta\mathbb{M}_\xi)_{pq} &= 4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{(H_s + \omega_s)}{(p^2 - s^2)(q^2 - s^2)} \right. \\ &\quad \left. + \frac{(H_p + \omega_p)}{(s^2 - p^2)(q^2 - p^2)} + \frac{(H_q + \omega_q)}{(s^2 - q^2)(p^2 - q^2)} \right] \end{aligned} \quad (37)$$

$$\mathbb{M}_\rho = 4|\vec{k}|^2 \int_{-\infty}^\infty \frac{ds}{2\pi} \frac{H_s + \omega_s}{\omega_s^4} + \frac{c_0}{2|\vec{k}|}. \quad (38)$$

In these formulas,

$$H_p = \omega_p \left(\frac{\omega_p + \kappa_L \coth b\omega_p}{\omega_p \coth b\omega_p + \kappa_L} \right), \quad \omega_p^2 = \vec{k}^2 + p^2. \quad (39)$$

The quantities G_{2L} and G_{2R} are the Green's functions or inverses of the kernels \mathbb{M}_L and \mathbb{M}_R , respectively. Explicitly, they are of the form

$$\begin{aligned} G_{2L}(p, q) &= \pi\delta(p - q) \frac{1}{2(H_p + \kappa_L)} \\ &\quad - \frac{1}{2(H_p + \kappa_L)} (\Delta\mathbb{M}_L)_{pq} \frac{1}{2(H_q + \kappa_L)} \\ &\quad + \int_0^\infty \frac{dp'}{\pi} \frac{1}{2(H_p + \kappa_L)} (\Delta\mathbb{M}_L)_{pp'} \\ &\quad \times \frac{1}{2(H_{p'} + \kappa_L)} (\Delta\mathbb{M}_L)_{p'q} \frac{1}{2(H_q + \kappa_L)} + \dots \end{aligned} \quad (40)$$

$$\begin{aligned} G_{2R}(p, q) &= \pi\delta(p - q) \frac{1}{2(\omega_p + \kappa_R)} \\ &\quad - \frac{1}{2(\omega_p + \kappa_R)} (\Delta\mathbb{M}_R)_{pq} \frac{1}{2(\omega_q + \kappa_R)} \\ &\quad + \int_0^\infty \frac{dp'}{\pi} \frac{1}{2(\omega_p + \kappa_R)} (\Delta\mathbb{M}_R)_{pp'} \\ &\quad \times \frac{1}{2(\omega_{p'} + \kappa_R)} (\Delta\mathbb{M}_R)_{p'q} \frac{1}{2(\omega_q + \kappa_R)} + \dots \end{aligned} \quad (41)$$

Finally, we also have

$$\begin{aligned} N_{\rho\xi}(\vec{k}, p) &= 4|\vec{k}|p \int_0^\infty \frac{ds}{\pi} \left[\frac{(H_s + \omega_s)}{\omega_s^2} - \frac{(H_p + \omega_p)}{\omega_p^2} \right] \\ &\quad \times \frac{1}{p^2 - s^2} \\ N_{2L\xi}(\vec{k}, p, p') &= -(\Delta\mathbb{M}_L)_{pq} \\ N_{2R\xi}(\vec{k}, p, p') &= -(\Delta\mathbb{M}_R)_{pq} \\ &= \frac{4pp'}{(p'^2 - p^2)} [(p'^2 + \vec{k}^2)I(\vec{k}, p') \\ &\quad - (q^2 + \vec{k}^2)I(\vec{k}, q)] \end{aligned} \quad (42)$$

$$\begin{aligned} N_{2L\rho}(\vec{k}, p) &= 4|\vec{k}|p \left[\int_0^\infty \frac{ds}{\pi} \left(\frac{H_s}{\omega_s^2} - \frac{H_p}{\omega_p^2} \right) \frac{1}{p^2 - s^2} + \frac{c_{2L}}{2|\vec{k}|\omega_p^2} \right] \\ N_{2R\rho}(\vec{k}, p) &= -4|\vec{k}|p \left[I(\vec{k}, p) + \frac{c_{2R}}{2|\vec{k}|\omega_p^2} \right], \end{aligned} \quad (43)$$

where

$$I(\vec{k}, p) = \frac{1}{2\pi p \sqrt{\vec{k}^2 + p^2}} \log \left(\frac{\sqrt{\vec{k}^2 + p^2} + p}{\sqrt{\vec{k}^2 + p^2} - p} \right). \quad (44)$$

The equalities $N_{2L\xi} = -\Delta\mathbb{M}_L$ and $N_{2R\xi} = -\Delta\mathbb{M}_R$ show that $N_{2L\xi}, N_{2R\xi}$ are of the same order in diffractive effect as the ΔM 's, an issue which is relevant when we do the expansions explained below.

C. Evaluation of the Casimir energy

We are now in a position to write down the free energy, which will have several terms corresponding to the contributions from the left and right bulk regions and integrations over $\phi_1, \phi_{2L}, \phi_{2R}$, etc. First, in extracting the free energy, we note that the \vec{k} dependence is the same for all terms and that the overall integration factor representing the trace is $L_0 L_3 (d^2k / (2\pi)^2)$. With $-\log Z_B = L_0 F$, the free energy is given by

$$F = F_{\text{bulk}} + F_I + F_{2L} + F_{2R} + F_\xi + F_\rho. \quad (45)$$

We have added in F_{bulk} , which is the contribution from the bulk determinants (from Z_L, Z_R), calculated with Dirichlet

conditions. The remaining terms in Eq. (45) arise from the boundary action and correspond to the terms on the right-hand side of Eq. (32) or the result of integrating over the terms on the right-hand side of Eq. (33). All these contributions have divergences corresponding to free space with no plates. They can be identified as the $b \rightarrow \infty$ limit of the expressions here. Thus, in the following calculations, the renormalization of all terms will be done by subtracting the $b \rightarrow \infty$ limit.

1. Note on the expansion scheme

The exact calculation of the relevant determinants and the free energy is still very involved, and an expansion scheme is needed to get a good approximation for several of the terms in Eq. (45). In our earlier work [3], we noticed that the relevant operator, say \mathbb{M} , had the structure $\mathbb{M}_{pq} = \mathbb{M}^{(0)}\delta(p-q) + \Delta\mathbb{M}_{pq}$, where $\mathbb{M}^{(0)}$ is diagonal as indicated. It was referred to as a pole term, and $\Delta\mathbb{M}$ was designated a cut term, based on the method of calculation we had used [3]. More appropriately, $\mathbb{M}^{(0)}$ was a ‘‘direct term,’’ giving the geometrical optics approximation while the ‘‘diffractive term’’ $\Delta\mathbb{M}$ captured the effects of diffraction [5]. We then carried out an expansion in powers of the diffractive term, i.e.,

$$\begin{aligned} \text{Tr log}(\mathbb{M}^{(0)} + \Delta\mathbb{M}) &= \text{Tr log} \mathbb{M}^{(0)} + \text{Tr}((\mathbb{M}^{(0)})^{-1}\Delta\mathbb{M}) \\ &\quad - \frac{1}{2} \text{Tr}((\mathbb{M}^{(0)})^{-1}\Delta\mathbb{M}(\mathbb{M}^{(0)})^{-1}\Delta\mathbb{M}) + \dots \end{aligned} \quad (46)$$

Although there is really no parameter controlling this expansion, numerically, the higher-and-higher-order diffractive contributions in Eq. (46) were smaller and smaller, and a sensible truncation was possible.

We propose to do a similar expansion here. The propagators in Eqs. (40) and (41), as we have presented them, naturally show this expansion. Likewise, $N_{2L\xi} = -\Delta\mathbb{M}_L$, $N_{2R\xi} = -\Delta\mathbb{M}_R$ are to be considered as of the first order in the diffractive effect. With this understanding, we can now proceed to the individual terms in Eq. (45).

2. Bulk contribution

The bulk contribution to the free energy is

$$F_{\text{bulk}} = \frac{1}{2}L_3(W_1 + W_2) \int \frac{d^3k}{(2\pi)^3} \log(1 - e^{-2b\omega}), \quad (47)$$

where W_1 is the width of the aperture and W_2 is the width of the half-plate on the right, with $L_2 = W_1 + W_2$.

3. Contribution from fields on the left plate (plate I)

For F_I , which is the contribution from the left plate, we have

$$\begin{aligned} F_I &= \frac{1}{2} \text{Tr log}(\omega_k \coth b\omega_k + \kappa_I) - (b \rightarrow \infty \text{ limit}) \\ &= \frac{1}{2}L_3(W_1 + W_2) \int \frac{d^3k}{(2\pi)^3} \log \left[\frac{\omega_k \coth b\omega_k + \kappa_I}{\omega_k + \kappa_I} \right]. \end{aligned} \quad (48)$$

We have also carried out the renormalization by subtracting the $b \rightarrow \infty$ limit.

4. Contribution from fields on the left side of the half-plate

The term F_{2L} corresponds to $\frac{1}{2} \text{Tr log} \mathbb{M}_L$ and can be simplified as

$$\begin{aligned} F_{2L} &= \frac{1}{2}L_3W_2 \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \log(H_p + \kappa_L) \\ &\quad + \frac{1}{2}L_3 \int \frac{d^2k}{(2\pi)^2} \text{Tr log} \left[1 + \frac{1}{2(H_p + \kappa_L)} \Delta\mathbb{M}_L \right] \\ &\quad - (b \rightarrow \infty \text{ limit}) \\ &= F_{2L}^{(0)} + F_{2L}^{(1)} + F_{2L}^{(2)} + \dots, \end{aligned} \quad (49)$$

where we have expanded the second term in a series in $\Delta\mathbb{M}_L$. In expanding the last term, we get the trace of products of $(1/2(H_p + \kappa_L))\Delta\mathbb{M}_L$. The ‘‘Tr’’ stands for setting the labels $p = q$ for the two end terms and integrating with dp/π . The first few terms in the expansion are

$$F_{2L}^{(0)} = \frac{1}{2}L_3W_2 \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \log \left(\frac{H_p + \kappa_L}{\omega_p + \kappa_L} \right), \quad (50)$$

$$F_{2L}^{(1)} = \frac{L_3}{2} \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{1}{2(H_p + \kappa_L)} (\Delta\mathbb{M}_L)_{pp} - (b \rightarrow \infty \text{ limit}), \quad (51)$$

$$\begin{aligned} F_{2L}^{(2)} &= -\frac{L_3}{4} \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{dq}{\pi} \frac{1}{2(H_p + \kappa_L)} (\Delta\mathbb{M}_L)_{pq} \\ &\quad \times \frac{1}{2(H_q + \kappa_L)} (\Delta\mathbb{M}_L)_{qp} - (b \rightarrow \infty \text{ limit}). \end{aligned} \quad (52)$$

5. Contribution from fields on the right side of the half-plate

The term F_{2R} corresponds to $\frac{1}{2} \text{Tr log} \mathbb{M}_R$; it has no b dependence, and hence it is eliminated by renormalization.

6. Contribution from fields on the aperture

The next term of interest is the contribution from the ξ integration. The $\xi\xi$ terms in Eq. (33) can be collected together as

$$\begin{aligned}
S_{\xi\xi} &= \frac{1}{2} \int_{\vec{k}, p, q} \xi(-\vec{k}, p) \left[\mathbb{M}_\xi(p, q) \right. \\
&\quad - \int_{p', q'} N_{2L\xi}(\vec{k}, p', p) G_{2L}(\vec{k}, p', q') N_{2L\xi}(\vec{k}, q', q) \\
&\quad - \int_{p', q'} N_{2R\xi}(\vec{k}, p', p) G_{2R}(\vec{k}, p', q') \\
&\quad \left. \times N_{2R\xi}(\vec{k}, q', q) \right] \xi(\vec{k}, q) \\
&= \frac{1}{2} \left[\int_{\vec{k}, p} \xi(-\vec{k}, p) 2(H_p + \omega_p) \xi(\vec{k}, p) \right. \\
&\quad \left. + \int_{\vec{k}, p, q} \xi(-\vec{k}, p) (\Delta\tilde{\mathbb{M}}_\xi)_{pq} \xi(\vec{k}, q) \right], \quad (53)
\end{aligned}$$

where

$$\begin{aligned}
(\Delta\tilde{\mathbb{M}}_\xi)_{pq} &= (\Delta\mathbb{M}_\xi)_{pq} \\
&\quad - \int_{p', q'} N_{2L\xi}(\vec{k}, p', p) G_{2L}(\vec{k}, p', q') N_{2L\xi}(\vec{k}, q', q) \\
&\quad - \int_{p', q'} N_{2R\xi}(\vec{k}, p', p) G_{2R}(\vec{k}, p', q') N_{2R\xi}(\vec{k}, q', q) \\
&\equiv (\Delta\tilde{\mathbb{M}}_\xi^{(1)})_{pq} + (\Delta\tilde{\mathbb{M}}_\xi^{(2)})_{pq} + \dots \quad (54)
\end{aligned}$$

$$(\Delta\tilde{\mathbb{M}}_\xi^{(1)})_{pq} = (\Delta\mathbb{M}_\xi)_{pq} \quad (55)$$

$$\begin{aligned}
(\Delta\tilde{\mathbb{M}}_\xi^{(2)})_{pq} &= - \int_{p'} N_{2L\xi}(\vec{k}, p', p) \frac{1}{2(H_{p'} + \kappa_L)} N_{2L\xi}(\vec{k}, p', q) \\
&\quad - \int_{p'} N_{2R\xi}(\vec{k}, p', p) \frac{1}{2(\omega_{p'} + \kappa_R)} N_{2R\xi}(\vec{k}, p', q). \quad (56)
\end{aligned}$$

For the contribution to free energy upon integration over the ξ 's, we find

$$\begin{aligned}
F_\xi &= \frac{1}{2} L_3 W_1 \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \log(H_p + \omega_p) \\
&\quad + \frac{1}{2} L_3 \int \frac{d^2k}{(2\pi)^2} \text{Tr} \log \left(1 + \frac{1}{2(H_p + \omega_p)} (\Delta\tilde{\mathbb{M}}_\xi) \right) \\
&\quad - (b \rightarrow \infty \text{ limit}). \quad (57)
\end{aligned}$$

We can simplify this further by noting that

$$\left[\frac{H_p + \omega_p}{(H_p + \omega_p)_{b \rightarrow \infty}} \right] = \left(\frac{1}{1 - e^{-2b\omega_p}} \right) \left(\frac{\omega_p + \kappa_I}{\omega_p \coth b\omega_p + \kappa_I} \right). \quad (58)$$

The term F_ξ thus splits up as

$$\begin{aligned}
F_\xi &= F_\xi^{(0)} + F_\xi^{(1)} + F_\xi^{(2)} + \dots \\
F_\xi^{(0)} &= -\frac{1}{2} L_3 W_1 \int \frac{d^3k}{(2\pi)^3} \log(1 - e^{-2b\omega_k}) \\
&\quad - \frac{1}{2} L_3 W_1 \int \frac{d^3k}{(2\pi)^3} \log \left[\frac{\omega_k \coth b\omega_k + \kappa_I}{\omega_k + \kappa_I} \right], \quad (59)
\end{aligned}$$

$$\begin{aligned}
F_\xi^{(1)} &= \frac{L_3}{2} \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{1}{2(H_p + \omega_p)} (\Delta\mathbb{M}_\xi)_{pp} \\
&\quad - (b \rightarrow \infty \text{ limit}), \quad (60)
\end{aligned}$$

$$\begin{aligned}
F_\xi^{(2)} &= -\frac{L_3}{4} \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{dq}{\pi} \left[\frac{1}{2(H_p + \omega_p)} (\Delta\mathbb{M}_\xi)_{pq} \right. \\
&\quad \left. \times \frac{1}{2(H_p + \omega_p)} (\Delta\mathbb{M})_{qp} \right] \\
&\quad + \frac{L_3}{2} \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{1}{2(H_p + \omega_p)} (\Delta\tilde{\mathbb{M}}_\xi^{(2)})_{pp} \\
&\quad - (b \rightarrow \infty \text{ limit}). \quad (61)
\end{aligned}$$

Notice that the terms in Eq. (59) cancel against similar terms in F_{bulk} and F_I .

7. Contribution from the field at the edge (field ρ)

The contributions we have evaluated so far correspond to the fields ϕ_{2L} , ϕ_{2R} , and ξ , which vanish at the edge of the half-plate. If the parameter $c_0 \rightarrow \infty$, these are the only terms we have. For finite values of c_0 , we have the contribution from ρ as well. To evaluate this term, we need the kernel for the $\rho\rho$ term in the boundary action. In addition to the $\rho\rho$ term manifestly displayed in Eq. (33), there will be additional terms from the integration over ξ because of the ξ - ρ terms in Eq. (33). For this, first define the propagator for ξ given by Eq. (53),

$$\begin{aligned}
G_\xi(\vec{k}) &= \pi \delta(p, q) \frac{1}{2(H_p + \omega_p)} \\
&\quad - \frac{1}{2(H_p + \omega_p)} (\Delta\tilde{\mathbb{M}}_\xi)_{pq} \frac{1}{2(H_q + \omega_q)} \\
&\quad + \int_0^\infty \frac{dp'}{\pi} \frac{1}{2(H_p + \omega_p)} (\Delta\tilde{\mathbb{M}}_\xi)_{pp'} \\
&\quad \times \frac{1}{2(H_{p'} + \omega_{p'})} (\Delta\tilde{\mathbb{M}}_\xi)_{p'q} \frac{1}{2(H_q + \omega_q)} + \dots \quad (62)
\end{aligned}$$

The integration over the ξ 's thus leads to the ρ -dependent terms,

$$S_\rho = \frac{1}{2} \int_{\vec{k}} \rho(-\vec{k}) [\mathbb{M}_\rho(\vec{k}) + \Delta\mathbb{M}_\rho] \rho(\vec{k}), \quad (63)$$

where

$$\begin{aligned}
 \Delta\mathbb{M}_\rho = & - \int_{p,q} N_{2L\rho}(\vec{k}, p) G_{2L}(\vec{k}, p, q) N_{2L\rho}(\vec{k}, q) \\
 & - \int_{p,q} N_{2R\rho}(\vec{k}, p) G_{2R}(\vec{k}, p, q) N_{2R\rho}(\vec{k}, q) \\
 & - \int_{p,q} \tilde{N}_{\rho\xi}(-\vec{k}, p) G_\xi(\vec{k}, p, q) \tilde{N}_{\rho\xi}(\vec{k}, q) \quad (64)
 \end{aligned}$$

$$\begin{aligned}
 \tilde{N}_{\rho\xi} = & N_{\rho\xi} - \int_{p',q'} N_{2L\xi}(\vec{k}, p', p) G_{2L}(\vec{k}, p', q') N_{2L\rho}(\vec{k}, q') \\
 & - \int_{p',q'} N_{2R\xi}(\vec{k}, p', p) G_{2R}(\vec{k}, p', q') N_{2R\rho}(\vec{k}, q'). \quad (65)
 \end{aligned}$$

The quantity $\Delta\mathbb{M}_\rho$ naturally has an expansion which follows from the expansion of the propagators G_{2L} , G_{2R} , and G_ξ . Notice that we start off with at least two powers of $N_{2L\xi}$, $N_{2R\xi}$, or $N_{\rho\xi}$, so that there is no first-order term. The expansion for $\Delta\mathbb{M}_\rho$ is thus

$$\begin{aligned}
 \Delta\mathbb{M}_\rho = & \Delta\mathbb{M}_\rho^{(2)} + \Delta\mathbb{M}_\rho^{(3)} + \dots \\
 \Delta\mathbb{M}_\rho^{(2)} = & - \int_p N_{2L\rho}(\vec{k}, p) \frac{1}{2(H_p + \kappa_L)} N_{2L\rho}(\vec{k}, p) \\
 & - \int_p N_{2R\rho}(\vec{k}, p) \frac{1}{2(\omega_p + \kappa_R)} N_{2R\rho}(\vec{k}, p) \\
 & - \int_p N_{\rho\xi}(-\vec{k}, p) \frac{1}{2(H_p + \omega_p)} N_{\rho\xi}(\vec{k}, p). \quad (66)
 \end{aligned}$$

All terms in these expressions depend on \vec{k} ; we have not indicated this explicitly in the arguments of various functions to avoid too much clutter. Once the integrations over p , q , etc., are done, $\mathbb{M}_\rho + \Delta\mathbb{M}_\rho$ is only a function of \vec{k} , so that the free energy resulting from Eq. (63) can be written as

$$F_\rho = \frac{L_3}{2} \int \frac{d^2k}{(2\pi)^2} \log(\mathbb{M}_\rho + \Delta\mathbb{M}_\rho). \quad (67)$$

Thus, approximations to this term will be of the form

$$\begin{aligned}
 F_\rho^{(0)} = & \frac{L_3}{2} \int \frac{d^2k}{(2\pi)^2} \log(\mathbb{M}_\rho) - (b \rightarrow \infty \text{ limit}) \\
 F_\rho^{(0)} + F_\rho^{(2)} = & \frac{L_3}{2} \int \frac{d^2k}{(2\pi)^2} \log(\mathbb{M}_\rho + \Delta\mathbb{M}_\rho^{(2)}) \\
 & - (b \rightarrow \infty \text{ limit}). \quad (68)
 \end{aligned}$$

D. Numerical estimates

The terms proportional to W_1 , the width of the aperture, cancel out between Eqs. (47), (48), and (59). Thus, we are left with bulk terms which only involve the facing area $L_3 W_2$ between the two plates. This is given by part of F_{bulk} in Eq. (47), part of F_I in Eq. (48), and $F_{2L}^{(0)}$ in Eq. (50) and is equal to

$$\begin{aligned}
 F_{\text{area}} = & \frac{L_3 W_2}{2} \int \frac{d^3k}{(2\pi)^3} \left[\log(1 - e^{-2b\omega_k}) \right. \\
 & \left. + \log\left(\frac{\omega_k \coth b\omega_k + \kappa_I}{\omega_k + \kappa_I}\right) + \log\left(\frac{H_k + \kappa_L}{\omega_k + \kappa_L}\right) \right]. \quad (69)
 \end{aligned}$$

There are also many edge-dependent terms to consider. The fields ξ , ϕ_{2L} vanish at the edge $x_2 = 0$. For an aperture and a half-plate of finite widths W_1 and W_2 , we should use discrete modes to evaluate the free energy, with a sum replacing the integral over p or k_2 . In the limit of large W_1 , W_2 , the sum over the discrete modes can be approximated by an integral, as in an Euler-Maclaurin summation formula, which gives the area-dependent term in Eq. (69). But there is also a subdominant edge-dependent term in the summation formula [3]; we will denote this term by $F_{\text{edge}}^{(0)}$.

The remaining edge-dependent terms in the energy are given by $F_{2L}^{(1)} + F_{2L}^{(2)} + \dots$ from Eqs. (51) and (52), etc.; $F_\xi^{(1)} + F_\xi^{(2)} + \dots$ from Eqs. (60) and (61), etc.; and F_ρ from Eq. (68). All these quantities depend on the parameters κ_I , κ_L , κ_R , c_0 , c_{2L} , and c_{2R} . We will now proceed to the numerical estimate of these integrals for some choices of these parameters.

1. Area-dependent contribution

The area-dependent part of the Casimir energy (69), from scaling out b , is of the form

TABLE I. The area-dependent contributions $\mathcal{E}_{\text{area}}$ for different boundary conditions.

m	0	1	2	3	4	5
$\mathcal{E}_{\text{RD}}(\infty, m) = \mathcal{E}_{\text{DR}}(m, \infty)$	0.875	0.092	-0.212	-0.381	-0.489	-0.564
$\mathcal{E}_{\text{RR}}(m, m)$	-1	-0.089	0.115	-0.200	-0.283	-0.354
m	6	7	8	9	10	∞
$\mathcal{E}_{\text{RD}}(\infty, m) = \mathcal{E}_{\text{DR}}(m, \infty)$	-0.620	-0.663	-0.698	-0.725	-0.749	-1
$\mathcal{E}_{\text{RR}}(m, m)$	-0.414	-0.464	-0.507	-0.544	-0.575	-1




FIG. 3 (color online). The area-dependent contribution to energy, $\mathcal{E}_{\text{area}}$, for different boundary conditions, as a function of m , showing an asymptotic approach to the Dirichlet-Dirichlet case.

$$F_{\text{area}} = \frac{L_3 W_2}{b^3} \frac{\pi^2}{1440} \mathcal{E}_{\text{area}}(b\kappa_I, b\kappa_L). \quad (70)$$

We will consider Dirichlet condition for the left plate ($\kappa_I \rightarrow \infty$), Robin condition for the right (with finite $b\kappa_L = m$)—or the other way ($\kappa_L \rightarrow \infty, b\kappa_I = m$ finite)—and then the Robin condition for both (with

$b\kappa_I = b\kappa_L = m$). The values for these cases are given in Table I. The Dirichlet-Robin and Robin-Dirichlet cases give the same area-dependent term for the energy. The Robin-Robin case with $m = 0$ corresponds to the Neumann condition on both plates. The limit of $m \rightarrow \infty$ should correspond to the case with Dirichlet conditions on both plates, which is, of course, well known. As expected, and as Fig. 3 shows, the values found here approach that value ($\mathcal{E} = -1$) asymptotically.

2. Edge contributions (without ρ)

The edge-dependent contributions from F_{2L} and F_{ξ} (including $F_{\text{edge}}^{(0)}$) are of the form

$$F_{\text{edge}} = \frac{L_3}{b^2} \mathcal{E}_{\text{edge}}(b\kappa_I, b\kappa_L, b\kappa_R). \quad (71)$$

We consider the cases of Robin-Dirichlet ($b\kappa_I = m = \text{finite value}, \kappa_L, \kappa_R \rightarrow \infty$), Dirichlet-Robin ($\kappa_I \rightarrow \infty, b\kappa_L = b\kappa_R = m = \text{finite value}$), and Robin-Robin ($b\kappa_I = b\kappa_L = b\kappa_R = m = \text{finite value}$). The values $\mathcal{E}_{\text{edge}}$ for a range of m are given in Tables II, III, and IV.

TABLE II. The edge contribution to the Casimir energy, Robin-Dirichlet case, without F_{ρ} .

m	0	1	2	3	4	5
$\mathcal{E}_{\text{edge, RD}}^0(m, \infty, \infty)$	0.00448	-0.00079	-0.00241	-0.00325	-0.00377	-0.00412
$\mathcal{E}_{\text{edge, RD}}^1(m, \infty, \infty)$	-0.00030	0.00005	0.00016	0.00022	0.00025	0.00028
$\mathcal{E}_{\text{edge, RD}}^2(m, \infty, \infty)$	0.00180	0.00121	0.00100	0.00088	0.00081	0.00075
$\mathcal{E}_{\text{edge, RD}}^{\text{total}}(m, \infty, \infty)$	0.00249	-0.00033	-0.00112	-0.00152	-0.00176	-0.00193
m	6	7	8	9	10	∞
$\mathcal{E}_{\text{edge, RD}}^0(\infty, m, m)$	-0.00439	-0.00456	-0.00471	-0.00483	-0.00493	-0.00598
$\mathcal{E}_{\text{edge, RD}}^1(\infty, m, m)$	0.00203	0.00212	0.00219	0.00224	0.00229	0.00277
$\mathcal{E}_{\text{edge, RD}}^2(\infty, m, m)$	0.00029	0.00031	0.00032	0.00032	0.00033	0.00040
$\mathcal{E}_{\text{edge, RD}}^{\text{total}}(\infty, m, m)$	-0.00205	-0.00214	-0.00221	-0.00227	-0.00232	-0.00280

TABLE III. The edge contribution to the Casimir energy, Dirichlet-Robin case, without F_{ρ} .

m	0	1	2	3	4	5
$\mathcal{E}_{\text{edge, DR}}^0(\infty, m, m)$	-0.01644	-0.01117	-0.00954	-0.00871	-0.00819	-0.00784
$\mathcal{E}_{\text{edge, DR}}^1(\infty, m, m)$	0.00790	0.00594	0.00516	0.00471	0.00441	0.00419
$\mathcal{E}_{\text{edge, DR}}^2(\infty, m, m)$	0.00180	0.00121	0.00100	0.00088	0.00081	0.00075
$\mathcal{E}_{\text{edge, DR}}^{\text{total}}(\infty, m, m)$	-0.00675	-0.00402	-0.00338	-0.00312	-0.00298	-0.00290
m	6	7	8	9	10	∞
$\mathcal{E}_{\text{edge, DR}}^0(\infty, m, m)$	-0.00759	-0.00740	-0.00725	-0.00712	-0.00702	-0.00598
$\mathcal{E}_{\text{edge, DR}}^1(\infty, m, m)$	0.00403	0.00390	0.00379	0.00371	0.00364	0.00277
$\mathcal{E}_{\text{edge, DR}}^2(\infty, m, m)$	0.00071	0.00068	0.00065	0.00063	0.00061	0.00040
$\mathcal{E}_{\text{edge, DR}}^{\text{total}}(\infty, m, m)$	-0.00285	-0.00282	-0.00280	-0.00278	-0.00277	-0.00280

TABLE IV. The edge contribution to the Casimir energy, Robin-Robin case, without F_ρ .

m	0	1	2	3	4	5
$\mathcal{E}_{\text{edge,RR}}^0(m, m, m)$	0.01495	-0.00084	-0.00340	-0.00441	-0.00492	-0.00521
$\mathcal{E}_{\text{edge,RR}}^1(m, m, m)$	-0.00515	0.00058	0.00187	0.00238	0.00263	0.00277
$\mathcal{E}_{\text{edge,RR}}^2(m, m, m)$	-0.00159	0.00006	0.00034	0.00043	0.00047	0.00049
$\mathcal{E}_{\text{edge,RR}}^{\text{total}}(m, m, m)$	0.00820	-0.00020	-0.00120	-0.00160	-0.00182	-0.00196
m	6	7	8	9	10	∞
$\mathcal{E}_{\text{edge,RR}}^0(m, m, m)$	-0.00540	-0.00552	-0.00561	-0.00568	-0.00573	-0.00598
$\mathcal{E}_{\text{edge,RR}}^1(m, m, m)$	0.00285	0.00289	0.00292	0.00294	0.00295	0.00277
$\mathcal{E}_{\text{edge,RR}}^2(m, m, m)$	0.00049	0.00050	0.00050	0.00050	0.00050	0.00040
$\mathcal{E}_{\text{edge,RR}}^{\text{total}}(m, m, m)$	-0.00206	-0.00213	-0.00220	-0.00224	-0.00228	-0.00280

We have calculated these to the second order in the expansion of the $\text{Tr} \log$ terms, as in Eqs. (50)–(52) and (59)–(61). These do not include the contribution from the ρ integration, which will be given separately. We also graphically display the leading-order term, the first-order term, the second-order term, and the total edge contribution to this order for the three cases in Figs. 4–6. The dashed line is




FIG. 4 (color online). The edge-dependent contributions from F_{2L} and F_ξ for the Robin-Dirichlet case, $\mathcal{E}_{\text{edge,DR}}(m, \infty, \infty)$.




FIG. 5 (color online). The edge-dependent contributions from F_{2L} and F_ξ for the Dirichlet-Robin case, $\mathcal{E}_{\text{edge,DR}}(\infty, m, m)$.

the total contribution, up to second order, and the dotted line is the Dirichlet-Dirichlet limit.

3. Edge contributions from ρ

The edge contribution from ρ is zero for the case of $c_0 \rightarrow \infty$. Thus, for the case of Robin conditions (or anything else) on the left plate and Dirichlet conditions on the half-plate, the results given so far suffice. We will consider the inclusion of the contribution from ρ for the Dirichlet-Robin case ($\kappa_L \rightarrow \infty$). For simplicity, we shall consider the case $c_{2L} = c_{2R} = 0$. We will need to specify c_0 as well. We take it to be $c_0 = (\kappa_L + \kappa_R)$, since this is what would be naturally considered if we take ρ to be part of ϕ_{2L} and ϕ_{2R} . (There is no compelling reason for this choice; it is one case worth considering and easy enough to calculate.) Thus, the contribution from ρ is of the form

$$F_\rho = \frac{L_3}{b^2} \mathcal{E}_\rho(b\kappa_L, b\kappa_R, bc_0). \quad (72)$$

We will calculate this to the second order as well, which means that we will evaluate the expression in Eq. (68). These values, for a range of parameters, are given in




FIG. 6 (color online). The edge-dependent contributions from F_{2L} and F_ξ for the Robin-Robin case, $\mathcal{E}_{\text{edge,RR}}(m, m, m)$.

TABLE V. The edge contribution from ρ to the Casimir energy.

m	0	1	2	3	4	5
$\mathcal{E}_{\rho,DR}(\infty, m, m, 2m)$	0.00336	0.00356	0.00311	0.00271	0.00239	0.00214
$\mathcal{E}_{\rho,RR}(m, m, m, 2m)$	-0.00553	0.00185	0.00102	0.00127	0.00133	0.00132
m	6	7	8	9	10	∞
$\mathcal{E}_{\rho,DR}(\infty, m, m, 2m)$	0.00193	0.00177	0.00162	0.00150	0.00140	0
$\mathcal{E}_{\rho,RR}(m, m, m, 2m)$	0.00128	0.00123	0.00118	0.00113	0.00108	0

Table V. We also show the contribution from ρ as compared to the other edge contributions as a function of the parameter m in graphs Figs. 7 and 8.

V. DISCUSSION

We have generalized our previous formulation for the calculation of diffractive effects in Casimir energy in the case of boundaries with edges and apertures to include general boundary conditions. As a specific example, we have analyzed the geometry of two parallel plates, one of which is a semi-infinite, so that there are edge effects and diffraction. We have considered a variety of boundary




FIG. 7 (color online). The edge-dependent contributions, including ρ , for the Dirichlet-Robin case, $\mathcal{E}_{\text{edge},DR}(\infty, m, m, 2m)$.




FIG. 8 (color online). The edge-dependent contributions, including ρ for the Robin-Robin case, $\mathcal{E}_{\text{edge},RR}(m, m, m, 2m)$.

conditions, and almost all the results are new, not calculated previously by any method. There are a few special cases for which results using other methods are known, and a comparison with the literature is possible.

The first case is where we have two full parallel plates. The Casimir energy for a massless scalar field subject to Robin boundary conditions on two infinitely long parallel plates has been analyzed before in Refs. [11,12]. Of particular interest is Ref. [12], where a boundary term was added to impose Robin conditions at the level of Green's functions. The idea is similar to ours, although our argument for such a term, which is based on the wave functional interpretation, is different and somewhat more general. If we ignore the diffractive contributions in our calculation and replace the facing area of the plates by the full area, our results are in agreement with Refs. [11,12]. (The graph in Ref. [12] shows \mathcal{E}/m^3 rather than \mathcal{E} as in our graph.)

The MIT group has analyzed, using methods of scattering theory, the edge contribution to the Casimir energy in a number of different geometries of boundaries with edges [9,10]. One particular case is exactly the geometry we have but restricted to Dirichlet-Dirichlet (DD) and Neumann-Neumann (NN) conditions. Our results for the DD, Neumann-Dirichlet (ND), Dirichlet-Neumann (DN), and NN cases are summarized in Table VI. These values for the DD and NN cases are in good agreement with the results in Refs. [7,9,10]. In particular, the numerical values obtained from the expressions given in Ref. [10] are $\mathcal{E}_{DD}^{\text{edge}} = -2.63 \times 10^{-3}$ and $\mathcal{E}_{NN}^{\text{edge}} = 2.97 \times 10^{-3}$. In our previous paper [3], we calculated the edge contribution up to the fifth order in diffraction, $\mathcal{E}_{DD}^{\text{edge}} = -2.68 \times 10^{-3}$, and the agreement with Ref. [10] is even better than is apparent from Table VI. In making this comparison, it is important to keep in mind that the Neumann boundary condition corresponds to integrating over all boundary fields, including ρ (with $\kappa_I = \kappa_L = \kappa_R = c_0 = 0$); hence, the contribution from ρ must also be taken into account in comparing the values. We also note that the DD case was initially treated using numerical worldline methods in Ref. [7]. The value obtained was $\mathcal{E}_{DD}^{\text{edge}} = -2.62 \times 10^{-3}$; again, the later calculations [3,9,10] are consistent with this result.

Another interesting result which emerges from our calculations is that the diffractive effects are always of the opposite sign to the nondiffractive (and leading) contribution.

TABLE VI. Summary for DD, ND, DN, and NN cases.

	DD	ND	DN	NN
$\mathcal{E}_{\text{area}}$	-1	$\frac{7}{8}$	$\frac{7}{8}$	-1
$\mathcal{E}_{\text{edge}}^0$	$-\frac{\xi(3)}{64\pi} = -0.00598$	$\frac{3}{4} \frac{\xi(3)}{64\pi} = 0.00448$	$-\frac{11}{4} \frac{\xi(3)}{64\pi} = -0.01645$	$\frac{10}{4} \frac{\xi(3)}{64\pi} = 0.01495$
$\mathcal{E}_{\text{edge}}^1$	0.00277	-0.00170	0.00790	-0.00515
$\mathcal{E}_{\text{edge}}^2$	0.00040	-0.00030	0.00180	-0.00060
$\mathcal{E}_{\text{edge}}^{\text{total}}$	-0.00280	0.0048	-0.00675	0.00820
$\mathcal{E}_{\text{edge},\rho}$	0	0	0.00336	-0.00553
$\mathcal{E}_{\text{edge}}$ (Total)	-0.00280	0.00249	-0.00339	0.00267

This had been noticed already in our previous work [3,5,6], where we considered Dirichlet-Dirichlet boundary conditions. This continues to hold with the more general boundary conditions discussed here, as is strikingly clear from the graphs for the edge contributions. There is presumably a general reason for this; it is a point worth exploring.

Finally, a short remark on the issue of negative eigenvalues: The bulk determinants in our approach are calculated with Dirichlet boundary conditions. There are no negative eigenvalues to worry about for this calculation. However, this issue is not totally eliminated; it has a lingering effect on the boundary action. Notice that the signs of κ_I , κ_L , etc., are important. These parameters occur in the boundary action, and negative values for these can

lead to instabilities. For example, the propagators in Eqs. (40) and (41) clearly display this possibility. We have to conclude that acceptable boundary conditions which encode the boundary effects of real material plates must be such that the operators \mathcal{K} have positive eigenvalues. One may still ask the question whether there is any meaningful physical interpretation for the mathematically acceptable case of negative eigenvalues. This will be discussed in a separate paper [15].

ACKNOWLEDGMENTS

This research was supported in part by NSF Grants No. PHY-1068172 and No. PHY-1213380 and by PSC-CUNY grants.

APPENDIX

In this Appendix, we work out some of the simplification of the terms in the boundary action. Using the mode expansions in Eqs. (24)–(27), we find the following:

$\varphi_I \varphi_I$ term

$$\frac{1}{2} \int \varphi_I(\omega \coth b\omega + \kappa_I) \varphi_I = \frac{1}{2} \int \frac{d^3k}{(2\pi)^3} \phi_1(-k) \mathcal{M}_1(k) \phi_1(k)$$

$$\mathcal{M}_1(k) = (\omega_k \coth b\omega_k + \kappa_I) \quad (\text{A1})$$

$\varphi_I \phi_{2L}$ term

$$\int \varphi_I(-\omega \operatorname{csch} b\omega) \phi_{2L} = \int \frac{d^3k}{(2\pi)^3} \frac{dp}{\pi} \phi_1(-k) N_{12L}(k, p) \phi_{2L}(\vec{k}, p)$$

$$N_{12L}(k, p) = \frac{2p}{p^2 - (k_2 + i\epsilon)^2} (\omega_k \operatorname{csch} b\omega_k) \quad (\text{A2})$$

$\varphi_I \xi$ term

$$\int \varphi_I(-\omega \operatorname{csch} b\omega) \xi = \int \frac{d^3k}{(2\pi)^3} \frac{dq}{\pi} \phi_1(-k) N_{1\xi}(k, q) \xi(\vec{k}, q)$$

$$N_{1\xi}(k, q) = \frac{2q}{q^2 - (k_2 - i\epsilon)^2} (-\omega_k \operatorname{csch} b\omega_k) \quad (\text{A3})$$

$\xi\xi$ term

$$\begin{aligned} \frac{1}{2} \int \xi(\omega + \omega \coth b\omega) \xi &= \frac{1}{2} \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{dq}{\pi} \xi(-\vec{k}, p) \mathcal{M}_\xi(\vec{k}, p, q) \xi(\vec{k}, q) \\ \mathcal{M}_\xi(\vec{k}, p, q) &= \int_{-\infty}^{\infty} \frac{dk_2}{2\pi} \frac{4pq}{[p^2 - (k_2 - i\epsilon)^2][q^2 - (k_2 + i\epsilon)^2]} (\omega_k + \omega_k \coth b\omega_k) \end{aligned} \quad (\text{A4})$$

$\phi_{2L}\phi_{2L}$ term

$$\begin{aligned} \frac{1}{2} \int \phi_{2L}(\omega \coth b\omega + \kappa_L) \phi_{2L} &= \frac{1}{2} \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{dq}{\pi} \phi_{2L}(-\vec{k}, p) \mathcal{M}_{2L}(\vec{k}, p, q) \phi_{2L}(\vec{k}, q) \\ \mathcal{M}_{2L}(\vec{k}, p, q) &= \int_{-\infty}^{\infty} \frac{dk_2}{2\pi} \frac{4pq}{[p^2 - (k_2 - i\epsilon)^2][q^2 - (k_2 + i\epsilon)^2]} (\omega_k \coth b\omega_k + \kappa_L) \end{aligned} \quad (\text{A5})$$

$\phi_{2R}\phi_{2R}$ term

$$\begin{aligned} \frac{1}{2} \int \phi_{2R}(\omega + \kappa_R) \phi_{2R} &= \frac{1}{2} \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{dq}{\pi} \phi_{2R}(-\vec{k}, p) \mathcal{M}_{2R}(\vec{k}, p, q) \phi_{2R}(\vec{k}, q) \\ \mathcal{M}_{2R}(\vec{k}, p, q) &= \int_{-\infty}^{\infty} \frac{dk_2}{2\pi} \frac{4pq}{[p^2 - (k_2 - i\epsilon)^2][q^2 - (k_2 + i\epsilon)^2]} (\omega_k + \kappa_R) \end{aligned} \quad (\text{A6})$$

$\xi\phi_{2L}$ term

$$\begin{aligned} \int \xi(\omega \coth b\omega) \phi_{2L} &= \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{dq}{\pi} \xi(-\vec{k}, p) \mathcal{Q}_{2L\xi}(\vec{k}, p, q) \phi_{2L}(\vec{k}, q) \\ \mathcal{Q}_{2L\xi}(\vec{k}, p, q) &= - \int_{-\infty}^{\infty} \frac{dk_2}{2\pi} \frac{4pq}{[p^2 - (k_2 - i\epsilon)^2][q^2 - (k_2 - i\epsilon)^2]} (\omega_k \coth b\omega_k) \end{aligned} \quad (\text{A7})$$

$\xi\phi_{2R}$ term

$$\begin{aligned} \int \xi(\omega) \phi_{2R} &= \int \frac{d^2k}{(2\pi)^2} \frac{dp}{\pi} \frac{dq}{\pi} \xi(-\vec{k}, p) \mathcal{N}_{2R\xi}(\vec{k}, p, q) \phi_{2R}(\vec{k}, q) \\ \mathcal{N}_{2R\xi}(\vec{k}, p, q) &= - \int_{-\infty}^{\infty} \frac{dk_2}{2\pi} \frac{4pq}{[p^2 - (k_2 - i\epsilon)^2][q^2 - (k_2 - i\epsilon)^2]} \omega_k. \end{aligned} \quad (\text{A8})$$

Before taking up the ρ -dependent terms, we will consider some simplification of the expressions so far. We start with \mathcal{M}_ξ and use the integral representation

$$\begin{aligned} \omega + \omega \coth b\omega &= \int_{-\infty}^{\infty} \frac{d\lambda}{\pi} \frac{\omega^2}{\lambda^2 + \omega^2} f(\lambda), \\ f(\lambda) &= \frac{2}{1 - e^{-2ib(\lambda - i\epsilon)}}. \end{aligned} \quad (\text{A9})$$

The contour for the λ integration is to be completed in the lower half-plane. Using this result in Eq. (A4), we can carry out the k_2 integration. Then, we do the λ integral to obtain

$$\mathcal{M}_\xi = 2\pi f(-i\omega_p) \omega_p \delta(p, q) + 4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{f(-i\omega_s) \omega_s}{(p^2 - s^2)(q^2 - s^2)} + \frac{f(-i\omega_p) \omega_p}{(s^2 - p^2)(q^2 - p^2)} + \frac{f(-i\omega_q) \omega_q}{(s^2 - q^2)(p^2 - q^2)} \right], \quad (\text{A10})$$

where $\omega_p = \sqrt{k_0^2 + k_3^2 + p^2}$, etc., and

$$f(-i\omega) \omega = \frac{2\omega}{1 - e^{-2b\omega}}. \quad (\text{A11})$$

The integration over ϕ_1 produces the additional term $-N_{1\xi}(-k)(\mathcal{M}_1)^{-1}N_{1\xi}$, which is written out as

$$\begin{aligned}
 -N_{1\xi}(\mathcal{M}_1)^{-1}N_{1\xi} &= -\int \frac{dk_2}{2\pi} \frac{4pq}{[p^2 - (k_2 - i\epsilon)^2][q^2 - (k_2 + i\epsilon)^2]} \frac{\omega^2}{(\sinh b\omega)^2(\omega \coth b\omega + \kappa_I)} \\
 &= -\int \frac{d\lambda}{\pi} \frac{\omega^2}{\lambda^2 + \omega^2} h(\lambda) \frac{4pq}{[p^2 - (k_2 - i\epsilon)^2][q^2 - (k_2 + i\epsilon)^2]} \\
 h(\lambda) &= \frac{i\lambda}{[\kappa_I(\sin b(\lambda - i\epsilon))^2 + (\lambda - i\epsilon) \sin b(\lambda - i\epsilon) \cos b(\lambda - i\epsilon)]}.
 \end{aligned} \tag{A12}$$

By carrying out the integrations as we did to obtain Eq. (A10), we find

$$\begin{aligned}
 -N_{1\xi}(\mathcal{M}_1)^{-1}N_{1\xi} &= -2\pi h(-iK_p)K_p\delta(p - q) \\
 &\quad - 4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{h(-iK_s)K_s}{(p^2 - s^2)(q^2 - s^2)} + \frac{h(-iK_p)K_p}{(s^2 - p^2)(q^2 - p^2)} + \frac{h(-iK_q)K_q}{(s^2 - q^2)(p^2 - q^2)} \right],
 \end{aligned} \tag{A13}$$

with

$$h(-iK)K = \frac{K^2}{(\sinh bK)^2(K \coth bK + \kappa_I)}. \tag{A14}$$

Combining Eqs. (A10) and (A13), we get

$$\begin{aligned}
 \mathbb{M}_\xi &= 2\pi(H_p + \omega_p)\delta(p - q) + (\Delta\mathbb{M}_\xi)_{pq} \\
 (\Delta\mathbb{M}_\xi)_{pq} &= 4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{(H_s + \omega_s)}{(p^2 - s^2)(q^2 - s^2)} + \frac{(H_p + \omega_p)}{(s^2 - p^2)(q^2 - p^2)} + \frac{(H_q + \omega_q)}{(s^2 - q^2)(p^2 - q^2)} \right].
 \end{aligned} \tag{A15}$$

In a similar way,

$$\begin{aligned}
 \mathbb{M}_R &= 2\pi(\omega_p + \kappa_R)\delta(p - q) + (\Delta\mathbb{M}_R)_{pq} \\
 (\Delta\mathbb{M}_R)_{pq} &= 4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{\omega_s}{(p^2 - s^2)(q^2 - s^2)} \right. \\
 &\quad \left. + \frac{\omega_p}{(s^2 - p^2)(q^2 - p^2)} + \frac{\omega_q}{(s^2 - q^2)(p^2 - q^2)} \right].
 \end{aligned} \tag{A16}$$

We follow a similar strategy for \mathcal{M}_{2L} using the integral representation

$$\begin{aligned}
 \omega \coth b\omega &= \int \frac{d\lambda}{\pi} \frac{\omega^2}{\lambda^2 + \omega^2} \tilde{f}(\lambda), \\
 \tilde{f}(\lambda) &= -i \cot b(\lambda - i\epsilon).
 \end{aligned} \tag{A17}$$

This leads to the expression

$$\begin{aligned}
 \mathcal{M}_{2L} &= 2\pi(\omega_p \coth b\omega_p + \kappa_L)\delta(p - q) \\
 &\quad + 4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{\omega_s \coth b\omega_s}{(p^2 - s^2)(q^2 - s^2)} \right. \\
 &\quad \left. + \frac{\omega_p \coth b\omega_p}{(s^2 - p^2)(q^2 - p^2)} + \frac{\omega_q \coth b\omega_q}{(s^2 - q^2)(p^2 - q^2)} \right].
 \end{aligned} \tag{A18}$$

We also get a term $-N_{12L}(-k)(\mathcal{M}_1)^{-1}N_{12L}$ from integration over ϕ_1 . Notice that N_{12L} is the same as $N_{1\xi}$ with $k_2 \rightarrow -k_2$ and an overall minus sign. These do not affect

the final expression, and we get $N_{12L}(-k)(\mathcal{M}_1)^{-1}N_{12L} = N_{1\xi}(\mathcal{M}_1)^{-1}N_{1\xi}$. Combining this with Eq. (A18), we get

$$\begin{aligned}
 \mathbb{M}_L &= 2\pi(H_p + \kappa_L)\delta(p - q) + (\Delta\mathbb{M}_L)_{pq} \\
 (\Delta\mathbb{M}_L)_{pq} &= 4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{H_s}{(p^2 - s^2)(q^2 - s^2)} \right. \\
 &\quad \left. + \frac{H_p}{(s^2 - p^2)(q^2 - p^2)} + \frac{H_q}{(s^2 - q^2)(p^2 - q^2)} \right].
 \end{aligned} \tag{A19}$$

The expression for $Q_{2L\xi}$ simplifies as

$$\begin{aligned}
 Q_{2L\xi} &= -4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{\omega_s \coth b\omega_s}{(p^2 - s^2)(q^2 - s^2)} \right. \\
 &\quad \left. + \frac{\omega_p \coth b\omega_p}{(s^2 - p^2)(q^2 - p^2)} + \frac{\omega_q \coth b\omega_q}{(s^2 - q^2)(p^2 - q^2)} \right],
 \end{aligned} \tag{A20}$$

which is just the second part of the expression for \mathcal{M}_{2L} . The integration over ϕ_1 also produces a new mixing term between ϕ_{2L} and ξ , given by $N_{1\xi}^T(\mathcal{M}_1)^{-1}N_{12L}$, which can be simplified as

$$\begin{aligned}
 N_{1\xi}^T(\mathcal{M}_1)^{-1}N_{12L} &= -4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{h(-iK_s)K_s}{(p^2 - s^2)(q^2 - s^2)} + \frac{h(-iK_p)K_p}{(s^2 - p^2)(q^2 - p^2)} \right. \\
 &\quad \left. + \frac{h(-iK_q)K_q}{(s^2 - q^2)(p^2 - q^2)} \right].
 \end{aligned} \tag{A21}$$

This expression can be combined with Eq. (A20) to get

$$\begin{aligned} N_{2L\xi} &\equiv (Q_{2L\xi} - N_{1\xi}^T(\mathcal{M}_1)^{-1}N_{12L})_{pq} \\ &= -4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{H_s}{(p^2 - s^2)(q^2 - s^2)} \right. \\ &\quad \left. + \frac{H_p}{(s^2 - p^2)(q^2 - p^2)} + \frac{H_q}{(s^2 - q^2)(p^2 - q^2)} \right] \\ &= -(\Delta\mathbb{M}_L)_{pq}. \end{aligned} \quad (\text{A22})$$

The expression for $N_{2R\xi}$ from Eq. (A8) can be simplified as follows:

$$\begin{aligned} N_{2R\xi} &= -4pq \int_{-\infty}^\infty \frac{dk_2}{2\pi} \frac{4pq}{[(k_2 - i\epsilon)^2 - p^2][(k_2 - i\epsilon)^2 - q^2]} \\ &\quad \times \int_{-\infty}^\infty \frac{d\lambda}{\pi} \frac{\omega^2}{\lambda^2 + \omega^2} \\ &= 4pq \int \frac{d\lambda}{2\pi} \frac{\lambda^2}{\sqrt{\lambda^2 + \vec{k}^2}(\lambda^2 + \omega_p^2)(\lambda^2 + \omega_q^2)} \\ &= 4pq \int \frac{d\lambda}{2\pi} \frac{1}{\sqrt{\lambda^2 + \vec{k}^2}} \left[\frac{\omega_p^2}{(\lambda^2 + \omega_p^2)} - \frac{\omega_q^2}{(\lambda^2 + \omega_q^2)} \right] \\ &\quad \times \frac{1}{p^2 - q^2} \\ &= 4pq \left[\frac{\omega_p^2 I(\vec{k}, p) - \omega_q^2 I(\vec{k}, q)}{p^2 - q^2} \right], \end{aligned} \quad (\text{A23})$$

where

$$\begin{aligned} I(\vec{k}, p) &= \int_0^\infty \frac{d\lambda}{\pi} \frac{1}{\sqrt{\lambda^2 + \vec{k}^2}(\lambda^2 + \omega_p^2)} \\ &= \frac{1}{2\pi} \frac{1}{p\omega_p} \log \left(\frac{\omega_p + p}{\omega_p - p} \right). \end{aligned} \quad (\text{A24})$$

By rewriting $1/\sqrt{\lambda^2 + \vec{k}^2}$ as we did for the others and simplifying, we can also see that

$$N_{2R\xi} = -(\Delta\mathbb{M}_R)_{pq}. \quad (\text{A25})$$

We now turn to the ρ -dependent terms.

$\rho\rho$ term

The kernel \mathcal{M}_ρ is $\omega \coth b\omega + \omega$; it is easier to simplify the $\rho\rho$ term after the integration over ϕ_1 . This yields the kernel $\mathbb{M}_\rho = \mathcal{M}_\rho - (\omega \operatorname{csch} b\omega)^2 (\mathcal{M}_1)^{-1} = H + \omega$. This gives directly

$$\mathbb{M}_\rho = 4|\vec{k}|^2 \int_{-\infty}^\infty \frac{ds}{2\pi} \frac{H_s + \omega_s}{\omega_s^4} + \frac{c_0}{2|\vec{k}|}. \quad (\text{A26})$$

$\rho\xi$ term

This term has $\omega + \omega \coth b\omega - N_{1\xi}(\mathcal{M}_1)^{-1}N_{1\rho} = H + \omega$ as the kernel, so we get

$$\begin{aligned} N_{\rho\xi} &= \int \frac{dk_2}{2\pi} \frac{2p}{p^2 - (k_2 - i\epsilon)^2} \frac{2|\vec{k}|}{\omega_k^2} (H_k + \omega_k) \\ &= \int \frac{d\lambda}{\pi} \frac{dk_2}{2\pi} \frac{4|\vec{k}|p}{[p^2 - (k_2 - i\epsilon)^2](\lambda^2 + \omega_k^2)} (f(\lambda) + h(\lambda)) \\ &= \int \frac{d\lambda}{2\pi} \frac{4|\vec{k}|p}{\sqrt{\lambda^2 + \vec{k}^2}(\lambda^2 + \omega_p^2)} (f(\lambda) + h(\lambda)) \\ &= \int_0^\infty \frac{ds}{\pi} \int_{-\infty}^\infty \frac{d\lambda}{\pi} \frac{4|\vec{k}|p}{(\lambda^2 + \omega_s^2)(\lambda^2 + \omega_p^2)} (f(\lambda) + h(\lambda)) \\ &= 4|\vec{k}|p \int_0^\infty \frac{ds}{\pi} \left[\frac{H_s + \omega_s}{\omega_s^2} - \frac{H_p + \omega_p}{\omega_p^2} \right] \frac{1}{p^2 - s^2}. \end{aligned} \quad (\text{A27})$$

$\phi_{2L\rho}$ term

The kernel for this term is $\omega \coth b\omega + c_{2L} - N_{1\rho}(\mathcal{M}_1)^{-1}N_{12L} = H + c_{2L}$ and gives

$$\begin{aligned} N_{2L\rho} &= \int \frac{dk_2}{2\pi} \frac{2p}{p^2 - (k_2 - i\epsilon)^2} \frac{2|\vec{k}|}{\omega^2} (H + c_{2L}) \\ &= 4|\vec{k}|p \left[\int_0^\infty \frac{ds}{\pi} \left(\frac{H_s}{\omega_s^2} - \frac{H_p}{\omega_p^2} \right) \frac{1}{p^2 - s^2} + \frac{c_{2L}}{2|\vec{k}|\omega_p^2} \right]. \end{aligned} \quad (\text{A28})$$

For the first term on the right hand side, we simplified as we did for $N_{\rho\xi}$, and for the second term, we did the k_2 integral completing the contour in the lower half-plane.

$\phi_{2R\rho}$ term

This is similar to the $\phi_{2L\rho}$ term, except that we have $\omega + c_{2R}$ instead of $H + c_{2L}$. We get

$$\begin{aligned} N_{2R\rho} &= - \int \frac{dk_2}{2\pi} \frac{4|\vec{k}|p}{[p^2 - (k_2 - i\epsilon)^2]\omega^2} (\omega + c_{2R}) \\ &= - \int \frac{d\lambda}{\pi} \frac{dk_2}{2\pi} \frac{4|\vec{k}|p}{[p^2 - (k_2 - i\epsilon)^2]\omega^2} \frac{\omega^2}{\omega^2 + \lambda^2} \\ &\quad - c_{2R} \int \frac{dk_2}{2\pi} \frac{4|\vec{k}|p}{[p^2 - (k_2 - i\epsilon)^2]\omega^2} \\ &= -4|\vec{k}|p \int \frac{d\lambda}{2\pi} \frac{1}{\sqrt{\lambda^2 + \vec{k}^2}(\lambda^2 + \omega_p^2)} - c_{2R} \frac{2p}{\omega_p^2} \\ &= -4|\vec{k}|p \left[I(\vec{k}, p) + \frac{c_{2R}}{2|\vec{k}|\omega_p^2} \right]. \end{aligned} \quad (\text{A29})$$

Finally, we note that in the calculation of the Casimir energy by expansion of the $\operatorname{Tr} \log$, the first-order correction involves the diagonal elements of the type Δ_{pp} . In some cases (such as $\Delta\mathbb{M}_L$, $\Delta\mathbb{M}_\xi$, etc.), Δ_{pq} is of the form

$$\Delta_{pq} = 4pq \int_0^\infty \frac{ds}{\pi} \left[\frac{W_s}{(p^2 - s^2)(q^2 - s^2)} + \frac{W_p}{(s^2 - p^2)(q^2 - p^2)} + \frac{W_q}{(s^2 - q^2)(p^2 - q^2)} \right]. \quad (\text{A30})$$

The $p = q$ limit can be extracted as

$$\Delta_{pp} = 4p^2 \int \frac{ds}{\pi} \left[\frac{W_s - W_p}{(s^2 - p^2)^2} - \left(\frac{\partial W}{\partial p^2} \right) \frac{1}{(s^2 - p^2)} \right]. \quad (\text{A31})$$

We will need this expression with the appropriate W 's to estimate the contribution to the energy numerically.

-
- [1] H.B.G. Casimir, Proc. K. Ned. Akad. Wet. **51**, 793 (1948); H.B.G. Casimir and D. Polder, *Phys. Rev.* **73**, 360 (1948).
- [2] For general reviews on the Casimir effect, see K.A. Milton, *J. Phys. Conf. Ser.* **161**, 012001 (2009); K.A. Milton, *The Casimir Effect: Physical Manifestations of Zero-Point Energy* (World Scientific, Singapore, 2001); M. Bordag, U. Mohideen, and V.M. Mostepanenko, *Phys. Rep.* **353**, 1 (2001); M. Bordag, G.L. Klimchitskaya, U. Mohideen, and V.M. Mostepanenko, *Advances in the Casimir Effect* (Oxford University Press, Oxford, 2009).
- [3] D. Kabat, D. Karabali, and V.P. Nair, *Phys. Rev. D* **81**, 125013(E) (2010); **84**, 129901 (2011).
- [4] D. Kabat, D. Karabali, and V.P. Nair, *Phys. Rev. D* **82**, 025014 (2010).
- [5] D. Kabat and D. Karabali, *Phys. Rev. D* **84**, 065029 (2011).
- [6] D. Karabali, *J. Phys. Conf. Ser.* **343**, 012053 (2012).
- [7] H. Gies and K. Klingmüller, *Phys. Rev. Lett.* **96**, 220401 (2006).
- [8] K. Klingmüller and H. Gies, *J. Phys. A* **41**, 164042 (2008); A. Weber and H. Gies, *Phys. Rev. D* **80**, 065033 (2009); H. Gies and A. Weber, *Int. J. Mod. Phys. A* **25**, 2279 (2010).
- [9] N. Graham, A. Shpunt, T. Emig, S.J. Rahi, R.L. Jaffe, and M. Kardar, *Phys. Rev. D* **81**, 061701 (2010); M.F. Maghrebi and N. Graham, *Europhys. Lett.* **95**, 14001 (2011); M.F. Maghrebi, *Phys. Rev. D* **83**, 045004 (2011); N. Graham, A. Shpunt, T. Emig, S.J. Rahi, R.L. Jaffe, and M. Kardar, *Phys. Rev. D* **83**, 125007 (2011).
- [10] M.F. Maghrebi, S.J. Rahi, T. Emig, N. Graham, R.L. Jaffe, and M. Kardar, *Proc. Natl. Acad. Sci. U.S.A.* **108**, 6867 (2011).
- [11] A. Romeo and A. A. Saharian, *J. Phys. A* **35**, 1297 (2002).
- [12] L. C. de Albuquerque and R. M. Cavalcanti, *J. Phys. A* **37**, 7039 (2004).
- [13] J. von Neumann, *Math. Ann.* **102**, 49 (1930).
- [14] M. Asorey, A. Ibort, and G. Marmo, *Int. J. Mod. Phys. A* **20**, 1001 (2005); M. Asorey, D. Garcia-Alvarez, and J. M. Munoz-Castaneda, *J. Phys. A* **39**, 6127 (2006); M. Asorey and J. M. Munoz-Castaneda, *J. Phys. A* **41**, 304004 (2008).
- [15] T.R. Govindarajan and V.P. Nair, in preparation.