

September 2013

# Specificity in the actions of the UBR1 ubiquitin ligase in the degradation of nuclear receptors<sup>✉</sup>

Rasheda Sultana

Maria A. Theodoraki

Avrom J. Caplan

## [How does access to this work benefit you? Let us know!](#)

Follow this and additional works at: [http://academicworks.cuny.edu/cc\\_pubs](http://academicworks.cuny.edu/cc_pubs)

---

### Recommended Citation

Sultana, R., Theodoraki, M. A. & Caplan, A. J. (2013). Specificity in the actions of the UBR1 ubiquitin ligase in the degradation of nuclear receptors<sup>✉</sup>. *FEBS Open Bio*, 3, 394-397. doi:10.1016/j.fob.2013.09.003.

This Article is brought to you for free and open access by the City College of New York at CUNY Academic Works. It has been accepted for inclusion in Publications and Research by an authorized administrator of CUNY Academic Works. For more information, please contact [AcademicWorks@cuny.edu](mailto:AcademicWorks@cuny.edu).



ELSEVIER



CrossMark

journal homepage: [www.elsevier.com/locate/febsopenbio](http://www.elsevier.com/locate/febsopenbio)

# Specificity in the actions of the UBR1 ubiquitin ligase in the degradation of nuclear receptors<sup>☆</sup>

Rasheda Sultana, Maria A. Theodoraki, Avrom J. Caplan\*

Department of Biology, The City College of New York, New York, NY 10031, USA

## ARTICLE INFO

### Article history:

Received 25 June 2013

Received in revised form 12 September 2013

Accepted 12 September 2013

### Keywords:

Ubiquitin ligase

Quality control

Degradation

Molecular chaperone

Hsp90

UBR1

## ABSTRACT

**The UBR1 ubiquitin ligase promotes degradation of proteins via the N-end rule and by another mechanism that detects a misfolded conformation. Although UBR1 was shown recently to act on protein kinases whose misfolding was promoted by inhibition of Hsp90, it was unknown whether this ubiquitin ligase targeted other client types of the chaperone. We analyzed the role of UBR1 in the degradation of nuclear receptors that are classical clients of Hsp90. Our results showed that UBR1 deletion results in impaired degradation of the glucocorticoid receptor and the androgen receptor but not the estrogen receptor  $\alpha$ . These findings demonstrate specificity in the actions of the UBR1 ubiquitin ligase in the degradation of Hsp90 clients in the presence of small molecule inhibitors that promote client misfolding.**

© 2013 The Authors. Published by Elsevier B.V. on behalf of Federation of European Biochemical Societies. All rights reserved.

## Introduction

The intracellular environment does not favor protein stability. Newly made proteins fold under conditions that are relatively hot and crowded, which favor misfolding or aggregation rather than acquisition of the correct conformation [1]. These environmental conditions necessitated the evolution of a large cohort of genes that protect newly made proteins or promote degradation of those that misfold. The process that is governed by these genes is known variously as protein quality control or proteostasis [2,3]. The cellular machinery that comprises the proteostasis gene network includes molecular chaperones that interact with newly made or misfolded proteins [4] and components of the ubiquitin/proteasome system (UPS) for disposal of misfolded proteins [5]. In addition, the autophagic machinery facilitates disposal of insoluble aggregates that accumulate upon failure or imbalance of molecular chaperones or UPS components as cells age or upon stress [6]. The proteostasis network is sensitive to environmental conditions and adjusts the expression of a variety of genes to protect against proteotoxic stress [7].

<sup>☆</sup> This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Abbreviations:** AR, androgen receptor; CHIP, C-terminal Hsp interacting protein; DMSO, dimethylsulfoxide; ER, estrogen receptor; GA, geldanamycin; GR, glucocorticoid receptor; HA, hemagglutinin; MEF, mouse embryonic fibroblasts; UPS, ubiquitin proteasome system.

\* Corresponding author. Address: Department of Biology, The City College of New York, Convent Avenue at 138th Street, New York, NY 10031, USA. Tel.: +1 (646) 664 8904; fax: +1 (212) 650 8585.

E-mail address: [avrom.caplan@cuny.edu](mailto:avrom.caplan@cuny.edu) (A.J. Caplan).

The Hsp90 molecular chaperone sits at a hub of the proteostasis network by integrating folding of newly made proteins and degradation of misfolded ones. The balance of these triage decisions can be altered by small molecule inhibitors of Hsp90, which promote client protein degradation [8]. Many of these small molecules are in clinical trials as chemotherapeutic agents because of the role Hsp90 plays in the folding and degradation of oncogenic protein kinases and nuclear receptors [9].

Although it is clear that Hsp90 inhibition results in rapid degradation of client proteins via the UPS, the mechanisms underlying the clearance process are not well characterized. For example, the ubiquitin ligase C-terminal Hsp interacting protein (CHIP), plays a role in the degradation of several Hsp90 clients including protein kinases and nuclear receptors [10]. CHIP is not alone in its capacity to promote degradation of misfolded Hsp90 clients and several ubiquitin ligases have now been characterized to act in this pathway in yeast and mammalian cells [11]. What is unclear, however, is the specificity by which these ubiquitin ligases promote degradation when Hsp90 is inhibited. A good example of this is the glucocorticoid receptor (GR). GR can be ubiquitinated *in vitro* by CHIP [12] or upon CHIP overexpression [13], although degradation of this receptor is not affected by deletion of CHIP in cells [14], suggesting that other ubiquitin ligases act in a redundant manner. In this report, we show that the ubiquitin ligase UBR1 promotes degradation of the GR and the androgen receptor (AR) but not the estrogen receptor (ER) in cells treated with an Hsp90 inhibitor. This finding is significant in view of the direct role of the AR in promoting growth of prostate cancer cells.

## Materials and methods

### Materials

Geldanamycin was purchased from InvivoGen (San Diego, CA), MG132 was purchased from Calbiochem (San Diego, CA), and both compounds were dissolved in 100% DMSO.

### Cell culture, transfection and plasmids

WT, *Ubr1*<sup>-/-</sup> and *Chip*<sup>-/-</sup> mouse embryonic fibroblast cells were maintained in DMEM medium supplemented with 10% heat-inactivated fetal Bovine serum (FBS) (Mediatech Inc., Herdon, VA), 100 units/ml penicillin, 100 µg/ml streptomycin (MP Biomedicals, LLC, France) and kept at 37 °C in 5% CO<sub>2</sub> incubator. The transfection was performed as described [15]. For each transfection reaction 4 µg of plasmid DNA was used unless indicated. The plasmid encoding human AR, HA tagged GR and ER-α were gift from Dr. Michael J. Garabedian (New York School of Medicine). The plasmid encoding the rat UBR1 was a gift from Dr. Hiroshi Handa (Integrated Research Institute, Tokyo Institute of Technology, Yokohama, Japan). The WT, *Ubr1*<sup>-/-</sup> cells were kind gift from Dr. Yong Tae Kwon (University of Pittsburg, Pennsylvania) and *Chip*<sup>-/-</sup> cells from Dr. Cam Patterson (University of North Carolina).


### Western blotting and antibodies

Cells were transfected or grown to 70–80% confluence and exposed to DMSO, GA and MG132 for indicated time and dose. Lysates were prepared using lysis buffer containing 50 mM Tris pH 7.5, 2% SDS, .25% Na-deoxycholate, 150 mM NaCl, 1 mM EDTA, 10% glycerol, 1 mM phenylmethylsulfonyl fluoride, 10 mM NaF, 1 mM Na<sub>3</sub>VO<sub>4</sub> and protease inhibitors (Complete mini, Roche Diagnostics, Indianapolis, IN). Lysates were sonicated for 3–4 times, 10 s each time. Protein concentration of lysates was determined using Bicinchoninic acid method (Pierce, Rockford, IL). Samples of 40 µg were analyzed in SDS-polyacrylamide gels and followed the same procedure described previously [15]. Antibodies used were: GR (MA1-510, ThermoScientific, Rockford, IL, USA), ER-α (sc-543, Santa Cruz Biotechnology, Inc.), AR [16]. PI3K (06-497, Millipore). Anti-HA (12CA5 from the Mount Sinai Hybridoma Facility, New York, NY) and UBR1 (Abcam Inc., Boston, MA).

## Results

In previous studies we demonstrated that mammalian UBR1 acted as part of the quality control apparatus that helps to clear misfolded protein kinases from cells treated with the Hsp90 inhibitor, geldanamycin (GA) [15]. UBR1 action was at least partially redundant with CHIP in this capacity. Based on these studies we tested whether UBR1 acted in the clearance of nuclear receptors, which represent another well-characterized class of Hsp90 client [17].

We first analyzed the ability of GA to promote degradation of the endogenously expressed GR in mouse embryonic fibroblasts (MEF). Treatment of cells with 100 nM GA resulted in a very rapid degradation of GR within a two-hour period and these levels were further diminished after 6 hours of treatment (Fig. 1A). GA also promoted degradation of GR in the *Ubr1*<sup>-/-</sup> cells but at a reduced rate, and there was still substantial GR remaining after 6 h of treatment. By contrast, *Chip*<sup>-/-</sup> cells behaved much like wild type MEFs with respect to GR degradation, as reported by others [14]. Similar findings were made in a dose-response assay where the levels of GR were barely affected at a dose of 100 nM in *Ubr1*<sup>-/-</sup> cells, whereas in wild type or *Chip*<sup>-/-</sup> MEFs the protein was mostly degraded (Fig. 1B). These findings suggest that UBR1 is important for the degradation of




**Fig. 1.** Geldanamycin dependent degradation of GR in WT, *Ubr1*<sup>-/-</sup> and *Chip*<sup>-/-</sup> MEF cells. (A) WT, *Ubr1*<sup>-/-</sup> and *Chip*<sup>-/-</sup> MEF cells were treated with 100 nM of GA for indicated times. 40 µg of total protein from each cell line were analyzed by SDS-PAGE and probed with anti-GR; PI3K was used as a loading control. (B) WT, *Ubr1*<sup>-/-</sup> and *Chip*<sup>-/-</sup> MEF cells were treated with different concentration of GA for 6 hours. 40 µg of total protein from each cell line were fractionated by SDS-PAGE and probed with anti-GR. PI3K levels were analyzed as a loading control.

GR upon Hsp90 inhibition and that additional ubiquitin ligases distinct from CHIP also act in this capacity. We analyzed whether UBR2, a UBR1 homolog, acted in a similar capacity to UBR1 but found that GR degradation was similar in *UBR2*<sup>-/-</sup> cells and wild type MEFs (data not shown).

To determine the UBR1 dependence on the degradation phenotype of *Ubr1*<sup>-/-</sup> cells, we used transfection to introduce wild type rat UBR1 (rUBR1) into the cells. An HA-tagged version of GR (HA-GR), or an empty vector as a control were co-transfected into *Ubr1*<sup>-/-</sup> cells (Fig. 2A). The effect of rUBR1 overexpression was to cause a large reduction in the levels of the transfected HA-GR even without GA treatment. Furthermore, treatment with GA did not lead to a substantial decrease in the HA-GR levels after 2 h. These findings suggested that HA-GR was sensitive to the levels of rUBR1 even in the absence of GA. The effect of GA was probed by a more prolonged treatment of 18 h (Fig. 2B), and in this case we observed that the GA-sensitivity of the transfected HA-GR was restored. Note also that rUBR1 levels were also sensitive to GA treatment as described previously [15]. HA-UBR1 levels were restored upon treatment with the proteasome inhibitor, MG132, indicating that UBR1 was acting via the UPS as expected, although this also occurred in the vector-transfected control cells, thus confirming redundancy in UBR1 action.

The large decrease in HA-GR levels upon co-transfection of rUBR1 (Fig. 2A) was studied further in a dose-response experiment (Fig. 2C). In this case we observed a direct correlation between increasing rUBR1 concentration and decreasing HA-GR levels, further suggesting that UBR1 controls GR levels both in the absence and presence of functional Hsp90.

The results described thus far suggest that UBR1 promotes degradation of GR and further studies addressed the specificity of that function with analysis of the human androgen receptor (hAR) and estrogen receptor α (hERα). hAR was transfected into the *Ubr1*<sup>-/-</sup> cells with either the plasmid overexpressing rat UBR1 or an empty vector. The findings (Fig. 3A) were very similar those observed for transfection of HA-GR (Fig. 2). In this case, there was a substantial decrease in hAR levels when rUBR1 was co-overexpressed in the *Ubr1*<sup>-/-</sup> cells compared with vector alone even in the absence of GA. In both cases, however, there was little further effects on degradation of hAR by GA




**Fig. 2.** Effect of UBR1 overexpression on GR levels. (A)  $Ubr1^{-/-}$  cells were transfected with human HA-tagged GR (HA-hGR) and pCMV empty vector or rat UBR1. After 24 h of transfection the cells were treated with 50 nM of GA or DMSO for the indicated time points and harvested. 40  $\mu$ g of cell lysates were analyzed in SDS-PAGE and probed with anti-HA ( $\alpha$ -HA) and UBR1 antisera. PI3K was used as loading control. \* in the PI3K blot indicates a non-specific band. (B)  $Ubr1^{-/-}$  cells were transfected with HA-hGR and empty vector or rat UBR1. After transfection cells were treated with DMSO or MG-132 (50  $\mu$ M) for 18 h and GA (100 nM) for 6 h. Cells were harvested 24 h after transfection. 40  $\mu$ g of cell lysates were analyzed in SDS-PAGE and probed with anti-HA ( $\alpha$ -HA), UBR1 and PI3K. (C)  $Ubr1^{-/-}$  cells were transfected with HA-hGR and different amounts of rat E3 ligase UBR1 plasmid DNA (0, 0.5, 2 and 4  $\mu$ g). Cell were harvested 24 h after transfection. 40  $\mu$ g of total cell lysates were fractionated by SDS-PAGE and probed with anti-HA ( $\alpha$ -HA) and UBR1 antisera. PI3K was used as a loading control. NT represents non-transfected  $Ubr1^{-/-}$  MEF cells in A–C.

treatment (Fig. 3A and B), although rUBR1 levels were sensitive at the highest treatment level of GA. Treatment of these cells with MG132, however, led to a strong resurgence of hAR levels that was particularly evident in the cells co-expressing rUBR1. These data strongly support a role for UBR1 in the quality control of hAR levels. By contrast, similar experiments with transfected human hER $\alpha$  suggest that UBR1 does not play a similar role in the degradation of this nuclear receptor (Fig. 3C). Co-transfection of rUBR1 did not affect hER $\alpha$  in the same manner it did with GR and hAR. Our combined findings therefore suggest specificity in the actions of UBR1 in the clearance of GR, AR and ER, with the AR displaying the most dependence on UBR1 actions.

## Discussion

Quality control ubiquitin ligases promote degradation of misfolded proteins. In yeast there exists a small network of quality control ubiquitin ligases that function in the degradation of Hsp90 clients that includes UBR1 and at least 7 others acting in the cytosol and the nucleus [11]. In mammalian cells, there are three ubiquitin ligases that are known to facilitate the clearance of protein kinases that misfold upon Hsp90 inhibition and these include CHIP, CUL5 and UBR1 (reviewed in [8]). It is striking, however, that neither CHIP nor CUL5 affect the degradation of GR when they are deleted or down-regulated [14,18], supporting the hypothesis that redundant ubiquitin ligases exist. In the studies presented in this report we identify UBR1 as having this specific function in GR degradation, and also show that UBR1 acts in the clearance of the hAR, but not hER $\alpha$ .

UBR1 is a well-characterized ubiquitin ligase that functions via N-end rule degradation and also via non-N-end rule pathways in protein quality control [19]. In yeast, Ubr1 acts via non-N-end rule pathways to promote degradation of misfolded proteins [20–22] and it seems likely that this will be the case for mammalian UBR1 based on its function in the clearance of nuclear receptors, as shown here, and protein kinases [15] that misfold upon Hsp90 inhibition. Furthermore, dipeptides [23,24] known to inhibit degradation via the N-end rule (Leu-Ala and Arg-Ala) had no effect on the degradation of GR and AR



**Fig. 3.** Effects of UBR1 overexpression on AR and ER- $\alpha$ . (A)  $Ubr1^{-/-}$  cells were transfected with the AR expressing plasmid and pCMV empty vector or rat UBR1. After 24 h of transfection the cells were treated with different concentrations of GA or vehicle for 24 h. 40  $\mu$ g of cell lysates were analyzed in SDS-PAGE and probed with AR and UBR1 antisera. PI3K was used as loading control. (B)  $Ubr1^{-/-}$  cells were transfected with AR and pCMV empty vector or rat UBR1. After transfection the cells were treated with DMSO, MG-132 (50  $\mu$ M) for 18 h and GA (100 nM) for 6 h. The cells were harvested 24 h after transfection and 40  $\mu$ g of cell lysates were analyzed by SDS-PAGE and probed with anti AR, UBR1 and PI3K. (C)  $Ubr1^{-/-}$  cells were transfected a plasmid expressing ER- $\alpha$  or pCMV empty vector or rat UBR1. After 24 h of transfection, the cells were treated with different concentrations of GA or DMSO for 24 h. 40  $\mu$ g of cell lysates were analyzed by SDS-PAGE and probed with AR and UBR1 antisera. PI3K was used as loading control. NT represents non-transfected  $Ubr1^{-/-}$  MEF cells in A–C.

(data not shown).

As mentioned above, yeast E3s that act in cytosolic quality control are highly redundant. In mammalian cells there is also some redundancy in the actions of E3 ligases in cellular quality control. In a previous study we observed redundancy between CHIP and UBR1 for the degradation of CDK4 and AKT protein kinases [15]. By contrast, there does not appear to be a role for CHIP in the degradation of GR that misfolds due to Hsp90 inhibition. CHIP can promote GR degradation when it is overexpressed, but as discovered by others, CHIP knockout cells have no impairment in misfolded GR degradation [14]. In this case, therefore, there appears to be specificity in E3 action. This specificity is further supported by our finding that overexpression of UBR1 had no effect on the degradation of ER- $\alpha$ , whereas downregulation of CHIP does inhibit ER- $\alpha$  degradation [25].

In conclusion, our findings provide further support for a conserved role for UBR1 in the degradation of misfolded proteins in mammalian cells. The finding that UBR1 acts on some misfolded nuclear receptors suggests it may have a role to play in a variety of human diseases of aging including cancer as well as Johanson–Blizzard syndrome [26].

## Acknowledgments

We thank Dr. Michael J. Garabedian, Dr. Hiroshi Handa, Dr. Yong Tae Kwon and Dr. Cam Patterson for the gift of plasmids and cell lines. This work was funded in part by grants from the National Institutes of Health (U54CA132378) and NCCR (5G12-RR03060) (CCNY).

## References

- [1] Ellis, R.J. (2007) Protein misassembly: macromolecular crowding and molecular chaperones. *Adv. Exp. Med. Biol.* 594, 1–13.
- [2] Balch, W.E., Morimoto, R.I., Dillin, A. and Kelly, J.W. (2008) Adapting proteostasis for disease intervention. *Science* 319, 916–919.
- [3] Wickner, S., Maurizi, M.R. and Gottesman, S. (1999) Posttranslational quality control: folding, refolding, and degrading proteins. *Science* 286, 1888–1893.
- [4] Hartl, F.U., Bracher, A. and Hayer-Hartl, M. (2011) Molecular chaperones in protein folding and proteostasis. *Nature* 475, 324–332.
- [5] Hochstrasser, M. (1996) Ubiquitin-dependent protein degradation. *Annu. Rev. Genet.* 30, 405–439.

- [6] Ketterer, N., Dreiseidler, M., Tawo, R. and Hohfeld, J. (2010) Chaperone-assisted degradation: multiple paths to destruction. *Biol. Chem.* 391, 481–489.
- [7] Richter, K., Haslbeck, M. and Buchner, J. (2010) The heat shock response: life on the verge of death. *Mol. Cell* 40, 253–266.
- [8] Theodoraki, M.A. and Caplan, A.J. (2012) Quality control and fate determination of Hsp90 client proteins. *Biochim. Biophys. Acta* 1823, 683–688.
- [9] Trepel, J., Mollapour, M., Giaccone, G. and Neckers, L. (2010) Targeting the dynamic HSP90 complex in cancer. *Nat. Rev. Cancer* 10, 537–549.
- [10] McDonough, H. and Patterson, C. (2003) CHIP: a link between the chaperone and proteasome systems. *Cell Stress Chaperones* 8, 303–308.
- [11] Theodoraki, M.A., Nillegoda, N.B., Saini, J. and Caplan, A.J. (2012) A network of ubiquitin ligases is important for the dynamics of misfolded protein aggregates in yeast. *J. Biol. Chem.* 287, 23911–23922.
- [12] Connell, P., Ballinger, C.A., Jiang, J., Wu, Y., Thompson, L.J., Hohfeld, J. et al. (2001) The co-chaperone CHIP regulates protein triage decisions mediated by heat-shock proteins. *Nat. Cell Biol.* 3, 93–96.
- [13] Cardozo, C.P., Michaud, C., Ost, M.C., Fliss, A.E., Yang, E., Patterson, C. et al. (2003) C-terminal Hsp-interacting protein slows androgen receptor synthesis and reduces its rate of degradation. *Arch. Biochem. Biophys.* 410, 134–140.
- [14] Morishima, Y., Wang, A.M., Yu, Z., Pratt, W.B., Osawa, Y. and Lieberman, A.P. (2008) CHIP deletion reveals functional redundancy of E3 ligases in promoting degradation of both signaling proteins and expanded glutamine proteins. *Human Mol. Genet.* 17, 3942–3952.
- [15] Sultana, R., Theodoraki, M.A. and Caplan, A.J. (2012) UBR1 promotes protein kinase quality control and sensitizes cells to Hsp90 inhibition. *Exp. Cell Res.* 318, 53–60.
- [16] Fang, Y., Fliss, A.E., Robins, D.M. and Caplan, A.J. (1996) Hsp90 regulates androgen receptor hormone binding affinity in vivo. *J. Biol. Chem.* 271, 28697–28702.
- [17] Pratt, W.B., Morishima, Y., Murphy, M. and Harrell, M. (2006) Chaperoning of glucocorticoid receptors *Handb. Exp. Pharmacol.*, 111–138.
- [18] Ehrlich, E.S. (2009) Regulation of Hsp90 client proteins by a Cullin5-RING E3 ubiquitin ligase. *Proc. Natl. Acad. Sci. U.S.A.* 106, 20330–20335.
- [19] Varshavsky, A. (2011) The N-end rule pathway and regulation by proteolysis. *Protein Sci.* 20, 1298–1345.
- [20] Eisele, F. and Wolf, D.H. (2008) Degradation of misfolded protein in the cytoplasm is mediated by the ubiquitin ligase Ubr1. *FEBS Lett.* 582, 4143–4146.
- [21] Heck, J.W., Cheung, S.K. and Hampton, R.Y. (2010) Cytoplasmic protein quality control degradation mediated by parallel actions of the E3 ubiquitin ligases Ubr1 and San1. *Proc. Natl. Acad. Sci. U.S.A.* 107, 1106–1111.
- [22] Nillegoda, N.B. (2010) Ubr1 and Ubr2 function in a quality control pathway for degradation of unfolded cytosolic proteins. *Mol. Biol. Cell* 21, 2102–2116.
- [23] Baker, R.T. and Varshavsky, A. (1991) Inhibition of the N-end rule pathway in living cells. *Proc. Natl. Acad. Sci. U.S.A.* 88, 1090–1094.
- [24] Du, F., Navarro-Garcia, F., Xia, Z., Tasaki, T. and Varshavsky, A. (2002) Pairs of dipeptides synergistically activate the binding of substrate by ubiquitin ligase through dissociation of its autoinhibitory domain. *Proc. Natl. Acad. Sci. U.S.A.* 99, 14110–14115.
- [25] Fan, M., Park, A. and Nephew, K.P. (2005) CHIP (carboxyl terminus of Hsc70-interacting protein) promotes basal and geldanamycin-induced degradation of estrogen receptor- $\alpha$ . *Mol. Endocrinol.* 19, 2901–2914.
- [26] Zenker, M. (2005) Deficiency of UBR1, a ubiquitin ligase of the N-end rule pathway, causes pancreatic dysfunction, malformations and mental retardation (Johanson-Blizzard syndrome). *Nat. Genet.* 37, 1345–1350.