

2015

## Toxins in Newtown Creek

Burl Yearwood

*CUNY La Guardia Community College*

Cho Cho Aung

*CUNY La Guardia Community College*

Ridima Pradhan

*CUNY La Guardia Community College*

Jennifer Vance

*CUNY La Guardia Community College*

[How does access to this work benefit you? Let us know!](#)

Follow this and additional works at: [https://academicworks.cuny.edu/lg\\_pubs](https://academicworks.cuny.edu/lg_pubs)

 Part of the [Environmental Sciences Commons](#)

---

### Recommended Citation

Yearwood, Burl; Aung, Cho Cho; Pradhan, Ridima; and Vance, Jennifer, "Toxins in Newtown Creek" (2015). *CUNY Academic Works*.  
[https://academicworks.cuny.edu/lg\\_pubs/103](https://academicworks.cuny.edu/lg_pubs/103)

This Article is brought to you for free and open access by the LaGuardia Community College at CUNY Academic Works. It has been accepted for inclusion in Publications and Research by an authorized administrator of CUNY Academic Works. For more information, please contact [AcademicWorks@cuny.edu](mailto:AcademicWorks@cuny.edu).

# Toxins in Newtown Creek

Burl Yearwood, Cho Cho Aung, Ridima Pradhan, Jennifer Vance\*

Natural Sciences, LaGuardia Community College, Long Island City, United States

**Abstract** Newtown Creek is a 3.5 mile river which empties into the East River in New York, United States. The river runs the boundary between the Queens and Brooklyn Burroughs. A thriving sight of industry, this river has been the location of much pollution for over a century. Battered by oil spills and oil seepage, raw sewage, trash carried by rain water, cement, animal fat, and a 15 meter sludge on the bottom of the riverbed, Newtown Creek is one of the most polluted waterways in the United States. In this article, data for nitrate, phosphate, lead, copper, dissolved iron, and chromate were collected over 6 months. Levels of lead were 10 times higher than the drinking water limit, and then diminished greatly over the 6 months.

**Keywords** Nitrate, Chromate, Phosphate, Lead, Copper, Iron, Newtown Creek

## 1. Introduction

With an extensive history of pollution, Newtown Creek has been the site of multiple oil refineries for over a century. Early articles describing the creek, include the explicit depictions of the pungent odor of oil [1]. The amount of oil estimated to have been leaked into the creek is between 17-30 million gallons [64-114 million liters], which is twice the amount of oil spilt by the infamous Exxon Valdez [2]. Dumping of cement and animal fat occurred as well. In fact the bottom of the creek is coated in 15 feet of sludge, which has been nicknamed “black mayonnaise” [3]. Since 1866, New York City has also contaminated the creek when combined sewer overflows carry raw sewage and garbage into the waterway [4].

In September, 2010 Newtown Creek was designated as a Region 2 superfund site by the Environmental Protection Agency (EPA). A superfund site is designated as a hazardous waste area, which the EPA will oversee the remediation. Either the polluting parties pay for the clean-up, or the government pays. In the case of Newtown Creek several oil companies are being held responsible for a portion of the payment [4, 5]. Since the superfund site designation, the EPA has been characterizing the toxins present in the river. The toxins included polychlorinated biphenyls (PCBs), volatile organic compounds (VOCs), pesticides, and heavy metals [4].

Our team investigated the presence of nitrate, phosphate, lead, copper, dissolved iron, and chromate in Newtown Creek over 6 months, from August to January of 2014-2015. Our aim was to quantify these pollutants and compare them to the accepted levels. Samples were taken from Whale

creek, which is part of Newtown Creek (Figure 1). The Newtown Creek Wastewater plant, Allocco Recycling Corporation, and Sims Metal Management are in this immediate area.




Figure 1. Map of Newtown Creek and Whale Creek

We investigated these toxins for several different reasons. High lead levels lead to developmental delays in children. In adults, lead contributes to kidney damage and high blood pressure. Copper can lead to liver or kidney damage and chromate can lead to allergic dermatitis. Nitrate contributes to blue baby syndrome in infants [6]. High dissolved iron levels have been found to be damaging to aquatic life [7]. High phosphate levels encourage algae growth, which is damaging to aquatic life. [8] In Newtown Creek, some fish, such as striped bass, American eel, and blue fish are present and unfortunately fished. Blue crabs are caught for recreation, much to the surmise of the EPA. The EPA is planning on testing the tissues of blue crab and striped bass for toxins in Phase 2 of the remedial investigation. People

\* Corresponding author:

jvance@lagcc.cuny.edu (Jennifer Vance)

Published online at <http://journal.sapub.org/env>

Copyright © 2015 Scientific & Academic Publishing. All Rights Reserved

have also been observed swimming in the creek with direct exposure to the water. There are only three public access points to the creek, but exposure to the water and sediment is of concern to the EPA [9].




Figure 2. Nitrate Levels




Figure 3. Copper Levels




Figure 4. Iron Levels




Figure 5. Chromate Levels




Figure 6. Phosphate Levels




Figure 7. Lead Levels

Table 1. Toxin Levels in Newtown Creek in ppm

Toxin	25-Jul	1-Aug	15-Aug	22-Aug	29-Aug	2-Sep	7-Oct	25-Oct	22-Nov	23-Jan
Nitrate	3.89	4.99	3.67	3.88	4.49	5.46	10.05	9.18	9.13	8.87
Phosphate	0.45	0.48	0.53	0.77	0.47	0.43	0.6	0.45	0.35	0.17
Chromate	0.03	0.05	0	0.13	0.07	0.1	0.13	0.06	0.47	0.14
Iron	0.51	0.05	0	0.13	0	0	0	0	2.36	0
Copper	0	0.09	0.03	0.03	0	0	0.49	0.14	0.21	0.76
Lead	-	0.16	0.27	0.158	0.176	0.121	0.123	0.093	0.027	0.039
Temperature	82 °F	84 °F	62 °F	76 °F	76 °F	90 °F	60 °F	66 °F	43 °F	40 °F

## 2. Materials and Methods

For this investigation, three 1L water samples from Whale Creek, which is part of Newtown Creek, by the Newtown Creek Nature Walk were collected and analysed the same day, except for the January 23 sample (3 days later).

The samples were tested for phosphate, nitrate, dissolved iron, chromate, and copper using the CHEMetrics instrumental kits and the CHEMetrics V-2000 Multi-Analyte Photometer. Lead was determined with the Lead TNTplus kit offered by HACH. The samples underwent a predigestion with the Metals Prep Set TNTplus kit. Samples were analysed using the UV-spectrophotometer, DR5000 by HACH.

## 3. Results and Discussion

The U.S. EPA Maximum Contaminant Level (MCL) for drinking water for nitrate ( $\text{NO}_3^-$ ) is 10 ppm [6]. The levels recorded from August to January were either below or about 10 ppm (Figure 2). High levels of nitrate consumed can lead to difficulties absorbing oxygen into the blood, and high levels of nitrate in a river can lead to large algae growth.

The U.S. EPA MCL for drinking water for copper is 1.3 ppm [6]. Copper was well below the limit (Figure 3).

British Columbia published a report for the protection of aquatic life that concluded that the maximum level for dissolved iron was 0.35 ppm [7]. In July (0.51 ppm) and November (2.36 ppm), the levels were above this threshold (Figure 4). The rest of the time, the levels were below.

The U.S. EPA MCL for drinking water for total Chromium is 0.1 ppm [6]. In late summer, August 22, and early fall, Oct. 7, the chromate ( $\text{CrO}_4^{2-}$ ) level (not total chromium) was slightly above at 0.13 ppm. In November, the level dramatically increased to 0.47 ppm. Iron also dramatically increased in November (Figure 5).

One source said that the EPA set the limit for rivers not flowing into a reservoir or lake to be 0.1 ppm phosphate [8]. This was in order to limit algae blooms. Unfortunately, our results found the phosphate levels to be much higher from 0.17 to 0.77 ppm (Figure 6).

The U.S. EPA MCL for drinking water for lead is 0.015 ppm [6]. From August to October, the levels were at least 10 times greater. Starting at the end of October, the lead levels began to decrease (Figure 7). As the months progressed the air temperature went down. Perhaps the lead is leaching out of the pipes and the warmer weather increases this effect.

## 4. Conclusions

Over 6 months, nitrate, phosphate, copper, iron, chromate, and lead were measured. The lead levels were at least ten

times greater than the MCL for lead, and diminished greatly in November. Perhaps the lead is being leached from pipes and the cooler temperatures mitigate this process. In addition the chromate level and iron levels spiked in November. Perhaps this was due to some form of pollution being added to the creek during that time period. We plan to continue to monitor these toxins at Newtown Creek.

## ACKNOWLEDGEMENTS

The authors gratefully acknowledge the support and funding by the City University of New York (CUNY) Research Scholars Program.

## REFERENCES

- [1] Hurley, A., 1994, Creating ecological wastelands oil pollution in New York City, 1870-1900. *Journal of Urban History*. 20, 340-363.
- [2] Navarro, M. (2010) Between Queens and Brooklyn, an Oil Spills Legacy. *New York Times*. [Online]. Available: <http://www.nytimes.com/2010/08/04/science/earth/04newtown.html?hp>.
- [3] Smith, P. A. (2013) Liquid Cow and Black Mayo. *New York Magazine*. [Online]. Available: <http://nymag.com/news/intelligencer/newtown-creek-2013-12/>.
- [4] U.S. Environmental Protection Agency. (2015) Newtown Creek. Retrieved March 17, 2015, from <http://www.epa.gov/region2/superfund/npl/newtowncreek/>.
- [5] Durkin, E. (2009) Exxon Mobil, BP, National Grid and Phelps Dodge to pay for Newtown Creek cleanup phase. *NY Daily News*. [Online] Available: <http://www.nydailynews.com/new-york/brooklyn/exxon-mobil-chevron-bp-national-grid-phelps-dodge-pay-newtown-creek-cleanup-phase-article-1.418224>.
- [6] United States Environmental Protection Agency. (2015) Drinking Water Contaminants. Retrieved March 17, 2015, from <http://water.epa.gov/drink/contaminants/>.
- [7] Phippen, B.; Horvath, C.; Nordin, R.; Nagpal, N., 2008 Ambient Water Quality Guidelines for Iron. Recommended Guidelines. 978-0-7726-5989-7 Retrieved March 17, 2015 from [http://www.env.gov.bc.ca/wat/wq/BCguidelines/iron/iron\\_overview.pdf](http://www.env.gov.bc.ca/wat/wq/BCguidelines/iron/iron_overview.pdf).
- [8] Water Sheds (2015) Phosphorus. Retrieved March 17, 2015 from <http://www.water.ncsu.edu/watershedss/info/phos.html>.
- [9] Anchor QEA, LLC (2014) Final Phase 2 Remedial Investigation Work Plan – Volume 1. Retrieved April 2, 2015 from [http://www.epa.gov/region2/superfund/npl/newtowncreek/pdf/phase\\_2\\_ri\\_workplan\\_volume1\\_2014\\_05\\_16.pdf](http://www.epa.gov/region2/superfund/npl/newtowncreek/pdf/phase_2_ri_workplan_volume1_2014_05_16.pdf).