

Spring 1961

## On the Order of a Bias

William (Bill) H. Williams  
*CUNY Hunter College*

[How does access to this work benefit you? Let us know!](#)

Follow this and additional works at: [http://academicworks.cuny.edu/hc\\_pubs](http://academicworks.cuny.edu/hc_pubs)

 Part of the [Physical Sciences and Mathematics Commons](#)

---

### Recommended Citation

Williams, William (Bill) H., "On the Order of a Bias" (1961). *CUNY Academic Works*.  
[http://academicworks.cuny.edu/hc\\_pubs/119](http://academicworks.cuny.edu/hc_pubs/119)

This Article is brought to you for free and open access by the Hunter College at CUNY Academic Works. It has been accepted for inclusion in Publications and Research by an authorized administrator of CUNY Academic Works. For more information, please contact [AcademicWorks@cuny.edu](mailto:AcademicWorks@cuny.edu).



---

On the Order of a Bias

Author(s): W. H. Williams

Source: *The American Mathematical Monthly*, Vol. 68, No. 2 (Feb., 1961), pp. 137-138

Published by: Mathematical Association of America

Stable URL: <http://www.jstor.org/stable/2312477>

Accessed: 03-12-2015 19:30 UTC

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



*Mathematical Association of America* is collaborating with JSTOR to digitize, preserve and extend access to *The American Mathematical Monthly*.

<http://www.jstor.org>

ON THE ORDER OF A BIAS

W. H. WILLIAMS,\* McMaster University

A quantity which often arises in statistics is  $u/v$  where  $u = (\bar{x} - \mu_x)^2$ ,  $v = \sum_{i=1}^n (x_i - \bar{x})^2$ , and  $\bar{x}$  is the mean of a sample  $(x_1, \dots, x_n)$  drawn from a population with mean  $\mu_x$ . The form arises, for example, in evaluating the variance of the linear regression estimator of a population mean, see Cochran [1]. It has been shown ([3], [4]) that  $u$  and  $v$  are independently distributed if and only if the  $x$  distribution is Gaussian. Much literature has been devoted to the effects of non-normality on  $u/v$ , most of which is directed at the distribution properties and analysis of Type I and Type II errors. In general  $E(u/v) = E(1/v)Eu + \text{Cov}(u, 1/v)$ , where  $E$  denotes mathematical expectation and  $\text{Cov}$  denotes the covariance of two random variables. If the higher moments are finite, it can be shown by an interesting straightforward method that the covariance term is of lower order in  $n$  than  $E(1/v)Eu$ .

If  $u/v$  is written as  $U(1 + \delta_u)(1 + \delta_v)^{-1}/V$  and  $1/v$  as  $(1 + \delta_v)^{-1}/V$  where  $Eu = U$ ,  $Ev = V \neq 0$ ,  $\delta_u = (u - U)/U$  and  $\delta_v = (v - V)/V$  then  $\text{Cov}(u, 1/v) = (U/V)E\{\delta_u(1 + \delta_v)^{-1}\} \doteq -\text{Cov}(u, v)/V^2$ . The approximation involves dropping the terms  $UE\{\delta_u\delta_v^{2+i}\}/V$ ,  $i = 0, 1, \dots$ . The terms inside the expectation cannot be greater than  $O(n^{-1})$  by the methods of Fisher [2]. Further,  $U = \sigma_x^2/n$  and  $V = (n-1)\sigma_x^2$  so that the neglected terms are no more than  $O(n^{-3})$ .

Next, let  $d_i = x_i - \mu_x$  so that  $Ed_i = 0$  and  $Ed_i^2 = \sigma_x^2$ , then

$$\text{Cov}(u, v) = \frac{1}{n^3} E \left\{ \left[ (n-1) \sum_{i=1}^n d_i^2 - \sum_{i \neq j} d_i d_j \right] \left[ \sum_{i=1}^n d_i^2 + \sum_{i \neq j} d_i d_j \right] \right\} - \frac{n-1}{n} \sigma_x^4.$$

which will reduce to

$$(1) \quad \frac{n-1}{n^2} \{Ed_i^4 - 3\sigma_x^4\}$$

\* Now at Bell Telephone Laboratories, Inc.

or alternatively, since by definition the variance of  $d_i^2$  is  $\text{Var}(d_i^2) = E d_i^4 - \sigma_x^4$ ,

$$(2) \quad \frac{n-1}{n^2} \{ \text{Var}(d_i^2) - 2\sigma_x^4 \}.$$

Hence,  $\text{Cov}(u, v)$  is  $O(n^{-1})$  and  $\text{Cov}(u, 1/v)$  is  $O(n^{-3})$ . Furthermore,  $Eu$  is  $O(n^{-1})$  and  $E(1/v)$  is  $O(n^{-1})$  so that  $EuE(1/v)$  is  $O(n^{-2})$ . Notice the interesting agreement of (1) and (2) with the known fact that  $\text{Cov}(u, v) = 0$  for normal populations. The straightforward approach used to derive these expressions can also be used to verify the stated orders of the terms neglected in the approximation.

#### References

1. W. G. Cochran, *Sampling Techniques*, New York, 1953.
2. R. A. Fisher, Moments and product moments of sampling distributions, *Proc. London Math. Soc.*, vol. 30, 1928, pp. 199–238.
3. R. C. Geary, Distribution of students ratio for non-normal samples, *J. Royal Statist. Soc.*, vol. 3, 1936, pp. 178–184 (Supplement).
4. E. Lukács, A characterization of the normal distribution, *Ann. Math. Statist.*, vol. 13, 1942, pp. 91–93.