

6-1-2010

Newborn infants learn during sleep

William P. Fifer
Columbia University

Dana L. Byrd
University of Florida

Michelle Kaku
Columbia University

Inge-Marie Eigsti
University of Connecticut - Storrs

Joseph R. Isler
Columbia University

See next page for additional authors

[How does access to this work benefit you? Let us know!](#)

Follow this and additional works at: https://academicworks.cuny.edu/jj_pubs

 Part of the [Developmental Psychology Commons](#)

Recommended Citation

Fifer, William P.; Byrd, Dana L.; Kaku, Michelle; Eigsti, Inge-Marie; Isler, Joseph R.; Grose-Fifer, Jillian; Tarullo, Amanda R.; and Balsam, Peter D., "Newborn infants learn during sleep" (2010). *CUNY Academic Works*.
https://academicworks.cuny.edu/jj_pubs/146

This Article is brought to you for free and open access by the John Jay College of Criminal Justice at CUNY Academic Works. It has been accepted for inclusion in Publications and Research by an authorized administrator of CUNY Academic Works. For more information, please contact AcademicWorks@cuny.edu.

Authors

William P. Fifer, Dana L. Byrd, Michelle Kaku, Inge-Marie Eigsti, Joseph R. Isler, Jillian Grose-Fifer, Amanda R. Tarullo, and Peter D. Balsam

Newborn infants learn during sleep

William P. Fifer^{a,b,c,1}, Dana L. Byrd^d, Michelle Kaku^a, Inge-Marie Eigsti^e, Joseph R. Isler^b, Jillian Grose-Fifer^f, Amanda R. Tarullo^a, and Peter D. Balsam^{a,g}

^aDepartment of Psychiatry, Columbia University, New York, NY, 10032; ^bDepartment of Pediatrics, Columbia University, New York, NY 10032; ^cNew York State Psychiatric Institute, New York, NY 10032; ^dDepartment of Psychology, University of Florida, Gainesville, FL 32611; ^eDepartment of Psychology, University of Connecticut, Storrs, CT 06269; ^fDepartment of Psychology, John Jay College of Criminal Justice, City University of New York, New York, NY 10019; and ^gDepartment of Psychology, Barnard College, Columbia University, New York, NY 10027

Communicated by Eric Richard Kandel, Columbia University, New York, NY, April 16, 2010 (received for review November 19, 2009)

Newborn infants must rapidly adjust their physiology and behavior to the specific demands of the novel postnatal environment. This adaptation depends, at least in part, on the infant's ability to learn from experiences. We report here that infants exhibit learning even while asleep. Bioelectrical activity from face and scalp electrodes was recorded from neonates during an eye movement conditioning procedure in which a tone was followed by a puff of air to the eye. Sleeping newborns rapidly learned the predictive relationship between the tone and the puff. Additionally, in the latter part of training, these infants exhibited a frontally maximum positive EEG slow wave possibly reflecting memory updating. As newborns spend most of their time sleeping, the ability to learn about external stimuli in the postnatal environment during non-awake states may be crucial for rapid adaptation and infant survival. Furthermore, because eyelid conditioning reflects functional cerebellar circuitry, this method potentially offers a unique approach for early identification of infants at risk for a range of developmental disorders including autism and dyslexia.

EEG | eyelid conditioning | neonate

During the first days of life, awake infants are capable of learning associations between oral motor patterns and altered milk flow (1) and can learn to alter sucking to produce a variety of reinforcers, including milk (2), their mother's voice (3, 4), or a sweet-tasting solution (5). Cross-sensory associative learning also has been demonstrated in awake neonates using paired auditory and visual stimuli (6, 7). Furthermore, awake newborns show Pavlovian conditioning to tactile (8) and taste stimuli (9, 10), as well as eyelid conditioning to paired auditory and tactile stimuli (11). These early adaptations to the postnatal environment have been well documented in awake infants, but as newborns spend the vast majority of their time asleep, the need and capacity to learn may not be confined to states of wakefulness.

Even while asleep, neonates are able to process external information actively. Scalp recordings of brain activity in sleeping neonates have demonstrated their capacity to differentiate between two sounds (12–14), indicating that infants are forming representations of specific stimuli and distinguishing between those stimuli during sleep. We report here that sleeping neonates not only process information about individual events, but also learn about relationships between them.

To investigate whether neonates can learn during sleep, we attempted to condition an eye movement response in 1- to 2-day-old infants while they slept. All infants were fed immediately before testing to increase the likelihood they would sleep through the entire procedure. Sleep status was confirmed using behavioral observations in conjunction with heart rate variability patterns, respiratory regularity, and video scoring of the infants' faces. Infants were videotaped while exposed to tones and puffs of air directed at the eyelid. In the experimental group, tones were a reliable signal that a puff of air was likely to be presented. In the control group, tones and puffs were presented at random times. In addition to video scored observational measures derived from previous studies with older awake infants (15, 16), electroencephalographic and muscle activity was also recorded

from a network of 124 scalp and face electrodes. Responses associated with both neural processing and eye movements to the tone and air puff were recorded and analyzed over the course of a single training session.

Results

In the experimental group, participants averaged a 4-fold increase in the likelihood of the conditioned eye movement response (EMR) by the end of training, whereas the likelihood of responding did not change in the control group (Fig. 1*B*). Statistical analyses were conducted on data aggregated into five blocks of 40 trials. A two-way analysis of variance showed a significant difference between groups, $F(1, 28) = 38.17, P < 0.05$, and a significant group- (experimental vs. control) by-block interaction, $F(4, 112) = 8.41, P < 0.05$. Separate ANOVAs for the two groups showed a significant increase in eyelid responses over trials in the experimental group, $F(4, 100) = 42.14, P < 0.05$, but not for the controls. The group average onset latency of eye movement responses to the tone on tone-alone trials decreased from 1,030 ms in the first block of the session to 870 ms in the final block, becoming significantly closer to the expected puff onset at 900 ms, $t(25) = 5.40, P < 0.005$. By the end of training, 24 of the 26 babies in the experimental group had, at a minimum, doubled the likelihood of making an EMR. In contrast, there was no change in the probability of an EMR in the control group (Fig. 1*B*). Because $\approx 92\%$ of the subjects in the experimental group conditioned, the chance of having 4 of 4 subjects that fail to condition (as in the control group) is < 0.00005 . Hence, 4 subjects provide an adequate control for this behavioral paradigm. For the total sample, EMR was observed on $> 90\%$ of puff-alone trials across all blocks, demonstrating that infants consistently were responsive to the unconditioned stimulus. There was no change in rate of unconditioned responding from the first to the second half of the session, $t(23) = 1.09$ (not significant).

Because the auditory evoked potential is relatively small in comparison with the ongoing EEG, at least 50 trials were required to extract a clear sensory evoked potential using averaging. Ten infants in the experimental group, but only 1 in the control group, had sufficient artifact-free EEG responses required for event-related potential (ERP) analysis. Consequently, statistical analysis was possible only in the experimental group. In this subset, separate averages were constructed for each infant for the first and the second half of the conditioning trials. Consistent with previous studies of newborns (17–19), these babies had clearly visible auditory evoked potentials (AEPs) with a positive peak at ≈ 300 ms (Fig. 2). The AEP was maximal in amplitude (3.5 μ V) over the central areas. Visual inspection of

Author contributions: W.P.F., D.L.B., I.-M.E., and P.D.B. designed research; W.P.F., D.L.B., and M.K. performed research; W.P.F., D.L.B., M.K., J.R.I., J.G.-F., A.R.T., and P.D.B. analyzed data; and W.P.F., D.L.B., M.K., I.-M.E., J.R.I., J.G.-F., A.R.T., and P.D.B. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: wpf1@columbia.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1005061107/-DCSupplemental.

spending most of the time cycling within sleep states. Thus, the perinatal capacity to learn while asleep may play a pivotal role in the infant's rapid adaptation to the postnatal environment.

Eyelid conditioning relies on an intact functioning cerebellum (32), and functional MRI (fMRI) shows cerebellar activation during delay eyelid conditioning in adults (33). Thus, the current study demonstrates the feasibility of using eyelid conditioning as an early screen for cerebellar function, which may help identify infants at risk for several neurodevelopmental disorders. Functional cerebellar abnormalities are implicated in dyslexia, attention deficit hyperactivity disorder, autism, and schizophrenia (34, 35), and deficits in cerebellar mediated eyelid conditioning are associated with all of these disorders (36–41). Cerebellar abnormalities in neurodevelopmental disorders such as autism are thought to emerge prenatally (42). Moreover, the third trimester of pregnancy is a period of rapid cerebellar development, and preterm birth disrupts this process (43, 44). Because disruption of cerebellar development interferes with acquisition of the conditioned eye movement response (45), the conditioning paradigm described here offers a unique approach to non-invasive cerebellar-based assessment of developmental risk in newborn infants. Additionally, whereas delay eyelid conditioning does not necessarily depend on the cerebral cortex (46), changes in frontal activity associated with conditioning may provide early information about cortical function, as well.

Materials and Methods

Thirty-four healthy term infants between 10 and 73 h of age were tested while asleep. After being fitted with cardio-respiratory sensors and a high-density EEG net, the infant was swaddled and placed supine in a bassinet. Tones were presented using two speakers, each placed next to the ears. A flexible tube was configured to deliver a puff of air to the outer canthus of the right eye (Fig. 1A).

Usable eye movement response data (without waking or crying or unacceptable amounts of electrical noise) were obtained from 30 infants (14 males and 16 females). Immediately following 3 tone-alone trials, 20 sets of 10

trials were presented to the experimental group ($n = 26$). Each set of 10 trials consisted of eight tones paired with an air puff, 1 puff-alone trial, and 1 tone-alone trial. The puff-alone trials allow the unconditioned response to be monitored over the course of training without the influence of a prior tone. The tone-alone trials allowed the detection of conditioned responses unaffected by the presentation of an air puff on paired trials. The control group ($n = 4$) was presented with the same number of stimuli as the experimental group in an equivalent amount of time but in a semirandom sequence.

The EMRs served as the primary measure of conditioned and unconditioned responses. Offline, the EMRs from four frontal electrodes along the eyebrow ridge were visually inspected during the 2000 ms after the conditioned stimulus onset on tone-alone trials and scored for the occurrence of a sharply rising electrical potential (20–200 μV) in the midline lead nearest the eye (E17), indicative of muscle activation associated with eye movement (47). The onset latency of the EMR also was scored for tone-alone trials. For puff-alone trials, EMR was scored during the 1000 ms after puff onset.

EEG activity was recorded using a high impedance system with 124 electrodes spaced at 1-cm intervals over the scalp, using a vertex reference. Data were sampled at 1,000 Hz. All analyses were performed offline using a linked mastoid reference. The EEG in response to all tone stimuli (i.e., tone-puff pairs and tone-alone stimuli) was analyzed from 128 ms before stimulus presentation until 900 ms after tone onset (just before puff presentation on paired trials). Electrodes with artifact-free data for at least 80% of the epochs were included in the analysis (*SI Materials and Methods*). We examined if patterns of brain activation associated with early sensory or later cognitive processing (13, 21) changed over training. Because these patterns of activation are very small in relation to the background EEG, many responses need to be averaged to extract the stimulus-specific response from ongoing brain activity. Two averaged EEG waveforms (event-related potentials) were created for each infant on the basis of the first and second halves of the trials for both sensory and cognitive responses to the tone.

ACKNOWLEDGMENTS. We thank Elizabeth Alf, Carmen Gennarini, Michael M. Myers, J. David Nugent, Michele Pasamba, Albany Perez, Kimon Violaris, and Jessica Wilson for help with many aspects of this project. This research was supported by the Sackler Institute of Developmental Psychobiology at Columbia University and by National Institutes of Health Grants R37 HD032774 (to W.P.F.), T32 MH018264 (to D.L.B., I.-M.E., and A.R.T.), and R01 MH068073 (to P.D.B.).

- Bosma JF (1972) *Oral Sensation and Perception: The Mouth of the Infant* (Charles C. Thomas, Springfield, IL).
- Papoušek H (1961) Conditioned head rotation reflexes in infants in the first months of life. *Acta Paediatr* 50:565–576.
- DeCasper AJ, Fifer WP (1980) Of human bonding: Newborns prefer their mothers' voices. *Science* 208:1174–1176.
- Moon C, Bever TG, Fifer WP (1992) Canonical and non-canonical syllable discrimination by two-day-old infants. *J Child Lang* 19:1–17.
- Siqueland ER, Lipsitt LP (1966) Conditioned head-turning in human newborns. *J Exp Child Psychol* 3:356–376.
- Crowell DH, Blurton LB, Kobayash LR, McFarland JL, Yang RK (1976) Studies in early infant learning: Classical conditioning of the neonatal heart rate. *Dev Psychol* 12:373–397.
- Stamps LE, Porges SW (1975) Heart rate conditioning in newborn infants: Relationships among conditionability, heart rate variability, and sex. *Dev Psychol* 11:424–431.
- Sullivan RM, et al. (1991) Olfactory classical conditioning in neonates. *Pediatrics* 87:511–518.
- Blass EM, Ganchrow JR, Steiner JE (1984) Classical conditioning in newborn humans 2–48 hours of age. *Infant Behav Dev* 7:223–235.
- Clifton RK (1974) Heart rate conditioning in the newborn infant. *J Exp Child Psychol* 18:9–21.
- Little AH, Lipsitt LP, Rovee-Collier C (1984) Classical conditioning and retention of the infant's eyelid response: Effects of age and interstimulus interval. *J Exp Child Psychol* 37:512–524.
- Cheour M, et al. (2002) Speech sounds learned by sleeping newborns. *Nature* 415:599–600.
- Kurtzberg D, Hilpert PL, Kreuzer JA, Vaughan HG, Jr (1984) Differential maturation of cortical auditory evoked potentials to speech sounds in normal fullterm and very low-birthweight infants. *Dev Med Child Neurol* 26:466–475.
- Sambeth A, Ruohio K, Alku P, Fellman V, Huotilainen M (2008) Sleeping newborns extract prosody from continuous speech. *Clin Neurophysiol* 119:332–341.
- Clafin DI, et al. (2002) Effect of delay interval on classical eyeblink conditioning in 5-month-old human infants. *Dev Psychobiol* 41:329–340.
- Herbert JS, Eckerman CO, Stanton ME (2003) The ontogeny of human learning in delay, long-delay, and trace eyeblink conditioning. *Behav Neurosci* 117:1196–1210.
- Barnet AB, Ohlrich ES, Weiss IP, Shanks B (1975) Auditory evoked potentials during sleep in normal children from ten days to three years of age. *Electroencephalogr Clin Neurophysiol* 39:29–41.
- Rotteveel JJ, Colon EJ, Stegeman DF, Visco YM (1987) The maturation of the central auditory conduction in preterm infants until three months post term. IV. Composite group averages of the cortical auditory evoked responses (ACRs). *Hear Res* 27:85–93.
- Shucard DW, Shucard JL, Thomas DG (1987) Auditory event-related potentials in waking infants and adults: A developmental perspective. *Electroencephalogr Clin Neurophysiol* 68:303–310.
- Courchesne E (1979) From infancy to adulthood: The neurophysiological correlates of cognition. *Clin Neurophysiol* 6:224–242.
- Pascalis O, de Haan M, Nelson CA, de Schonen S (1998) Long-term recognition memory for faces assessed by visual paired comparison in 3- and 6-month-old infants. *J Exp Psychol Learn Mem Cogn* 24:249–260.
- Aarons L (1976) Sleep-assisted instruction. *Psychol Bull* 83:1–40.
- Hensch TK (2004) Critical period regulation. *Annu Rev Neurosci* 27:549–579.
- Scher MS (2008) Ontogeny of EEG-sleep from neonatal through infancy periods. *Sleep Med* 9:615–636.
- Fransson P, et al. (2009) Spontaneous brain activity in the newborn brain during natural sleep—an fMRI study in infants born at full term. *Pediatr Res* 66:301–305.
- Richardson HL, Walker AM, Horne RS (2009) Maternal smoking impairs arousal patterns in sleeping infants. *Sleep* 32:515–521.
- Harper RM (2003) Impaired arousals and sudden infant death syndrome: Preexisting neural injury? *Am J Respir Crit Care Med* 168:1262–1263.
- Kinney HC (2009) Brainstem mechanisms underlying the sudden infant death syndrome: Evidence from human pathologic studies. *Dev Psychobiol* 51:223–233.
- Mitchell EA, Williams SM, Taylor BJ (1999) Use of duvets and the risk of sudden infant death syndrome. *Arch Dis Child* 81:117–119.
- Lipsitt LP (1982) Infancy and life-span development. *Hum Dev* 25:41–48.
- Paluszynska DA, Harris KA, Thach BT (2004) Influence of sleep position experience on ability of prone-sleeping infants to escape from asphyxiating microenvironments by changing head position. *Pediatrics* 114:1634–1639.
- Ohyama T, Nores WL, Murphy M, Mauk MD (2003) What the cerebellum computes. *Trends Neurosci* 26:222–227.
- Cheng DT, Disterhoft JF, Power JM, Ellis DA, Desmond JE (2008) Neural substrates underlying human delay and trace eyeblink conditioning. *Proc Natl Acad Sci USA* 105:8108–8113.
- Rae C, et al. (1998) Metabolic abnormalities in developmental dyslexia detected by ^1H magnetic resonance spectroscopy. *Lancet* 351:1849–1852.
- Hoppenbrouwers SS, Schutter DJ, Fitzgerald PB, Chen R, Daskalakis ZJ (2008) The role of the cerebellum in the pathophysiology and treatment of neuropsychiatric disorders: A review. *Brain Res Rev* 59:185–200.

