

1-2018

Secular Trends in Regional Differences in Nutritional Biomarkers and Self-Reported Dietary Intakes among American Adults: National Health and Nutrition Examination Survey (NHANES) 1988–1994 to 2009–2010

Ashima K. Kant
CUNY Queens College

Barry I. Graubard
National Institutes of Health

[How does access to this work benefit you? Let us know!](#)

Follow this and additional works at: https://academicworks.cuny.edu/qc_pubs

 Part of the [Nutritional Epidemiology Commons](#)

Recommended Citation

Kant, Ashima K. and Barry I. Graubard. "Secular Trends in Regional Differences in Nutritional Biomarkers and Self-Reported Dietary Intakes among American Adults: National Health and Nutrition Examination Survey (NHANES) 1988–1994 to 2009–2010." *Public Health Nutrition*, published online 10 January 2018, doi: 10.1017/S1368980017003743 . Reprinted in CUNY Academic Works.

This Article is brought to you for free and open access by the Queens College at CUNY Academic Works. It has been accepted for inclusion in Publications and Research by an authorized administrator of CUNY Academic Works. For more information, please contact AcademicWorks@cuny.edu.



Secular trends in regional differences in nutritional biomarkers and self-reported dietary intakes among American adults: National Health and Nutrition Examination Survey (NHANES) 1988–1994 to 2009–2010

Ashima K Kant^{1,*} and Barry I Graubard²

¹Queens College of the City University of New York, Remsen Hall, Room 306E, Flushing, NY 11367, USA:

²Biostatistics Branch, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA

Submitted 6 March 2017; Final revision received 13 October 2017; Accepted 14 November 2017

Abstract

Objective: To understand the contribution of regional differentials in dietary exposures to regional gradients in health, we examined 20-year trends in the association of US census region of residence with nutritional biomarkers and dietary intakes of American adults.

Design: Observational.

Setting: The biomarker and 24 h dietary recall data were from the National Health and Nutrition Examination Surveys (NHANES) conducted during 1988–1994 and 1999–2010. The US census region was operationalized as Northeast, Midwest, South and West. Nutritional biomarker outcomes were serum folate, vitamins B₆, B₁₂, C, D and E, and carotenoids; dietary outcomes were intakes of nutrients, food groups and eating patterns.

Subjects: US adults, $n > 8000$ – $40\,000$ for biomarkers and $>43\,000$ for dietary outcomes.

Results: The interactions of survey time period and region were not significant for the examined biomarker and dietary outcomes, indicating similar secular trends among regions. The main effect of region was significant for all nutritional biomarkers except serum vitamin B₆, most dietary micronutrients, food groups and eating patterns ($P < 0.001$). The mean serum folate, vitamins B₁₂, C and E, and all carotenoid (except lycopene) biomarker levels, and intakes of dietary fibre, vitamins A, E, C and B₆, folate, K, Ca, Mg and Fe, fruits, vegetables and whole grains, were higher in the West and Northeast regions, relative to the South and Midwest regions.

Conclusions: Overall, the regional gradients in dietary exposure, expressed objectively as biomarkers or as self-reported nutrient and food group intakes, paralleled trajectories reported for health outcomes and were remarkably persistent over time.

Keywords
Census region
NHANES
Secular trends
Biomarkers
Dietary intakes
Diet quality
Food groups
Eating patterns
Health disparities

There are known regional differences in mortality from all causes and from stroke, heart disease and cancer in the US population^(1–6). The mortality rates are highest in the South, followed by the Midwest, Northeast and the West. Over the past several decades, US mortality rates have declined overall⁽⁷⁾; however, the regional gradients in mortality rates have persisted^(4–6). The likely reasons for these regional health disparities are complex and include social and physical determinants, which also relate to individual risk behaviours⁽⁸⁾. In an attempt to understand the reasons for regional differentials in health, regional differences in established risk factors for leading causes of

morbidity and mortality – obesity, physical inactivity, smoking, hypertension – have been examined in several publications^(9–14). These reports generally show regional gradients in risk factors that are similar to those observed for risk of mortality.

Diet plays a role in both prevention and management of leading causes of mortality^(15,16). Both quantitative and qualitative dietary characteristics may potentially modify disease risk and risk factors. Given the large body of investigation of diet and health associations, surprisingly few published studies have systematically addressed regional variation in dietary attributes^(17–22); fewer still

*Corresponding author: Email ashima.kant@qc.cuny.edu



have examined regional differences in secular trends in dietary intakes⁽¹⁴⁾. Because self-reported dietary intake of free-living individuals is measured with error⁽²³⁾, biomarkers of dietary exposure provide an alternative to self-reported intakes⁽²⁴⁾. Circulating micronutrient concentrations also reflect nutrient bioavailability, which reflects the mix of dietary constituents and supplements consumed and individual factors⁽²⁴⁾. To our knowledge, however, there is little published information on regional differences in key nutritional biomarkers of dietary exposure even though nutrient biomarkers have the advantage of objectivity. With few exceptions, the diets of free-living individuals contain multiple sources of nutrients. For further understanding of dietary aspects that require intervention, it is important to examine dietary intakes and biomarkers concurrently, but none such studies are currently available. Given these gaps, we used nationally representative data to examine secular trends (1988–2010) in differences among US census regions in nutritional biomarkers and self-reported dietary intakes.

Methods

We used public-domain nutritional and disease biomarker data⁽²⁵⁾ and restricted geographic data⁽²⁶⁾ for the National Health and Nutrition Examination Survey (NHANES) 1988–1994 and the continuous NHANES 1999–2010. The City University of New York, Human Research Protection Program, did not consider the study human subjects research. The NHANES surveys, conducted by the National Center for Health Statistics (NCHS) of the Centers for Disease Control and Prevention, include a stratified, multistage probability sampling design to select a nationally representative sample of the US non-institutionalized population. The surveys include an at-home interview of each sampled person and a complete medical examination in a specially equipped mobile examination centre (MEC). The MEC visit includes a dietary interview and collection of anthropometric and biochemical measures. The response rates for samples examined in the MEC in these surveys were >70%⁽²⁷⁾.

Exposure assessment (geographic variables)

Information on census region and urbanization of area of residence of the NHANES survey participants is not available in the public domain. Therefore, public-domain data were merged with the available restricted-access census region and urbanization of area of residence data by the Research Data Center (RDC) of the NCHS. The census region was operationalized as Northeast, Midwest, South and West according to the US Census Bureau definition⁽²⁸⁾. Because the NHANES are not designed to provide census division, state or county-level data, this level of geographic aggregation is therefore the only one practicable. There are documented differences in access to

healthy foods in rural *v.* urban areas⁽¹¹⁾; therefore, the urbanization of area of residence was a covariate in our analysis and was operationalized as (i) urban, large metro, (ii) urban, fringe metro, (iii) urban, medium/small metro and (iv) non-metropolitan, according to the NCHS urban/rural classification⁽²⁹⁾. Urban/rural linkage of NHANES respondents is available only in the restricted-access data for use at the RDC.

Biomarker outcomes

Serum total and HDL cholesterol, C-reactive protein, folate (both serum and red blood cell (RBC)), carotenoids, and vitamins B₆, B₁₂, C, D and E were examined as outcomes. The biomarkers examined include those that reflect chronic disease risk and nutrient exposures. Each biomarker was not measured in all examined surveys. The methods of assay for each biomarker and the survey years in which the biomarker was measured are described in the documentation available for each survey⁽²⁵⁾, and are summarized in the Appendix. Per NCHS recommendations⁽²⁵⁾, for serum and RBC folate and serum 25-hydroxy-cholecalciferol, we used data harmonized using calibration to adjust for differences in assay methodology over the surveys examined. For vitamins B₆ and C, changes in assay methods across survey years were not harmonized and made it inadvisable to examine trends; in such cases we followed NCHS guidelines in using the data from later surveys⁽²⁵⁾.

Dietary outcomes

Each NHANES used computer-assisted 24 h recall methodology to collect an in-person recall⁽²⁵⁾. Trained dietary interviewers administered the 24 h recall during the MEC exam⁽²⁵⁾. Estimated intakes of energy and nutrients available in the public-domain dietary data for each sample person were derived from the 24 h recall⁽²⁵⁾. We used these data to examine intakes of energy, percentage of energy from macronutrients, dietary fibre, the micronutrients vitamins A, B₆, B₁₂, folate, C and E, and the minerals Na, K, Ca, Mg and Fe. The 2015 Dietary Guidelines for Americans Advisory Committee has identified these nutrients as either shortfall or of public health concern⁽¹⁶⁾. As an approximate indicator of energy under-reporting⁽³⁰⁾, we also computed the ratio of reported 24 h intake to estimated basal energy expenditure for each respondent in all surveys.

The 24 h recall of dietary intake has also been used to create a Pyramid servings database (NHANES 1988–1994), a MyPyramid Equivalents Database (MPED; NHANES 1999–2004) or a Food Patterns Equivalents Database (FPED; NHANES 2005–2010) for each respondent, by the US Department of Agriculture^(31,32). We used these data to obtain summary estimates of serving equivalents of total fruits, whole fruits, vegetables, dairy, grains, whole grains, meat, added sugar, discretionary solid fat, and combined energy from of discretionary fat, added sugar and alcohol.



These food groups represent those encouraged and those discouraged by current and past dietary guidance⁽¹⁶⁾. There is also increasing interest in the potential role of eating patterns in health promotion⁽³³⁾. Information on the name of each eating episode reported and time of its consumption was also obtained during the recall. Using previously described methods^(34,35), we used this information to derive eating pattern outcomes of number of main meal or non-main meal eating episodes, breakfast report in the recall, time interval between the first and the last eating episode of the day and interval between eating episodes.

Analytic sample

Biomarker outcomes

In each NHANES survey, respondents aged ≥ 20 years with information on at least one biomarker of interest were eligible for inclusion in the analytic sample (n 45 155). We excluded pregnant and lactating women (n 1710), and those missing census region information (n 15), for a total analytic sample of 43 430. However, numbers available for the various biomarkers differ depending on the survey cycle in which the biomarker was measured (see Appendix).

Dietary outcomes

All respondents aged ≥ 20 years with a reliable 24 h recall in the NHANES 1988–1994 and 1999–2010 were eligible for inclusion in the study (n 45 112). We excluded pregnant and lactating women (n 1748), respondents reporting no energy intake in the recall (n 4) and those with missing information on region of residence (n 5). The final analytic sample included 43 345 men and women. Final analytic data sets were created by merge of the restricted-access geographic data with the public-domain dietary, biomarker and covariate data at the RDC of the NCHS.

Analytic approach

Descriptive analysis

We describe the sample-weighted percentage and associated 95 % CI for respondents in four US census regions by various sociodemographic and lifestyle characteristics separately for the analytic samples that examined biomarker and dietary outcomes.

Hypothesis testing

We used regression methods to examine regional differences in time trends in each available biomarker or dietary variable as a continuous outcome. We followed NCHS analytic guidelines⁽³⁶⁾ and combined survey years from the continuous NHANES to increase reliability of results. The survey years combined differed based on availability of the measured biomarker in the survey years (see Appendix).

Covariates

Because residents of different US regions sampled across time may differ in sociodemographic and lifestyle characteristics

with putative associations with the outcomes examined in the present study, we operationalized several covariates for use in multiple linear regression models. Information on these covariates was available in the NHANES public-domain or restricted-access database. For both dietary and biomarker outcomes, the covariates included: mid-survey year, urbanization of area of residence, sex, age, race/ethnicity, years of education, family income as a ratio of income to poverty threshold (PIR), BMI (kg/m^2), self-reported chronic disease status and calendar month period of the MEC exam. For biomarker outcomes, additional covariates included: reported dietary supplement use, serum cotinine (a marker of tobacco exposure) and hours of fasting before phlebotomy. For serum vitamin E and carotenoids as outcomes, serum cholesterol, and for total and HDL cholesterol outcomes, alcohol drinking status were additional covariates. Models for all dietary nutrient and food group outcomes also included total energy intake.

To test whether the relationship of region with biomarker concentrations or dietary intakes changed across survey years, we included a region \times survey interaction term in multiple covariate-adjusted linear regression models. In the absence of significant interactions, we excluded the interaction term and examined the main effect of region for each outcome. In all hypothesis testing, combined survey years were operationalized both as a categorical and as a trend variable. For all outcomes available in three combined survey groups, we used midpoint of the combined survey years as a trend variable. Trends could be examined for all dietary outcomes, all biomarkers except carotenoids, and vitamins B₆ and C. All biomarkers except total or HDL cholesterol and vitamin C were log transformed due to skewness prior to analyses; for these transformed variables, we present back-transformed geometric means in results tables.

The analyses were conducted on-site at the RDC, using the statistical software package SAS version 9.2 and SAS-callable SUDAAN software version 11.0.1, and included the appropriate sample weights to adjust for survey non-response and probability of selection^(36,37). The results presented in tables are geometric (biomarkers) or arithmetic means computed as adjusted means (predicted margins)⁽³⁸⁾, 95 % CI and P values associated with the Wald F test for each hypothesis. This global test was used to determine statistically significant interactions and main effects. In the case of a significant main effect of region, the narrative of results compares the 95 % CI for region-specific adjusted means to identify the general direction of differences of biomarker and dietary variables between regions.

We do not present main effect of survey for several reasons. First, the primary focus of our study is regional differences in dietary and biomarker outcomes. Second, changes in assay methods or laboratory sites for biomarkers, and recall methodology for dietary intakes are confounded with survey effect; however, the methods in each survey were similar for all regions. Moreover, other reports,

including a systematic examination by the NCHS, have examined secular trends in nutritional biomarkers and dietary intakes, thus the information is not novel^(39–41).

Effect modification

In separate models, we tested the three-way interactions of region \times survey \times sex, region \times survey \times age, region \times survey \times PIR, and region \times survey \times race/ethnicity to examine whether any variation in secular trend of each biomarker or dietary outcome across census regions was modified by sex, age, income and ethnicity. If a three-way interaction term was significant, we then examined secular trends stratified by the sociodemographic variable.

Adjustment for multiplicity of hypothesis testing

Given the variety of biomarker and dietary outcomes examined in the present study, we used the Bonferroni correction to adjust for multiple comparisons, based on the number of variables in each table and the hypotheses tested. For biomarker results presented in Table 2, the Bonferroni-corrected *P* value was ≤ 0.001 (for thirteen biomarkers and two tests of hypotheses, and two biomarkers with one hypothesis test each, $0.05/28 = 0.0018$). For dietary energy and nutrient results presented in Table 3, the Bonferroni-corrected *P* value was ≤ 0.001 (for twenty-one dietary outcomes in the table and two tests of hypothesis each, $0.05/42 = 0.0012$). For dietary food group, composition and diet quality results presented in Table 4, the Bonferroni-corrected *P* value was ≤ 0.001 (for eighteen dietary outcomes in the table and two tests of hypothesis each, $0.05/36 = 0.0014$). For eating behaviour results presented in Table 5, the Bonferroni-corrected *P* value was ≤ 0.005 (for five eating behaviour outcomes in the table and two tests of hypothesis each, $0.05/10 = 0.005$).

For tests of effect modification for biomarker and dietary outcomes examined in Tables 2–5, the Bonferroni-corrected values were determined as described above, except that four hypotheses tested three-way interactions (region \times survey \times sex, region \times survey \times age, region \times survey \times race/ethnicity, region \times survey \times PIR) for each variable. For outcomes in Tables 2, 3, 4 and 5, respectively, the corresponding Bonferroni-corrected *P* values for these tests were 0.001, 0.0006, 0.0007 and 0.002. We present the actual *P* values for the global tests in results Tables 2–5, but the identification of statistically significant interactions and main effects of region in the results narrative is based on Bonferroni-corrected *P* values.

Results

Sociodemographic and lifestyle characteristics of respondents

Expectedly, the sociodemographic profile of respondents in each census region reveals differences in distribution of race/ethnicity, PIR of $<130\%$, season of MEC examination,

BMI of $\geq 30 \text{ kg/m}^2$, self-reported chronic disease, education of <12 years, urbanization of residence and supplement use (Table 1 for biomarker analytic sample and online supplementary material, Supplemental Table 1 for the diet analytic sample). Highest percentage residing in large metropolitan areas was from the West, while percentage in the non-metropolitan areas was higher in the Midwest and the South. The Midwest had the highest percentage of non-Hispanic Whites, the South region had the highest percentage of non-Hispanic Blacks; the highest percentages of Mexican-Americans and Others were in the West region. The South region had higher percentage with <12 years of education and PIR of $<130\%$. More of the respondents from the South and the West regions were examined from November to April. The lowest percentage of respondents with BMI of $\geq 30 \text{ kg/m}^2$ or of current smokers or with any self-reported chronic disease was in the West region.

Biomarker outcomes

Did the association of region with each biomarker change over time?

The weighted, unadjusted mean and 95% CI of each biomarker concentration, by region, by survey year, is shown in the online supplementary material, Supplemental Table 2. In covariate-adjusted models, the interaction term for region \times survey did not reach the Bonferroni-corrected level of significance (≤ 0.001) for any of the examined biomarkers (Table 2). For serum vitamins C and B₆, the interaction was not examined because only one combined survey period was available.

Did biomarker concentrations, averaged over surveys, differ among US census regions?

In the absence of significant region \times survey interactions, we examined the main effect of census region for each biomarker (Table 2). The main effect of region across surveys was not significant for serum total cholesterol and vitamin B₆ ($P > 0.001$). For all remaining biomarkers, there were significant differences among regions ($P < 0.001$). The highest mean concentrations of serum (and RBC) folate, vitamins B₁₂, C and E, and the carotenoids α -carotene, β -carotene, β -cryptoxanthin and lutein plus zeaxanthin were in the West and/or the Northeast regions, while the lowest concentrations were in the South region. The mean HDL cholesterol concentration was lower, but C-reactive protein concentration was higher, in the South region. The Midwest region had the highest mean serum vitamin D concentration; both the South and the West had lower concentration than the Midwest.

Effect modification testing for biomarker outcomes

For serum β -carotene concentration, the three-way interaction of region \times survey \times age was significant ($P < 0.0002$). In age-stratified analysis, regional differences in serum

Table 1 Characteristics of adult respondents (weighted percentages and 95% CI) surveyed in the National Health and Nutrition Examination Survey (NHANES) 1988–1994 to 2009–2010, by US census region

	All regions (N 43 430)		Northeast (N 6685)		Midwest (N 8824)		South (N 17 309)		West (N 10 612)	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI
Survey cycle										
NHANES 1988–1994	30.7	28.9, 32.5	34.9	30.0, 40.0	30.5	25.9, 35.6	29.4	26.1, 32.9	29.3	21.5, 38.6
NHANES 1999–2004	33.3	31.6, 35.1	30.7	23.9, 38.4	31.4	23.5, 40.4	36.3	31.7, 41.2	32.8	24.7, 42.2
NHANES 2005–2010	36.0	34.1, 37.9	34.5	28.6, 40.8	38.1	31.4, 45.2	34.3	30.7, 38.0	37.8	30.2, 46.1
Urbanization of place of residence										
Large metropolitan	29.5	25.4, 33.8	23.4	13.8, 36.9	20.0	14.7, 26.5	27.0	19.4, 36.1	48.9	37.1, 60.8
Fringe metropolitan	18.8	14.5, 24.1	32.2	19.7, 47.8	16.2	8.0, 30.2	19.8	13.3, 28.4	9.2	4.6, 17.7
Medium/small metropolitan	33.5	28.2, 39.3	39.4	27.4, 53.0	34.8	22.7, 49.3	29.6	22.3, 38.2	33.5	22.3, 47.0
Non-metropolitan	18.2	15.2, 21.5	4.9	1.0, 21.0	28.9	21.2, 38.0	23.6	18.5, 29.7	8.4	3.3, 19.6
Sex										
Men	49.0	48.6, 49.5	48.9	48.2, 49.7	49.1	48.2, 50.1	49.1	48.3, 49.9	48.9	47.8, 50.0
Women	51.0	50.5, 51.4	51.1	50.0, 51.8	50.9	49.9, 51.8	50.9	50.1, 51.7	51.1	50.0, 52.2
Race/ethnicity										
Non-Hispanic White	73.1	71.1, 74.9	79.1	73.6, 83.7	84.8	82.0, 87.3	66.7	62.7, 70.4	65.4	60.9, 69.5
Non-Hispanic Black	10.6	9.7, 11.6	9.0	6.7, 12.0	8.6	6.9, 10.6	16.6	14.7, 18.8	4.3	3.4, 5.4
Mexican-American	6.7	5.9, 7.7	0.83	0.51, 1.35	2.6	1.9, 3.5	7.1	5.2, 9.5	15.6	13.5, 18.1
Other	9.6	8.5, 10.8	11.0	8.4, 14.4	4.0	3.1, 5.1	9.6	7.6, 12.0	14.7	12.3, 17.5
Age-group (years)										
20–39	39.6	38.7, 40.6	39.3	37.3, 41.4	37.3	35.3, 39.2	40.9	39.1, 42.7	40.5	38.0, 43.0
40–59	37.0	36.3, 37.8	36.8	35.4, 38.1	37.6	36.2, 39.0	37.0	35.6, 38.4	36.7	34.8, 38.7
≥ 60	23.3	22.4, 24.3	23.9	22.1, 25.7	25.2	23.6, 26.8	22.1	20.7, 23.6	22.8	20.4, 25.3
Education (years)										
< 12	21.3	20.4, 22.3	19.7	18.0, 21.6	18.1	16.4, 19.8	24.8	22.9, 26.8	20.6	18.6, 22.7
12	27.8	27.1, 28.6	29.1	26.5, 31.8	30.9	29.2, 32.7	28.2	27.0, 29.4	22.9	21.3, 24.6
Some college	27.0	26.3, 27.7	24.3	22.9, 25.8	28.1	26.6, 29.6	25.4	24.2, 26.6	30.9	29.3, 32.4
≥ College	23.8	22.6, 25.0	26.9	24.0, 30.0	22.9	20.6, 25.5	21.7	19.9, 23.6	25.6	23.1, 28.3
Poverty income ratio (PIR; %)										
< 130	18.5	17.4, 19.5	16.8	15.0, 18.8	16.5	14.6, 18.6	21.4	19.2, 23.6	17.3	15.3, 19.3
130–349	36.6	35.6, 37.6	35.4	32.9, 38.0	38.0	35.8, 40.3	37.4	35.6, 39.1	34.8	32.9, 36.7
≥ 350	38.3	37.0, 39.7	39.1	36.1, 42.2	39.6	36.9, 42.4	35.1	32.6, 37.6	41.6	39.1, 41.1
Missing	6.6	6.1, 7.2	8.7	7.2, 10.4	5.8	5.0, 6.8	6.2	5.3, 7.2	6.4	5.5, 7.4
Month of MEC exam										
November–April	46.1	41.9, 50.4	33.0	27.5, 39.1	32.9	26.7, 39.7	55.1	46.4, 63.4	57.3	45.4, 68.4
May–October	53.9	49.6, 58.1	67.0	60.9, 72.5	67.1	60.3, 73.3	44.9	36.6, 53.6	42.7	31.6, 54.6
BMI (kg/m ²)										
< 25	36.7	35.8, 37.6	37.1	34.6, 39.8	35.1	33.1, 37.2	36.5	35.2, 37.8	38.6	37.1, 40.2
25–< 30	33.7	33.0, 34.4	34.3	33.0, 35.7	33.8	32.1, 35.5	32.8	31.7, 33.9	34.7	33.4, 36.0
≥ 30	29.5	28.7, 30.4	28.5	26.5, 30.7	31.1	29.4, 32.9	30.7	29.4, 32.1	26.7	25.1, 28.4
Smoking status										
Never smoked	49.5	48.6, 50.5	48.0	45.6, 50.4	48.0	46.1, 49.9	49.4	47.9, 51.0	52.7	50.2, 55.2
Former smoker	25.2	24.5, 25.9	26.7	25.1, 28.4	25.2	23.8, 26.7	23.7	22.8, 24.7	26.3	24.7, 27.9
Current smoker	25.3	24.5, 26.1	25.3	23.3, 27.4	26.8	25.4, 28.1	26.9	25.5, 28.3	21.0	19.4, 22.6
Reported supplement use	48.6	47.7, 49.6	46.8	45.0, 48.6	48.3	46.5, 50.1	46.1	44.5, 47.6	54.8	52.5, 56.9
Any self-reported chronic disease	31.7	30.9, 32.6	30.7	29.0, 32.5	32.6	30.4, 34.8	33.2	31.7, 34.6	29.4	27.8, 31.0
Alcohol drinking status										
Never	11.4	10.6, 12.3	8.5	7.4, 9.6	9.5	8.1, 11.0	14.2	13.2, 15.3	11.4	8.9, 14.5
Former	18.9	18.1, 19.7	18.1	16.2, 20.3	19.2	17.3, 21.2	20.3	19.0, 21.7	16.9	15.1, 18.9
Current	64.2	62.9, 65.5	67.5	64.9, 70.0	66.8	63.6, 69.9	59.5	57.7, 61.3	66.2	63.1, 69.1
Unknown	5.5	5.1, 5.9	5.9	4.9, 7.0	4.5	4.0, 5.2	6.0	5.2, 6.8	5.5	4.6, 6.5

MEC, mobile examination centre.

Table 2 Covariate*-adjusted geometric means† and 95 % CI of disease and nutritional biomarkers, by US census region, averaged over survey years: US adults, National Health and Nutrition Examination Survey (NHANES) 1988–1994 to 1999–2010

Biomarker	Main effect of US census region averaged over surveys								P value (region × survey)	P value (region)
	Northeast		Midwest		South		West			
	Mean	95 % CI	Mean	95 % CI	Mean	95 % CI	Mean	95 % CI		
Disease biomarkers										
Total cholesterol (mg/dl; n 42 262)	201	199, 203	200	199, 201	201	200, 203	202	200, 203	0.1	0.4
HDL cholesterol (mg/dl; n 42 147)	52.4	51.6, 53.0	51.5	50.7, 52.2	51.2	50.7, 51.6	53.0	52.5, 53.5	0.1	<0.0001
C-reactive protein (mg/dl; n 27 298)	0.17	0.16, 0.18	0.18	0.17, 0.18	0.20	0.19, 0.20	0.17	0.16, 0.18	0.1	<0.0001
Nutritional biomarkers										
Serum folate (nmol/l; n 42 248)	30.8	29.4, 32.2	29.4	28.3, 30.5	28.5	27.5, 29.5	30.8	29.7, 31.9	0.02	0.0007
RBC folate (nmol/l; n 41 832)	992	974, 1009	999	974, 1024	972	956, 988	1045	1020, 1071	0.02	<0.0001
Vitamin B ₁₂ (pmol/l; n 24 066)	334	326, 348	336	330, 342	329	323, 336	350	341, 356	0.7	0.0004
Vitamin C (µmol/l; n 82 75)	55.7	53.4, 58.1	52.3	49.1, 55.5	51.7	50.0, 53.2	57.3	55.2, 59.3	NA	0.0004
Vitamin B ₆ (nmol/l; n 14 770)	51.6	48.7, 54.6	50.8	48.5, 53.2	48.0	45.8, 50.2	52.4	49.4, 55.6	NA	0.007
Vitamin D (nmol/l; n 37 704)	60.7	56.7, 62.8	61.4	60.0, 62.9	56.6	55.0, 58.2	57.2	55.8, 58.6	0.1	<0.0001
Vitamin E (µmol/l; n 31 274)	27.5	27.0, 27.9	26.9	26.6, 27.3	26.8	26.5, 27.1	27.9	27.4, 28.4	0.1	<0.0001
α-Carotene (µmol/l; n 27 563)	0.05	0.05, 0.06	0.05	0.04, 0.05	0.05	0.04, 0.05	0.06	0.06, 0.06	0.2	0.0002
β-Carotene‡ (µmol/l; n 27 203)	0.26	0.25, 0.28	0.25	0.23, 0.26	0.25	0.24, 0.26	0.28	0.26, 0.29	0.2	0.0002
β-Cryptoxanthin (µmol/l; n 27 525)	0.13	0.12, 0.13	0.13	0.12, 0.13	0.13	0.12, 0.13	0.15	0.14, 0.15	0.6	<0.0001
Lutein + zeaxanthin (µmol/l; n 27 562)	0.30	0.29, 0.31	0.27	0.27, 0.28	0.28	0.27, 0.29	0.30	0.29, 0.31	0.2	0.0001
Lycopene (µmol/l; n 27 562)	0.39	0.38, 0.40	0.39	0.37, 0.40	0.36	0.36, 0.37	0.36	0.35, 0.37	0.7	0.0002

RBC, red blood cell; NA, not available (because multiple survey cycles for trend analysis were not available); MEC, mobile examination centre.

*All estimates are adjusted means and 95 % CI from multiple linear regression models with each biomarker as a continuous outcome. Independent variables were: US census region (Northeast, Midwest, South, West), survey years (total and HDL cholesterol, serum and RBC folate, 1988–1994, 1999–2004, 2005, 2010; C-reactive protein, 1999–2002, 2003–2006, 2007–2010; vitamins D and E, 1988–1994, 2001–2004, 2005–2010; vitamin B₁₂, 1988–1994, 1999–2002, 2003–2006; serum vitamin C, 2003–2006; serum vitamin B₆, 2005–2010; and serum carotenoids, 1988–1994, 2001–2006), sex, race/ethnicity (Non-Hispanic Black, Non-Hispanic White, Mexican-American, Others), urban/rural residence (large metropolitan, small/medium metropolitan, fringe metropolitan, non-metropolitan), poverty income ratio, % (<130, 130–349, >349, missing), education, years (<12, 12, some college, ≥ college), age, years (20–39, 40–59, ≥ 60), BMI, kg/m² (<25, 25–<30, ≥30, missing), reported supplement use (yes/no), serum cotinine (continuous), period of MEC exam (May–October, November–April), hours of fasting before phlebotomy (continuous), self-reported chronic disease (yes/no). For total and HDL cholesterol, alcohol drinking status (never, former, current, unknown), and for carotenoids and vitamin E, serum total cholesterol were additional covariates.

†Arithmetic mean for total and HDL cholesterol and vitamin C. For all other biomarkers, the table shows back-transformed geometric means.

‡Significant interaction of region × survey × age (P=0.0002). Age-stratified results are presented in the online supplementary material, Supplemental Table 3.



Table 3 Covariate-adjusted means* and 95% CI of dietary nutrient intakes, by US census region, averaged over survey years: US adults, National Health and Nutrition Examination Survey (NHANES) 1988–1994 to 1999–2010

Dietary variable	Main effect of US census region averaged over surveys								P value (region × survey)	P value (region)
	Northeast		Midwest		South		West			
	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI		
24 h Energy (kJ)	9046	8870, 9222	9125	9021, 9230	9079	8958, 9196	9196	9017, 9376	0.6	0.6
24 h Energy (kcal)	2162	2120, 2204	2181	2156, 2206	2170	2141, 2198	2198	2155, 2241	0.6	0.6
Energy from protein (%)	15.7	15.5, 15.9	15.5	15.3, 15.7	15.5	15.3, 15.6	15.5	15.3, 15.7	0.6	0.4
Energy from carbohydrates (%)	50.1	49.5, 50.8	49.5	48.9, 50.0	49.5	49.1, 49.9	49.4	48.8, 50.0	0.5	0.3
Energy from fat (%)	32.7	32.3, 33.0	33.4	32.9, 33.8	33.6	33.2, 33.9	33.6	33.3, 34.0	0.5	0.0005
Energy from saturated fat (%)	10.9	10.7, 11.1	11.1	10.9, 11.3	11.0	10.8, 11.1	11.1	10.9, 11.2	0.9	0.4
Fibre (g)	16.1	15.6, 16.5	15.9	15.5, 16.3	15.8	15.5, 16.0	17.6	17.0, 18.1	0.1	< 0.0001
Fibre† (g)	16.1	15.8, 16.5	15.9	15.5, 16.2	15.8	15.5, 16.1	17.4	17.0, 17.9	0.1	< 0.0001
Alcohol (g)	10.5	9.4, 11.7	11.6	10.4, 12.8	10.6	9.6, 11.7	10.9	9.5, 12.2	0.3	0.6
Ratio of energy intake to calculated basal energy expenditure (n 41 853)	1.36	1.33, 1.38	1.36	1.35, 1.38	1.36	1.34, 1.39	1.37	1.34, 1.38	0.5	0.8
Vitamin A† (µg RAE)	799	768, 829	715	691, 739	718	693, 744	782	748, 815	0.1	< 0.0001
Retinol† (µg)	484	457, 511	445	429, 461	438	418, 458	465	437, 493	0.4	0.04
Vitamin E† (mg α-tocopherol)	7.70	7.45, 7.96	7.33	7.15, 7.51	7.44	7.25, 7.62	7.97	7.72, 8.22	0.4	< 0.0001
Vitamin C† (mg)	101	96, 106	92	88, 96	88	85, 92	100	96, 104	0.3	< 0.0001
Vitamin B ₆ † (mg)	1.97	1.94, 2.01	1.95	1.92, 1.98	1.90	1.86, 1.93	1.98	1.93, 2.03	0.5	0.0006
Folate† (µg)	385	374, 395	362	353, 371	361	354, 367	380	369, 390	0.02	0.0001
Vitamin B ₁₂ † (µg)	5.75	5.32, 6.18	5.12	4.96, 5.28	5.15	4.99, 5.31	5.37	5.09, 5.65	0.1	0.02
Na† (mg)	3493	3455, 3552	3506	3457, 3554	3528	3481, 3574	3486	3439, 3533	0.01	0.3
K† (mg)	2810	2776, 2845	2776	2735, 2817	2727	2697, 2759	2839	2801, 2877	0.7	< 0.0001
Ca† (mg)	919	896, 942	897	879, 915	856	841, 871	913	893, 934	0.7	< 0.0001
Mg† (mg)	297	292, 302	293	288, 297	290	287, 294	308	303, 313	0.3	< 0.0001
Fe† (mg)	16.0	15.6, 16.4	15.4	15.1, 15.7	15.3	15.1, 15.5	15.8	15.5, 16.1	0.6	0.0007

RAE, retinol activity equivalents; MEC, mobile examination centre.

*All estimates are adjusted means and 95% CI from multivariable linear regression models with each dietary variable as a continuous outcome. Independent variables were: US census region (Northeast, Midwest, South, West), survey year (1988–1994, 1999–2004, 2005–2010), sex, race/ethnicity (Non-Hispanic Black, Non-Hispanic White, Mexican-American, Others), urban/rural residence (large metropolitan, small-medium metropolitan, fringe metropolitan, non-metropolitan), poverty income ratio, % (<130, 130–349, >349, missing), education, years (<12, 12, some college, ≥ college), age, years (20–39, 40–59, ≥ 60), BMI, kg/m² (<25, 25–<30, ≥30, missing), period of MEC exam (May–October, November–April), week day of recalled intake (Monday–Thursday, Friday–Sunday) and self-reported chronic disease (yes/no). n 43 177 with complete covariate information.

†Models included energy intake (kcal/kJ) as a covariate.



Table 4 Covariate-adjusted* means and 95 % CI of food group servings and selected measures of diet quality and composition, by US census region, averaged over survey years: US adults, National Health and Nutrition Examination Survey (NHANES) 1988–1994 to 1999–2010

Dietary variable	Main effect of US census region averaged over surveys								P value (region × survey)	P value (region)
	Northeast		Midwest		South		West			
	Mean	95 % CI	Mean	95 % CI	Mean	95 % CI	Mean	95 % CI		
Total vegetables† (cup-equivalents)	1.67	1.63, 1.71	1.63	1.58, 1.68	1.59	1.55, 1.63	1.70	1.64, 1.75	0.04	0.001
Dark green vegetables† (cup-equivalents)	0.13	0.11, 0.14	0.09	0.08, 0.11	0.12	0.11, 0.13	0.14	0.12, 0.16	0.3	0.0002
Total fruits† (cup-equivalents)	1.13	1.07, 1.18	1.01	0.95, 1.07	0.86	0.82, 0.91	1.05	0.99, 1.11	0.1	<0.0001
Whole fruit† (cup-equivalents)	0.70	0.67, 0.73	0.68	0.64, 0.72	0.52	0.49, 0.55	0.66	0.62, 0.71	0.02	<0.0001
Total dairy† (cup-equivalents)	1.66	1.61, 1.72	1.65	1.59, 1.70	1.50	1.46, 1.54	1.64	1.57, 1.71	0.3	<0.0001
Total meat† (ounce-equivalents)	6.02	5.83, 6.21	6.00	5.83, 6.17	6.21	6.05, 6.36	5.92	5.76, 6.07	0.5	0.04
Total grains† (ounce-equivalents)	6.91	6.71, 7.11	6.59	6.47, 6.71	6.62	6.51, 6.72	6.73	6.56, 6.89	0.1	0.02
Whole grains† (ounce-equivalents)	0.72	0.67, 0.77	0.76	0.71, 0.82	0.68	0.65, 0.71	0.83	0.78, 0.89	0.2	<0.0001
Amount of all foods and beverages (g)	2486	2437, 2534	2568	2525, 2611	2499	2461, 2536	2501	2441, 2561	0.008	0.03
No. of all unique foods and beverages	14.5	14.2, 14.8	14.0	13.8, 14.2	13.9	13.7, 14.1	14.1	13.9, 14.4	0.04	0.005
No. of unique foods from the major food groups	9.1	8.9, 9.3	8.8	8.6, 8.9	8.7	8.5, 8.8	8.9	8.7, 9.1	0.02	0.002
Energy density of foods only (kJ/g)	7.95	7.82, 8.03	8.24	8.12, 8.33	8.24	8.16, 8.33	7.95	7.87, 8.08	0.5	<0.0001
Energy density of foods only (kcal/g)	1.90	1.87, 1.92	1.97	1.94, 1.99	1.97	1.95, 1.99	1.90	1.88, 1.93	0.5	<0.0001
Added sugar† (teaspoon-equivalents)	19.1	18.5, 19.7	20.4	19.5, 21.2	20.5	20.0, 21.0	18.7	17.9, 19.6	0.01	<0.0001
Fibre (g/100 g carbohydrates)	6.34	6.25, 6.43	6.36	6.24, 6.44	6.34	6.24, 6.44	6.95	6.80, 7.11	0.1	<0.0001
Fibre (g/1000 kcal (4184 kJ))	7.85	7.70, 8.01	7.74	7.56, 7.91	7.71	7.58, 7.83	8.41	8.21, 8.62	0.3	<0.0001
% of 24 h energy from beverages	19.4	18.9, 20.0	19.6	18.9, 20.4	19.9	19.4, 20.4	19.4	18.8, 20.1	0.8	0.6
Discretionary solid fat (g)	44.4	43.2, 45.7	45.8	44.7, 46.8	46.2	45.2, 47.2	45.8	44.4, 47.2	0.6	0.2
Energy from solid fat, added sugars and alcohol (kJ)	3059	2987, 3130	3230	3159, 3305	3234	3167, 3305	3121	3025, 3222	0.1	0.0005
Energy from solid fat, added sugars and alcohol (kcal)	731	714, 748	772	755, 790	773	757, 790	746	723, 770	0.1	0.0005

MEC, mobile examination centre.

*All estimates are adjusted means and 95 % CI from multiple linear regression models with each dietary variable as a continuous outcome. Independent variables were: US census region (Northeast, Midwest, South, West), survey year (1988–1994, 1999–2004, 2005–2010), sex, race/ethnicity (Non-Hispanic Black, Non-Hispanic White, Mexican-American, Others), urban/rural residence (large metropolitan, small-medium metropolitan, fringe metropolitan, non-metropolitan), poverty income ratio, % (<130, 130–349, >349, missing), education, years (<12, 12, some college, ≥ college), age, years (20–39, 40–59, ≥ 60), BMI, kg/m² (<25, 25–<30, ≥30, missing), period of MEC exam (May–October, November–April), week day of recalled intake (Monday–Thursday, Friday–Sunday), and self-reported chronic disease (yes/no). *n* 43 177 with complete covariate information.

†Models included energy intake (kcal/kJ) as a covariate.

Table 5 Covariate-adjusted* estimates and 95% CI of eating episode patterns, by US census region, averaged over survey years: US adults, National Health and Nutrition Examination Survey (NHANES) 1988–1994 to 1999–2010

Eating behaviour variable	Main effect of US census region averaged over surveys												P value (region)	
	Northeast			Midwest			South			West				P value (region × survey)
	Mean	95% CI		Mean	95% CI		Mean	95% CI		Mean	95% CI			
No. of main meal† eating episodes	2.7	2.7, 2.8	2.7	2.6, 2.73	2.7	2.7, 2.73	2.7	2.7, 2.74	2.7	2.7, 2.74	2.7	2.7, 2.74	0.1	0.2
No. of non-main meal‡ eating episodes	2.2	2.15, 2.36	2.3	2.27, 2.37	2.1	2.07, 2.17	2.2	2.15, 2.28	2.2	2.15, 2.28	2.2	2.15, 2.28	0.01	<0.0001
% reported breakfast	85	83, 86	80	78, 81	81	80, 82	83	82, 85	83	80, 82	83	82, 85	0.3	<0.0001
Length of the eating period§ (h)	12.5	12.3, 12.7	12.5	12.4, 12.6	12.3	12.2, 12.3	12.2	12.1, 12.3	12.2	12.2, 12.3	12.2	12.1, 12.3	0.1	0.001
Interval between eating episodes (h)	2.66	2.62, 2.70	2.65	2.62, 2.67	2.71	2.69, 2.73	2.64	2.61, 2.67	2.64	2.69, 2.73	2.64	2.61, 2.67	0.04	0.0002

MEC, mobile examination centre.
 *All estimates are adjusted means and 95% CIs from multiple linear regression models with each dietary variable as a continuous outcome. Independent variables were US census region (Northeast, Midwest, South, West), survey year (1988–1994, 1999–2004, 2005–2010), sex, race/ethnicity (Non-Hispanic White, Mexican-American, Others), urban/rural residence (large metropolitan, small-medium metropolitan, fringe metropolitan, non-metropolitan), poverty income ratio, % (<130, 130–349, >349, missing), education, years (<12, 12, some college, ≥ college), age, years (20–39, 40–59, ≥ 60), BMI, kg/m² (<25, 25–<30, ≥30, missing), period of MEC exam (May–October, November–April), week day of recalled intake (Monday–Thursday, Friday–Sunday) and self-reported chronic disease (yes/no). n 43 177 with complete covariate information.
 †Mention of an eating episode labelled as breakfast, lunch, brunch, supper and dinner, or equivalent in Spanish.
 ‡Mention of eating episodes not labelled as a main meal as described above.
 §Length of the eating period/sum of main meal and non-main meal eating episodes.
 || Interval between the time of the first reported episode and the time of the last reported eating episode.

β-carotene concentration were significant in 20–39- and 40–59-year-olds ($P \leq 0.002$); in ≥ 60-year-olds regional differences were not significant (see online supplementary material, Supplemental Table 3). None of the other examined three-way interactions were significant.

Dietary outcomes

Did the association of region with dietary intakes or eating behaviours change over time?

The weighted, unadjusted mean and 95% CI of dietary outcomes, by region, by survey year, are shown in the online supplementary material, Supplemental Tables 4–6. In multiple covariate-adjusted models, the region × survey interaction term was not significant ($P > 0.001$) for all examined dietary nutrient, food group, diet quality and eating behaviour outcomes (Tables 3–5).

Did reported dietary intakes and eating behaviours, averaged over surveys, differ among US census regions?

Energy and nutrient intakes. Mean intakes of energy, percentage of energy from protein, carbohydrates and saturated fat, grams of alcohol, vitamin B₁₂ and Na did not differ among regions ($P > 0.001$; Table 3). Intakes of dietary fibre and all other examined micronutrients (vitamins A, B₆, C, E and folate, and the minerals K, Ca, Mg and Fe) were significantly different ($P < 0.001$) among regions. For these nutrients, highest mean intakes were in the West or the Northeast region, while lowest mean intakes were associated with the South region. The ratio of reported energy intake to estimated basal energy expenditure (a possible measure of energy under-reporting) did not differ among US census regions.

Food group serving equivalents and estimates of diet composition and quality. Mean servings of vegetables, fruits, whole fruits, total dairy and whole grains, energy density (kcal/g, kJ/g) of foods, added sugar, g fibre/100 g carbohydrates, g fibre/1000 kcal (4184 kJ), and combined energy from discretionary solid fat, added sugar and alcohol were significantly different among regions ($P \leq 0.001$; Table 4). Mean amount or number of all reported foods and beverages, percentage of energy from beverages, grams of discretionary solid fat, and ounce-equivalents of meat and total grains did not differ among regions ($P > 0.001$). Relative to the West and the Northeast regions, in the South and the Midwest regions, the energy density of reported foods, added sugar, and combined energy from solid fats, added sugar and alcohol (nutrient-poor foods) were higher, while g fibre/100 g carbohydrates, g fibre/1000 kcal (4184 kJ), vegetable, fruit and whole grain intakes were lower.

Eating pattern outcomes. All examined eating behaviours, except the number of main meal eating episodes, were significantly different among regions ($P < 0.005$; Table 5). The South region reported lower mean number of non-main meal eating episodes and longer intervals



between eating episodes. Smaller percentages of those in the South and the Midwest reported breakfast. The length of the eating period (difference between the first and the last reported eating episode of the day) was longer in the Northeast and the Midwest relative to the other two regions.

Effect modification testing for dietary outcomes

The three-way interactions of survey \times region \times sex, survey \times region \times race/ethnicity, survey \times region \times age, and survey \times region \times PIR were not significant at the Bonferroni-corrected P values of <0.0006 , <0.0007 and <0.002 for the dietary outcomes presented in Tables 3, 4 and 5, respectively.

Discussion

In our study, region was a strong correlate of nearly all examined nutritional biomarkers, micronutrients and food groups, with higher mean concentrations and self-reported intakes in the West and Northeast, followed by the Midwest and the South. These results point to a distinct profile of generally higher-risk dietary patterns in the South and Midwest regions. The study findings point to similarities in directions of regional gradients in nutritional biomarkers and dietary intakes with gradients reported for health outcomes. For example, age-adjusted mortality rates from all causes, circulatory diseases and all cancers combined show differentials by census region and division for both sexes^(1–6). The mortality rates are highest in the South, followed by the Midwest, Northeast and the West. The prevalence of obesity (BMI ≥ 30 kg/m²) among American adults, based on self-reported body weight and height in the Behavioral Risk Factor Surveillance System (BRFSS), shows regional variation similar to that of mortality mentioned above⁽⁹⁾. Diabetes prevalence also has been reported to fall along similar geographic–regional lines⁽¹¹⁾.

The absence of a significant interaction of survey with region for the examined outcomes suggests that regional differences in dietary intakes and nutritional biomarker concentrations were maintained over time. To our knowledge, the only published report on secular trends in regional differences in a dietary outcome used data from the BRFSS 1994–2007 and found that fruit and vegetable consumption was unchanged over the examined period⁽¹⁴⁾. We also found little evidence of modification of the region and survey interaction by sex, age, income or race/ethnicity, as the three-way interaction terms to test the effect did not reach the Bonferroni-adjusted level of significance.

To our knowledge, the present study is the first to report concurrent examination of secular trends in nutritional biomarkers and dietary intakes across US census regions. Because serum concentrations of most of the examined nutrients (except vitamin D) represent the end result of nutrient consumption and metabolism⁽²⁴⁾, these findings point to possible differences in exposure to examined nutrients, which was confirmed by dietary intakes of food

group sources of micronutrients. Although overall energy intakes did not differ among regions, energy density of reported foods was higher in the at-risk South and Midwest regions. These findings are in accord with lower servings of fruits, vegetables and whole grains, and poorer carbohydrate quality reflected in g fibre/100 g carbohydrates and higher added sugar intakes. Due to differences in dietary assessment and analytic methodology, our results are not directly comparable to the few published reports. Overall, our results on dietary intakes are consistent with prior reports of gradients across US census regions for fruits and vegetables^(17,42), nutrient intakes⁽¹⁷⁾ and intake of sugar-sweetened beverages⁽²²⁾. The US Department of Agriculture's 1977–78 Nationwide Food Consumption Survey indicated that the South census region of the country had the smallest percentage of individuals reporting carotenoid-rich deep-yellow vegetables and all citrus or non-citrus fruits and juices, but the highest percentage who reported sugar⁽¹⁷⁾. Moreover, micronutrient intakes relative to the recommended standards were lowest in the South⁽¹⁷⁾. Both food and nutrient intakes were most favourable in the West census region⁽¹⁷⁾. We note that these are unadjusted estimates; to our knowledge, regional differences in dietary attributes to account for socio-demographic variables that are differentially distributed among regions has not been published with these US Department of Agriculture data. Analysis of the 2011 BRFSS data found that odds of mention of at least one serving of fruit were lowest in the South, while odds for mention of at least one vegetable were lowest in the Midwest⁽⁴²⁾. The odds of consuming at least one serving of a regular sweetened beverage were also higher in the South region in the 2010 National Health Interview Survey⁽²²⁾.

These dietary profiles suggest qualitative differences in the types of foods reported across US regions. The surprising persistence of regional differentials over more than two decades suggests little change in drivers of these differentials. Possible contributors to regional dietary differentials include differences in food access and learned food preferences unique to regional foodways. Although we cannot exclude the possibility of some residual confounding, we note that our analyses adjusted for many of the factors that may relate to food access, such as race/ethnicity, income, education and urbanization of area of residence. The US food system includes marketing of national brands across the country, thus assuring marketing of similar products (e.g. Na content of foods) nationwide⁽⁴³⁾. However, regional demand for some foods may contribute to differences in the market share of various products across regions⁽⁴³⁾. Regional demand likely reflects learned preferences for types of foods, their methods of preparation, frequency of their consumption and the combinations in which they are consumed, which in turn can contribute to regional differences in food intake. Therefore, our results are in accord with prior observations about the powerful role of regional food



saliency and foodways in determining food consumption^(44–46), which present a formidable challenge to adoption of recommended food-related behaviours.

Although our study included a nationally representative sample of US adults and used objective biomarkers of dietary exposure along with self-reported dietary intakes, we acknowledge that the census region is a broad categorization of region and assumes a degree of homogeneity within a region. The only available national estimates of both biochemical and dietary outcomes for the US population come from the NHANES, and the NHANES sample design does not permit evaluation of state- and county-specific estimates. To our knowledge, there are no other national estimates of time trends in both biochemical and dietary outcomes for the regional exposure examined in the present study. The breadth of outcomes examined herein was intentional, to provide a comprehensive profile of regional dietary exposures. We used a Bonferroni correction to reduce the Type 1 error (false positives) due to testing of multiple hypotheses. The potential cost of this conservative correction is the likely increase in the Type II error (false negatives)⁽⁴⁷⁾ and may have resulted in negative findings, especially for the three-way interactions examined in our study. This approach may partially explain the lack of concordance between our study finding of no significant effect modification by race/ethnicity, income and education and those reported in prior studies^(20,21,42,44).

All methods of dietary assessment are prone to systematic and random measurement errors⁽²³⁾. While the nutritional biomarkers due to their objectivity do not have the same measurement errors found in self-reported dietary intakes, they are not error free⁽²⁴⁾. Despite the known measurement errors⁽⁴⁸⁾, the study findings of concordance between dietary intakes and serum biomarker concentrations provide a broad validation of population estimates of self-reported food group and micronutrient intakes in the NHANES.

Conclusion

In conclusion, regional gradients in dietary exposure expressed objectively as biomarkers or as self-reported nutrient and food group intakes followed trajectories reported for health outcomes and were remarkably persistent over time. As suggested by Ricketts⁽⁶⁾, given that state, county, city and racial/ethnic political constituencies are powerful drivers of resource allocation, to address regional disparities in health outcomes, national policies are essential to target regional health behaviours and resources. Persistent differentials in dietary exposures suggest several areas of intervention that need to target added sugars, whole grains, total and whole fruits, and vegetables. Development of such interventions requires further study of regional foodways, regionally acceptable food options, and an understanding of the socio-ecological barriers to dietary change.

Acknowledgements

Acknowledgements: The authors thank Lisa Licitra Kahle for expert SAS and SUDDAN programming support, and Nataliya Kravet, NCHS, for facilitating use of the RDC (Hyattsville, MD, USA) on-site facilities for data analysis. **Disclosure:** Preliminary findings were presented at the Experimental Biology Meeting, San Diego, CA, USA, 2–6 April 2016. **Financial support:** This work was supported in part by the National Institute on Aging, National Institutes of Health (A.K.K., grant number AG046540); and by the intramural research programme of the Department of Health and Human Services, National Cancer Institute, National Institutes of Health (B.I.G.). The funders had no role in the design, analysis or writing of this article. **Conflict of interest:** None. **Authorship:** A.K.K. conceptualized the study question, designed the research, analysed the data, wrote the paper, and had primary responsibility for the final content. B.I.G. provided guidance on the study design and analytic strategy, and reviewed the paper for important intellectual content. **Ethics of human subject participation:** The City University of New York, Human Research Protection Program, did not consider the study human subjects research.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1368980017003743>

References

1. Pickle LW, Mungiole M & Gillum RF (1997) Geographic variation in stroke mortality in blacks and whites in the United States. *Stroke* **28**, 1639–1647.
2. Pickle LW & Gillum RF (1999) Geographic variation in cardiovascular disease mortality in US blacks and whites. *J Natl Med Assoc* **91**, 545–556.
3. US Cancer Statistics Working Group (2015) *United States Cancer Statistics: 1999–2012 Incidence and Mortality Web-based Report*. Atlanta, GA: Department of Health and Human Services, Centers for Disease Control and Prevention, and National Cancer Institute.
4. National Vital Statistics System (2015) Age-adjusted death rates for stroke, by US census regions – United States, 1970–2013. *MMWR Morb Mortal Wkly Rep* **64**, 371; available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6413a9.htm>
5. National Vital Statistics System (2015) Age-adjusted death rates from cancer, by US census region, and year – United States, 1970–2013. *MMWR Morb Mortal Wkly Rep* **64**, 774; available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6428a8.htm>
6. Ricketts TC (2007) Geography and health disparity. Woodrow Wilson International Center for Scholars Conference Paper. <http://www.wilsoncenter.org/sites/default/files/Ricketts1.pdf> (accessed July 2015).
7. Xu JQ, Murphy SL, Kochanek KD *et al.* (2016) Deaths: final data for 2013. *Natl Vital Stat Rep* **64**, issue 2, 1–119.
8. Secretary's Advisory Committee on Health Promotion and Disease Prevention Objectives for 2020 (2010) Healthy



- People 2020: an opportunity to address the societal determinants of health in the United States. <http://www.healthypeople.gov/2010/hp2020/advisory/SocietalDeterminantsHealth.html> (accessed February 2015).
9. Centers for Disease Control and Prevention (2010) State-specific obesity prevalence among adults – United States, 2009. *MMWR Morb Mortal Wkly Rep* **59**, 951–955; available at <http://www.cdc.gov/mmwr/pdf/wk/mm59e0803.pdf>
 10. Centers for Disease Control and Prevention (2008) Prevalence of self-reported physically active adults – United States, 2007. *MMWR Morb Mortal Wkly Rep* **57**, 1297–1300; available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5748a1.htm#tab>
 11. Centers for Disease Control and Prevention (2013) CDC health disparities and inequalities report, United States, 2013. *MMWR Morb Mortal Wkly Rep* **62**, Suppl. 3, 1–184; available at <http://www.cdc.gov/mmwr/pdf/other/su6203.pdf>
 12. Obisesan TO, Vargas CM & Gillum RF (2000) Geographic variation in stroke risk in the United States. Region, urbanization, and hypertension in the Third National Health and Nutrition Examination Survey. *Stroke* **31**, 19–25.
 13. Clark CR, Coull B, Berkman LF *et al.* (2011) Geographic variation in cardiovascular inflammation among healthy women in the Women's Health Study. *PLoS One* **6**, e27468.
 14. Troost JP, Rafferty AP, Luo Z *et al.* (2012) Temporal and regional trends in the prevalence of healthy lifestyle characteristics: United States, 1994–2007. *Am J Public Health* **102**, 1392–1398.
 15. World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases. Report of a Joint WHO/FAO Expert Consultation. WHO Technical Report Series* no. 916. Geneva: WHO.
 16. US Department of Agriculture (2015) Scientific Report of the 2015 Dietary Guidelines Advisory Committee. <https://www.health.gov/dietaryguidelines/2015-scientific-report/PDFs/Scientific-Report-of-the-2015-Dietary-Guidelines-Advisory-Committee.pdf> (accessed July 2016).
 17. US Department of Agriculture (1985) *Food and Nutrient Intakes: Individuals in Four Regions, 1977–78. Nationwide Food Consumption Survey 1977–78. Report I-3*. Hyattsville, MD: Nutrition Monitoring Division, Human Nutrition Information Service, US Department of Agriculture.
 18. Hajjar I & Kotchen T (2003) Regional variations of blood pressure in the United States are associated with regional variations in dietary intakes: the NHANES-III data. *J Nutr* **133**, 211–214.
 19. Dubowitz T, Heron M, Bird CE *et al.* (2008) Neighborhood socioeconomic status and fruit and vegetable intake among whites, blacks, and Mexican Americans in the United States. *Am J Clin Nutr* **87**, 1883–1891.
 20. Newby PK, Noel SE, Grant R *et al.* (2011) Race and region are associated with nutrient intakes among black and white men in the United States. *J Nutr* **41**, 296–303.
 21. Newby PK, Noel SE, Grant R *et al.* (2012) Race and region have independent and synergistic effects on dietary intakes in black and white women. *Nutr J* **11**, 25.
 22. Park S, McGuire LC & Galuska DA (2015) Regional differences in sugar-sweetened beverage intake among US adults. *J Acad Nutr Diet* **115**, 1996–2002.
 23. Food and Nutrition Board, Institute of Medicine (2000) *Dietary Reference Intakes: Applications in Dietary Assessments*. Washington, DC: National Academy Press.
 24. Potischman N (2003) Biologic and methodologic issues for nutritional biomarkers. *J Nutr* **133**, Suppl. 3, 875S–880S.
 25. Centers for Disease Control and Prevention, National Center for Health Statistics (2013) National Health and Nutrition Examination Surveys. Questionnaire, Datasets, and Related Documentation. <http://www.n.cdc.gov/nchs/nhanes/Default.aspx> (accessed July 2015).
 26. Centers for Disease Control and Prevention, Research Data Center (2012) National Health and Nutrition Examination Surveys (NHANES) Geocodes. http://www.cdc.gov/rdc/geocodes/geowt_nhanes.htm (accessed July 2015).
 27. Centers for Disease Control and Prevention, National Center for Health Statistics (2012) National Health and Nutrition Examination Surveys. Survey Response Rates and Population Totals. <http://www.n.cdc.gov/Nchs/Nhanes/ResponseRates.aspx> (accessed July 2015).
 28. US Census Bureau (n.d.) Geographic Terms and Concepts – Census Divisions and Census Regions. http://www.census.gov/geo/reference/gtc/gtc_census_divreg.html (accessed February 2015).
 29. Centers for Disease Control and Prevention, National Center for Health Statistics (n.d.) NCHS Urban Rural Scheme for Counties. http://www.cdc.gov/nchs/data_access/urban_rural.htm (accessed July 2015).
 30. Livingstone MB & Black AE (2003) Markers of the validity of reported energy intake. *J Nutr* **133**, Suppl. 3, 895S–920S.
 31. Centers for Disease Control and Prevention, National Center for Health Statistics (2000) Third National Health and Nutrition Examination Survey. Healthy Eating Index. <http://www.n.cdc.gov/nchs/Data/Nhanes3/6A/hei-acc.pdf> (accessed July 2015).
 32. US Department of Agriculture, Agricultural Research Service (n.d.) Food Patterns Equivalents Database. <http://www.ars.usda.gov/northeast-area/beltsville-md/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/fped-databases/> (accessed July 2015).
 33. St-Onge MP, Ard J, Baskin ML *et al.* (2017) Meal timing and frequency: implications for cardiovascular disease prevention: a scientific statement from the American Heart Association. *Circulation* **135**, e96–e121.
 34. Kant AK & Graubard BI (2013) Family income and education were related with 30-year time trends in dietary and meal behaviors of American children and adolescents. *J Nutr* **143**, 690–700. Erratum in: *J Nutr* **143**, 1348.
 35. Kant AK & Graubard BI (2015) 40-year trends in meal and snack eating behaviors of American adults. *J Acad Nutr Diet* **115**, 50–63.
 36. Johnson CL, Paulose-Ram R, Ogden CL *et al.* (2013) National health and nutrition examination survey: analytic guidelines, 1999–2010. *Vital Health Stat* **2**, issue 161, 1–24.
 37. Korn EL & Graubard BI (1999) *Analysis of Health Surveys*. New York: John Wiley and Sons.
 38. Graubard BI & Korn EL (1999) Predictive margins with survey data. *Biometrics* **55**, 652–659.
 39. Centers for Disease Control and Prevention, National Center for Environmental Health, Division of Laboratory Sciences (2012) Second National Report on Biochemical Indicators of Diet and Nutrition in the US Population. http://www.cdc.gov/nutritionreport/pdf/Nutrition_Book_complete508_final.pdf (accessed February 2015).
 40. Kant AK & Graubard BI (2006) Secular trends in patterns of self-reported food consumption of adult Americans: NHANES 1971–1975 to NHANES 1999–2002. *Am J Clin Nutr* **84**, 1215–1223.
 41. Kant AK & Graubard BI (2007) Secular trends in the association of socio-economic position with self-reported dietary attributes and biomarkers in the US population: National Health and Nutrition Examination Survey (NHANES) 1971–1975 to NHANES 1999–2002. *Public Health Nutr* **10**, 158–167.
 42. Tichenor N & Conrad Z (2016) Inter- and independent effects of region and race/ethnicity on variety of fruit and vegetable consumption in the USA: 2011 Behavioral Risk Factor Surveillance System (BRFSS). *Public Health Nutr* **19**, 104–113.
 43. Lee AK, Schieb IJ, Yuan K *et al.* (2015) Sodium content in packaged foods by census division in the United States,



2009. *Prev Chronic Dis* **12**, E43 Erratum in: *Prev Chronic Dis* **2**, E139.
44. Latshaw BA (2009) Food for thought: race, region, identity, and foodways in the American South. *South Cult* **15**, 106–128.
45. Yang Y, Buys DR, Judd SE *et al.* (2013) Favorite foods of older adults living in the Black Belt Region of the United States. Influences of ethnicity, gender, and education. *Appetite* **63**, 18–23.
46. Tucker KL, Maras J, Champagne C *et al.* (2005) A regional food-frequency questionnaire for the US Mississippi Delta. *Public Health Nutr* **8**, 87–96.
47. Perneger TV (1998) What's wrong with Bonferroni adjustments. *BMJ* **316**, 1236–1238.
48. Subar AF, Freedman LS, Tooze JA *et al.* (2015) Addressing current criticism regarding the value of self-report dietary data. *J Nutr* **145**, 2639–2645.

Appendix

Summary of available nutritional and disease biomarkers and laboratory assay methodology: US adults, National Health and Nutrition Examination Survey (NHANES) 1988–1994 to 1999–2010

Biomarker	Surveys available and grouped for trend analysis	N	Laboratory assay method
Serum total cholesterol	1988–1994, 1999–2004, 2005–2010	42 821	Enzymatic assay
Serum HDL cholesterol	1988–1994, 1999–2004, 2005–2010	42 296	Heparin manganese precipitation (1988–1994 and 1999–2002); direct immunoassay (2003–2010)
C-reactive protein	1999–2002, 2003–2006, 2007–2010	27 571	Latex-enhanced nephelometry
Serum and RBC folate	1988–1994, 1999–2004, 2005–2010	42 994 (serum) 42 568 (RBC)	Quantaphase II radioassay kit (1988–2006); microbiological method (2007–2010) Radioassay results harmonized to microbiological method (NCHS)
Serum carotenoids (α-carotene, β-carotene, β-cryptoxanthin, lutein/zeaxanthin, lycopene)	1988–1994, 2001–2006 (trend not examined; difference was examined)	~26 516	Isocratic HPLC
Vitamin D (25-hydroxycholecalciferol)	1988–1994, 2001–2004, 2005–2010	38 297	Diasorin RIA (1988–2006); LC (2007–2010) RIA harmonized to LC using regression equations
Vitamin E (α-tocopherol)	1988–1994, 2001–2004, 2005–2010	31 803	Isocratic HPLC
Vitamin C	2003–2006 (trend not examined)	8309	Isocratic HPLC
Vitamin B ₆ (plasma pyridoxal-5'-phosphate)	2005–2010 (trend not examined)	14 813	Reversed-phase HPLC with fluorometric detection
Vitamin B ₁₂	1988–1994, 1999–2002, 2003–2006	24 447	Quantaphase II radioassay kit

RBC, red blood cell; NCHS, National Center for Health Statistics.