

City University of New York (CUNY)

## CUNY Academic Works

---

Publications and Research

City College of New York

---

2015

### Predicting antidepressant treatment without controlling for depression is ill-advised

Renzo Bianchi  
*Université de Neuchâtel*

Irvin Sam Schonfeld  
*CUNY Graduate Center*

Eric Laurent  
*Université de Franche-Comté*

[How does access to this work benefit you? Let us know!](#)

More information about this work at: [https://academicworks.cuny.edu/cc\\_pubs/324](https://academicworks.cuny.edu/cc_pubs/324)

Discover additional works at: <https://academicworks.cuny.edu>

---

This work is made publicly available by the City University of New York (CUNY).  
Contact: [AcademicWorks@cuny.edu](mailto:AcademicWorks@cuny.edu)



## Letter to the Editor

# Predicting antidepressant treatment without controlling for depression is ill-advised




---

**Keywords:**

Antidepressant treatment  
Burnout  
Conceptual overlap  
Confounding variable  
Depression  
Methodology  
Third variable issue

---

Madsen et al. (2015) examined the prospective association between work-related burnout and antidepressant treatment in human service workers. The authors defined work-related burnout based on exhaustion symptoms assessed in the occupational context. They concluded that *work-related burnout is an important risk factor for antidepressant treatment*. While we read the authors' article with interest, we think that the internal validity of their study is threatened by the absence of assessment and statistical control of depression. This state of affairs undermines the conclusions drawn by the authors.

Given that the aim of the reported study was to examine the link between burnout and antidepressant treatment, the non-consideration of depression (both in terms of depressive symptoms and depressive disorders) is particularly problematic. Because of this omission, the authors cannot establish that burnout *independently* affects antidepressant intake. Put differently, whether burnout accounts for variance in antidepressant intake that is not explained by depression cannot be determined, leaving the study results largely uninterpretable.

Many studies have shown that distinguishing burnout from depression is challenging (Ahola et al., 2014; Bianchi and Laurent, 2015; Bianchi et al., 2014, 2015a, 2015b; Hintsa et al., in press; Rydmark et al., 2006; Wang et al., 2012). For example, in a recent, nosology-focused study, no diagnostically significant differences were found between burned out workers and clinically depressed patients in terms of reported depressive symptoms (Bianchi et al., 2013). At an etiological level, (chronic) work stress—from which burnout is supposed to result (Weber and Jaekel-Reinhard, 2000)—is also known to be an important depressogenic factor

(McTernan et al., 2013; Netterstrøm et al., 2008; Niedhammer et al., 2015; Tennant, 2001; Wang, 2005) and has been associated with antidepressant medication (Virtanen et al., 2007) and suicide (Amagasa et al., 2005). In this context of research suggesting between-syndrome overlap, it would have been especially interesting to observe whether burnout still predicted antidepressant treatment once depression was controlled for (see also Shirom, 2005).

Assessing depression would have also helped clarify the clinical meaning of the cutoff scores used by the authors to categorize levels of burnout. Indeed, although the authors lament a lack of clinical reference points in past burnout research, they themselves use categorization criteria for burnout whose clinical underpinning is elusive. For instance, when identifying workers with “high burnout” using a cutoff score of 50/100 on the Copenhagen Burnout Inventory—corresponding to burnout symptoms experienced “sometimes or somewhat” (see Kristensen et al., 2005, p. 200), the authors rely on a convenience cutpoint rather than a clinically-informed one. These categorization strategies add to the uncertainty surrounding the interpretation of the study results.

All in all, the link observed by Madsen et al. (2015) between burnout and antidepressant intake may well conceal an expected relationship between depression and antidepressant intake. The impossibility of ruling out this reasonable alternative hypothesis renders the study inconclusive. We recommend that future research on (psychotherapeutic or pharmacologic) treatments for burnout incorporate measures of depression in order to facilitate research advance on burnout's characterization and burnout-depression overlap.

## Contributors

Each of the three authors has contributed to the preparation of the article (e.g., writing, reviewing, editing) until a consensus was reached on the final version of the manuscript. All authors have approved the final version of the manuscript.

## Role of the funding source

No funding source involved.

## Conflicts of interest

None.

## Acknowledgement

None.

DOI of original article: <http://dx.doi.org/10.1016/j.jpsychires.2015.07.006>.

## References

- Ahola, K., Hakonen, J., Perhoniemi, R., Mutanen, P., 2014. Relationship between burnout and depressive symptoms: a study using the person-centred approach. *Burn. Res.* 1, 29–37.
- Amagasa, T., Nakayama, T., Takahashi, Y., 2005. Karojisatsu in Japan: characteristics of 22 cases of work-related suicide. *J. Occup. Health* 47, 157–164.
- Bianchi, R., Boffy, C., Hingray, C., Truchot, D., Laurent, E., 2013. Comparative symptomatology of burnout and depression. *J. Health Psychol.* 18, 782–787.
- Bianchi, R., Laurent, E., 2015. Emotional information processing in depression and burnout: an eye-tracking study. *Eur. Arch. Psychiatr. Clin. Neurosci.* 265, 27–34.
- Bianchi, R., Schonfeld, I.S., Laurent, E., 2014. Is burnout a depressive disorder? A re-examination with special focus on atypical depression. *Int. J. Stress Manag.* 21, 307–324.
- Bianchi, R., Schonfeld, I.S., Laurent, E., 2015a. Burnout-depression overlap: a review. *Clin. Psychol. Rev.* 36, 28–41.
- Bianchi, R., Schonfeld, I.S., Laurent, E., 2015b. Is burnout separable from depression in cluster analysis? A longitudinal study. *Soc. Psychiatry Psychiatr. Epidemiol.* 50 (6), 1005–1011.
- Hintsala, T., Elovainio, M., Jokela, M., Ahola, K., Virtanen, M., Pirkola, S., 2015. Is there an independent association between burnout and increased allostatic load? Testing the contribution of psychological distress and depression. *J. Health Psychol.* (in press).
- Kristensen, T.S., Borritz, M., Villadsen, E., Christensen, K.B., 2005. The Copenhagen Burnout Inventory: a new tool for the assessment of burnout. *Work Stress* 19, 192–207.
- Madsen, I.E.H., Lange, T., Borritz, M., Rugulies, R., 2015. Burnout as a risk factor for antidepressant treatment—a repeated measures time-to-event analysis of 2936 Danish human service workers. *J. Psychiatr. Res.* 65, 47–52.
- McTernan, W.P., Dollard, M.F., LaMontagne, A.D., 2013. Depression in the workplace: an economic cost analysis of depression-related productivity loss attributable to job strain and bullying. *Work Stress* 27, 321–338.
- Netterstrøm, B., Conrad, N., Bech, P., Fink, P., Olsen, O., Rugulies, R., et al., 2008. The relation between work-related psychosocial factors and the development of depression. *Epidemiol. Rev.* 30, 118–132.
- Niedhammer, I., Malard, L., Chastang, J.F., 2015. Occupational factors and subsequent major depressive and generalized anxiety disorders in the prospective French national SIP study. *BMC Public Health* 15, 200.
- Rydmark, I., Wahlberg, K., Ghatan, P.H., Modell, S., Nygren, Å., Ingvar, M., et al., 2006. Neuroendocrine, cognitive and structural imaging characteristics of women on longterm sickleave with job stress-induced depression. *Biol. Psychiatry* 60, 867–873.
- Shirom, A., 2005. Reflections on the study of burnout. *Work Stress* 19, 263–270.
- Tennant, C., 2001. Work-related stress and depressive disorders. *J. Psychosom. Res.* 51, 697–704.
- Virtanen, M., Honkonen, T., Kivimäki, M., Ahola, K., Vahtera, J., Aromaa, A., et al., 2007. Work stress, mental health and antidepressant medication findings from the health 2000 study. *J. Affect Disord.* 98, 189–197.
- Wang, J., 2005. Work stress as a risk factor for major depressive episode(s). *Psychol. Med.* 35, 865–871.
- Wang, J., Smailes, E., Sareen, J., Schmitz, N., Fick, G., Patten, S., 2012. Three job-related stress models and depression: a population-based study. *Soc. Psychiatry Psychiatr. Epidemiol.* 47, 185–193.
- Weber, A., Jaekel-Reinhard, A., 2000. Burnout syndrome: a disease of modern societies? *Occup. Med.* 50, 512–517.

Renzo Bianchi\*

University of Franche-Comté, Department of Psychology, 30-32 rue Mégevand, 25030 Besançon Cedex, France

Irvin Sam Schonfeld

The City College of the City University of New York, Department of Psychology, Convent Avenue at 138th Street, New York, NY, 10031, USA

E-mail address: ischonfeld@ccny.cuny.edu.

Eric Laurent

University of Franche-Comté, Department of Psychology, 30-32 rue Mégevand, 25030, Besançon Cedex, France

E-mail address: eric.laurent@univ-fcomte.fr.

\* Corresponding author. Université de Franche-Comté, Laboratoire de Psychologie (EA 3188), 30-32 rue Mégevand, 25030 Besançon Cedex, France.

E-mail address: renzo.bianchi@univ-fcomte.fr (R. Bianchi).

15 May 2015