

2016

Burnout and depression in psychiatric residents

Irvin Sam Schonfeld
CUNY Graduate Center


Eric Laurent
Université de Franche-Comté

Pierre Vandel
Université de Franche-Comté

Renzo Bianchi
Université de Neuchâtel

How does access to this work benefit you? Let us know!

Follow this and additional works at: http://academicworks.cuny.edu/cc_pubs

 Part of the [Clinical Psychology Commons](#), [Health Psychology Commons](#), [Medical Education Commons](#), [Psychiatry and Psychology Commons](#), and the [Public Health Commons](#)

Recommended Citation

Schonfeld, I.S., Laurent, E., Vandel, P., & Bianchi, R. (2016). Burnout and depression in psychiatric residents. *Canadian Journal of Psychiatry*, 61, 737-738. doi:10.1177/0706743716664333

This Article is brought to you for free and open access by the City College of New York at CUNY Academic Works. It has been accepted for inclusion in Publications and Research by an authorized administrator of CUNY Academic Works. For more information, please contact AcademicWorks@cuny.edu.

Burnout and Depression in Psychiatric Residents

The Canadian Journal of Psychiatry /
La Revue Canadienne de Psychiatrie
2016, Vol. 61(11) 737-738
© The Author(s) 2016
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/0706743716664333
TheCJP.ca | LaRCP.ca



Dear Editor:

A recent article¹ in this journal described the results of a study of burnout in psychiatric residents. Using a 1-item scale to assess burnout, the investigators found that 21% of the residents were symptomatic. Aside from problems inherent in the absence of binding or consensual criteria to diagnose burnout, the article ignored research that connects burnout and depression.

Mounting evidence, including evidence from research on health professionals,^{2,3} has linked burnout and depression and suggested that burnout is a depressive syndrome.⁴⁻⁶ Studies conducted in France⁵ and the United States⁶ found that teachers with high levels of burnout symptoms, compared to colleagues with few symptoms, were much more likely to experience the full array of depressive symptoms, including the most severe (e.g., suicidal ideation). In fact, in the French and US samples, burnout was assessed with the most commonly employed burnout instruments.^{7,8} In both studies, when measurement error was controlled, burnout and depressive symptoms⁹ correlated very highly ($r \approx .80$). Moreover, burnout and depression have both been etiologically associated with unresolvable stress. Burnout is assumed to be a product of unresolvable job stress. Unresolvable job stress has been causally related to depression.¹⁰ Burnout and depression also share similar dispositional risk factors (e.g., neuroticism) and overlap in terms of allostatic load, an index of the cumulative biological cost of experienced psychosocial adversity.^{3,11,12}

We therefore submit that in evaluating the distress experienced by overburdened psychiatric residents, investigators assess a problem with which psychiatry is already well familiar, namely, depression. Given the overlap of burnout with depression and the diagnostic blur surrounding burnout, we recommend that depression, rather than burnout, be assessed in occupational health research. In contrast to burnout, depression is nosologically well characterized and diagnosable using clinically validated instruments. To etiologically connect depression with work, the investigator can ask participants whether they mainly attribute their depressive symptoms to work-related problems.

Irvin Sam Schonfeld, PhD, MPH
*Department of Psychology, The City College and the
Graduate Center of the City University of New York,
New York, NY, USA*
ischonfeld@ccny.cuny.edu

Eric Laurent, PhD
*Department of Psychology and Laboratory of Psychology
(EA 3188), University of Franche-Comté, Besançon, France*

Pierre Vandell, MD
*Department of Psychiatry, University Hospital of Besançon,
and Laboratory of Neuroscience (EA 481) University of
Franche-Comté, Besançon, France*

Renzo Bianchi, PhD
*Institute of Work and Organizational Psychology, University
of Neuchâtel, Neuchâtel, Switzerland*

References

1. Kealy D, Halli P, Ogrodniczuk JS, Hadjipavlou G. Burnout among Canadian psychiatry residents: a national survey. *Can J Psychiatry*. 2016;61(11):732-736.
2. Wurm W, Vogel K, Hofmann P, et al. Depression-burnout overlap in physicians. *PLoS One*. 2016;11:e0149913.
3. Ahola K, Hakanen J, Perhoniemi R, Mutanen P. Relationship between burnout and depressive symptoms: a study using the person-centred approach. *Burnout Res*. 2014;1:29-37.
4. Bianchi R, Schonfeld I, Laurent E. Is burnout a depressive disorder? A reexamination with special focus on atypical depression. *Int J Stress Manag*. 2014;21:307-324.
5. Schonfeld I, Bianchi R. Burnout and depression: two entities or one? *J Clin Psychol*. 2016;72:22-37.
6. Bianchi R, Schonfeld I, Laurent E. Burnout does not help predict depression among French school teachers. *Scand J Work Environ Health*. 2015;41:565-568.
7. Maslach C, Jackson SE, Leiter MP. *Maslach Burnout Inventory manual*. 3rd ed. Palo Alto (CA): Consulting Psychologists Press; 1996.

8. Shirom A, Melamed S. A comparison of the construct validity of two burnout measures in two groups of professionals. *Int J Stress Manag.* 2006;13:176-200.
9. Kroenke K, Spitzer RL. The PHQ-9: a new depression diagnostic and severity measure. *Psychiatr Ann.* 2002;32:509-515.
10. Tennant C. Work-related stress and depressive disorders. *J Psychosom Res.* 2001;51:697-704.
11. Hintsala T, Elovainio M, Jokela M, et al. Is there an independent association between burnout and increased allostatic load? Testing the contribution of psychological distress and depression. *J Health Psychol.* 2016;21:1576-1586.
12. Bianchi R, Schonfeld IS. Burnout is associated with a depressive cognitive style. *Person Indiv Dif.* 2016 Jan 15. [Epub ahead of print].