City University of New York (CUNY)

CUNY Academic Works

Publications and Research

City College of New York

1982

Nonperturbative proof of the non-Abelian anomalies

A. P. Balachandran
Syracuse University

G. Marmo
Syracuse University

V. P. Nair
CUNY City College

C. G. Trahern
Syracuse University

[How does access to this work benefit you? Let us know!](#)

More information about this work at: https://academicworks.cuny.edu/cc_pubs/378

Discover additional works at: <https://academicworks.cuny.edu>

This work is made publicly available by the City University of New York (CUNY).
Contact: AcademicWorks@cuny.edu

Nonperturbative proof of the non-Abelian anomalies

A. P. Balachandran, G. Marmo,* V. P. Nair, and C. G. Trahern

Physics Department, Syracuse University, Syracuse, New York 13210

(Received 16 September 1981)

We give a nonperturbative derivation of non-Abelian anomalies.

I. INTRODUCTION

The non-Abelian axial anomalies^{1,2} have played an increasingly important role in the study of gauge theories. A nonperturbative derivation of these anomalies however has not appeared in the published literature. In this note, we try to supply such a derivation. (Similar results have been obtained by S. Coleman and B. Grossman³ using different techniques.)

Our approach to the problem relies on some recent work of Vergeles and Fujikawa.⁴ They have pointed out that the anomaly in any axial-vector current is a result of the noninvariance of the fermion functional measure in the path integral under the corresponding axial transformation. It can in fact be expressed in terms of the associated Jacobian determinant. Here, we shall evaluate this determinant using ζ -function regularization.⁵ The final results agree with those of Bardeen.¹

In Sec. II we recall the work of Ref. 4 and write down the formal expression for the anomalies. The Wess-Zumino consistency conditions⁶ are derived from the properties of Jacobians. In Sec. III the continuation of the chiral group $U_L(N) \times U_R(N)$ to Euclidean space is discussed. We find that it must be continued to $GL(N, C)$ to maintain the Hermiticity of the Dirac operator. In Sec. IV we outline the evaluation of the Jacobian. Section V concludes the paper with some brief comments.

II. THE ANOMALY AS A JACOBIAN

We consider the gauge theory of N massless Dirac fields ψ^a ($a=1, 2, \dots, N$) where the gauged group G is the chiral group $U_L(N) \times U_R(N)$. The field $\psi=(\psi^1, \psi^2, \dots, \psi^N)$ transforms according to the representation $(N, 1) + (1, N)$ of this group. A basis for the Lie algebra $\underline{G}$ of G in this representation is $i\lambda^a, i\gamma_5\lambda^a$ where λ^a are the Gell-Mann matrices. The vector and axial-vector gauge fields are

$$V_\mu = -iV_\mu^a \lambda^a, \quad A_\mu = -iA_\mu^a \lambda^a \quad (2.1)$$

so that the covariant derivative of ψ is

$$D_\mu \psi \equiv (\partial_\mu + V_\mu + \gamma_5 A_\mu) \psi. \quad (2.2)$$

According to Ref. 4, the fermion functional measure $d\psi d\bar{\psi}$ is not in general invariant under the gauge transformations associated with G . Thus if

$$\psi^\pm = \exp \left[\frac{1 \pm \gamma_5}{2} \lambda^a \theta_a \right] \psi, \quad \theta_a = \theta_a(x), \quad (2.3)$$

then (Tr will denote trace over Dirac and internal indices as well as over x , while tr will denote trace over Dirac and/or internal indices only)

$$\begin{aligned} d\psi^\pm d\bar{\psi}^\pm &= d\psi d\bar{\psi} \text{Det}[e^{\mp i \gamma_5 \lambda^a \theta_a}] \\ &= d\psi d\bar{\psi} e^{\mp i \text{Tr} \lambda^a \theta_a \gamma_5}. \end{aligned} \quad (2.4)$$

This anomalous transformation law is the cause of the anomalous nonconservation laws

$$\begin{aligned} \int d^4x \theta_a(x) \partial^\mu l_\mu^a &= \text{Tr} \lambda^a \theta_a \gamma_5, \\ \int d^4x \theta_a(x) \partial^\mu r_\mu^a &= -\text{Tr} \lambda^a \theta_a \gamma_5 \end{aligned} \quad (2.5)$$

of the currents l_μ^a and r_μ^a for the left-handed and right-handed transformations.

As a prelude to the derivation of the Wess-Zumino consistency conditions, we note that the trace in (2.4) depends on the potentials. Thus $d\psi d\bar{\psi}$ also depends on the potentials. We exhibit this fact by writing

$$\begin{aligned} d\psi d\bar{\psi} &= d\mu(\psi, W), \\ W_\mu &= V_\mu + \gamma_5 A_\mu. \end{aligned} \quad (2.6)$$

Let us also denote the gauge transform of W by an element g of the gauge group as $g \circ W$. For instance, if

$$\begin{aligned} g &= e^\eta, \\ \eta &= i \frac{1 + \gamma_5}{2} \lambda^a \theta_a + i \frac{1 - \gamma_5}{2} \lambda^a \phi_a, \end{aligned} \quad (2.7)$$

then

$$(g \circ W)_\mu \equiv g W_\mu g^{-1} + g \partial_\mu g^{-1}. \quad (2.8)$$

The transformation law for the functional measure can now be expressed as

$$d\mu(g\psi, g \circ W) = d\mu(\psi, W) e^{\beta(g, W)}, \quad (2.9)$$

where for (2.7),

$$\beta(g, W) = -i \text{Tr} \lambda^a (\theta_a - \phi_a) \gamma_5. \quad (2.10)$$

Since

$$d\mu((gg')\psi, (g \circ g') \circ W) = d\mu(g(g'\psi), g \circ (g' \circ W)) \quad (2.11)$$

we find

$$\beta(gg', W) = \beta(g, g' \circ W) + \beta(g', W) \quad (2.12)$$

(modulo 2π).

Equation (2.12) is an integrated form of the Wess-Zumino conditions. To derive the latter, we start from the identity

$$\begin{aligned} \beta(gg'g^{-1}, W) &= \beta(g, g' \circ g^{-1} \circ W) \\ &+ \beta(g', g^{-1} \circ W) + \beta(g^{-1}, W) \end{aligned} \quad (2.13)$$

which follows from (2.12). Set $g = e^\eta$, $g' = e^\epsilon$ and consider η and ϵ to be small. Then up to leading terms,

$$\beta(1 + [\eta, \epsilon] + \dots, W) = \delta_\eta \beta(1 + \epsilon + \dots, W) - \delta_\epsilon \beta(1 + \eta + \dots, W), \quad (2.14)$$

where $\delta_\epsilon \beta$ is the change in β due to an infinitesimal gauge transformation with parameter ϵ :

$$\delta_\epsilon \beta(s, W) = \text{term linear in } \epsilon \text{ in } \beta(s, W - [\epsilon, W] - \partial\epsilon) - \beta(s, W). \quad (2.15)$$

Now, for any group element s of the form $1 + \xi + O(\xi^2)$, $\beta(s, W)$ and $\beta(e^\xi, W)$ are the same up to the linear term in ξ and are given by (2.10) and (2.7). Thus (2.14) is just the Wess-Zumino condition.

III. THE DIRAC OPERATOR AND CHIRAL GROUP IN EUCLIDEAN SPACE

The regularization of the trace in (2.4) will involve us in the consideration of the Euclidean-space Dirac operator

$$\gamma_\lambda D_\lambda = \gamma_\lambda (\partial_\lambda + V_\lambda + \gamma_5 A_\lambda) \quad (3.1)$$

in the Hilbert space of functions with scalar product

$$(\psi_1, \psi_2) = \int d^4x \psi_1^\dagger(x) \psi_2(x). \quad (3.2)$$

Our manipulations with the heat kernel will require this operator to be anti-Hermitian. The anti-Hermiticity of $\gamma \cdot D$ implies the properties

$$V_\lambda^\dagger = -V_\lambda, \quad A_\lambda^\dagger = A_\lambda \quad (3.3)$$

of V_λ and A_λ in Euclidean space. Thus the Hermiticity property of A_λ has to be changed in the passage from Minkowski to Euclidean space.

If this Hermiticity property is to be preserved by gauge transformations in Euclidean space, then in (2.7) we have to regard $\theta_a + \phi_a$ as real and $\theta_a - \phi_a$ as pure imaginary when we work in Euclidean

space. This changes the group from $U_L(N) \times U_R(N)$ to $GL(N, C)$. This change is similar to the change that affects the group acting on the spinor index of a Dirac field in going from Minkowski to Euclidean space: the group is $SL(2, C)$ in Minkowski and $SU(2) \times SU(2)$ in Euclidean space.

If the gauged group G in Minkowski space is an arbitrary subgroup of $U_L(N) \times U_R(N)$, then we do not know how to continue it to Euclidean space. For instance if $G = U_L(N)$, then in (2.7), $\phi_a = 0$. The natural Euclidean continuation of η (with $\phi_a = 0$) is $i[\frac{1}{2}(1 + i\gamma_5)]\lambda^a \theta_a$ (θ_a being real). However $i[\frac{1}{2}(1 + i\gamma_5)]\lambda^a$ do not span a Lie algebra.

IV. EVALUATION OF THE ANOMALY

We follow standard methods for the evaluation of the anomaly $\text{Tr} \lambda^a \theta_a \gamma_5$.⁵ It is continued to Euclidean space (where $\theta_a^* = -\theta_a$) and evaluated by ζ -function regularization. It is finally continued back to Minkowski space (where $\theta_a^* = \theta_a$).

We now quickly review the ζ -function regularization method. In Euclidean space, let ϕ_n be the eigenfunctions of $\gamma \cdot D$:

$$\begin{aligned} \gamma \cdot D \phi_n &= i \lambda_n \phi_n, \\ (\phi_n, \phi_m) &= \delta_{nm}, \\ \sum_n \phi_n(x) \phi_n^\dagger(x') &= \delta^4(x - y) 1. \end{aligned} \quad (4.1)$$

Then (formally),

$$\text{Tr} \lambda^a \theta_a \gamma_5 = \int d^4x \theta_a(x) \lim_{\substack{s \rightarrow 0 \\ x \rightarrow y}} \text{tr} \lambda^a \gamma_5 \xi(s, x, y), \quad (4.2)$$

$$\xi(s, x, y) \equiv \sum_n \frac{\phi_n(x) \phi_n^\dagger(y)}{\lambda_n^{2s}}.$$

The trace on the right side (tr with a lower case t) is only over Dirac and internal indices.

The ξ function can be written in terms of the heat kernel

$$h(t, x, y) = \langle x | e^{t(\gamma \cdot D)^2} | y \rangle : \quad (4.3)$$

$$\xi(s, x, y) = \frac{1}{\Gamma(s)} \int_0^\infty dt t^{s-1} h(t, x, y).$$

The heat kernel has the asymptotic expansion⁵

$$h(t, x, y) = \frac{1}{16\pi^2 t^2} \exp \left[-\frac{(x-y)^2}{4t} \right] \times \sum_{n=0}^\infty a_n(x, y) t^n \quad (4.4)$$

for small t . Inserting (4.4) in (4.3), we find that

$$\xi(0, x, x) = \frac{a_2(x, x)}{16\pi^2}$$

and that

$$\text{Tr} \lambda^a \theta_a \gamma_5 = \frac{1}{16\pi^2} \int d^4x \theta_a(x) \text{tr} \lambda^a \gamma_5 a_2(x), \quad (4.5)$$

$$a_2(x) \equiv a_2(x, x).$$

The coefficients a_n can be evaluated recursively using the heat equation

$$\partial_t h(t, x, y) = (\gamma \cdot D)^2 h(t, x, y) \quad (4.6)$$

and the boundary condition

$$h(0, x, y) = \delta^4(x-y) 1. \quad (4.7)$$

It is known⁵ that if there is a differential operator

$$\mathcal{D}_\mu = \partial_\mu + P_\mu \quad (4.8)$$

such that

$$X = (\gamma \cdot D)^2 - \mathcal{D}^2 \quad (4.9)$$

involves no differential operator, then

$$a_2(x) = \frac{1}{2} X^2 + \frac{1}{12} Y_{\mu\nu} Y_{\mu\nu} + \frac{1}{6} \mathcal{D}^2 X, \quad (4.10)$$

where

$$\mathcal{D}_\mu X = \partial_\mu X + [P_\mu, X]$$

and

$$Y_{\mu\nu} = [\mathcal{D}_\mu, \mathcal{D}_\nu]. \quad (4.11)$$

Thus in our problem, we have to find a P_μ such that X in (4.9) involves no differential operator and then evaluate the anomaly from (4.10) and (4.5). A suitable P_μ and the corresponding X are

$$P_\mu = V_\mu + \gamma_5 A_\mu - \gamma_5 \gamma \cdot A \gamma_\mu, \quad (4.12)$$

$$X = \frac{1}{4} [\gamma_\mu, \gamma_\nu] F_{\mu\nu} + \gamma_5 \{ \partial_\mu (\gamma \cdot A \gamma_\mu) + [V_\mu, \gamma \cdot A] \gamma_\mu \} + 2A^2,$$

where

$$F_{\mu\nu} = V_{\mu\nu} + \gamma_5 A_{\mu\nu},$$

$$V_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu + [V_\mu, V_\nu] + [A_\mu, A_\nu], \quad (4.13)$$

$$A_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [V_\mu, A_\nu] - [V_\nu, A_\mu].$$

The calculation of the anomaly is now straightforward though tedious. We find (here

$$[A, B]_\pm = AB \pm BA)$$

$$\frac{1}{16\pi^2} \text{tr} \lambda^a \gamma_5 a_2(x) = \text{tr} \lambda^a (G^{(1)} + G^{(2)}),$$

$$G^{(1)} = -\frac{1}{8\pi^2} \epsilon_{\mu\nu\alpha\beta} \left[\frac{1}{4} V_{\mu\nu} V_{\alpha\beta} + \frac{1}{12} A_{\mu\nu} A_{\alpha\beta} - \frac{2}{3} (A_\mu A_\nu V_{\alpha\beta} + A_\mu V_{\nu\alpha} A_\beta + V_{\mu\nu} A_\alpha A_\beta) + \frac{8}{3} A_\mu A_\nu A_\alpha A_\beta \right], \quad (4.14)$$

$$G^{(2)} = -\frac{1}{16\pi^2} \left\{ \frac{4}{3} [(D_\mu^\nu A_\nu + D_\nu^\nu A_\mu), A_\mu A_\nu] + -\frac{2}{3} [D^\nu \cdot A, A^2] + \frac{4}{3} [A_\mu, D_\lambda^\nu V_{\mu\lambda}] - \frac{1}{3} [A_{\mu\lambda}, V_{\mu\lambda}] - \right.$$

$$\left. + \frac{2}{3} D_\rho^\nu D_\rho^\nu (D^\nu \cdot A) + 4A_\lambda (D^\nu \cdot A) A_\lambda \right\},$$

$$D_\mu^\nu f \equiv \partial_\mu f + [V_\mu, f]_-.$$

The term involving $G^{(1)}$ is identical to Bardeen's expression.¹ The terms involving $G^{(2)}$ on the other hand can be removed from the divergence equations (2.5) (and hence Ward-Takahashi identities) by following an idea discussed by Bardeen.¹ Thus we add the following counterterm $\Delta\mathcal{L}$ to the Lagrangian density:

$$\begin{aligned}\Delta\mathcal{L} = & -\frac{1}{8\pi^2} \text{tr} \left\{ \frac{2}{3} (D_\mu^\nu A_\nu)^2 \right. \\ & - (D^\nu A)^2 - \frac{2}{3} [A_\mu, A_\nu]_-^2 \\ & \left. + \frac{2}{3} (A_\mu A_\nu A_\mu A_\nu) + \frac{1}{2} V_{\mu\nu}^2 \right\} .\end{aligned}\quad (4.15)$$

Under an axial transformation $\psi \rightarrow e^{i\gamma_5 \lambda^a \theta_a} \psi$, the first-order variation of

$$\int d^4x \Delta\mathcal{L} \quad (4.16)$$

is precisely

$$2 \int d^4x \theta_a(x) \text{tr} \lambda^a G^{(2)}(x) .$$

Thus the infinitesimal variation of

$$d\mu(\psi, W) e^{i \int d^4x \Delta\mathcal{L}}$$

under an axial transformation does not involve $G^{(2)}$. In other words, the Ward-Takahashi identities are not affected by $G^{(2)}$ when the counterterm $\Delta\mathcal{L}$ is added to the Lagrangian density. ($\Delta\mathcal{L}$ is

invariant under the vector gauge group. Therefore, it does not generate anomalies for the vector currents.)

V. CONCLUDING REMARKS

The expression in (4.14) for the anomaly fulfills the consistency conditions. This follows from the fact that the contribution from $G^{(1)}$ can be verified to do so by an explicit calculation,⁶ while the contribution from $G^{(2)}$ does so since it can be reproduced by the variation of the action (4.16) (cf. Ref. 6). We feel that there should be a proof of such results which uses only the formal properties of the ζ -function regularization and the heat equation. We also feel that the potential P_μ in (4.12) should be capable of a simple interpretation. Our attempts at the resolution of these problems have not however been successful.

ACKNOWLEDGMENTS

We thank P. P. Divakaran, S. Rajeev, and J. Schechter for discussions. G. Marmo thanks Joshua Goldberg and Kameshwar Wali for their warm hospitality at Syracuse University where this work was done. The work of A.P.B., V.P.N., and C.G.T. was supported by the U.S. Department of Energy under Contract No. DE-AC02-76ER03533. The work of G.M. was supported by INFN, Italy.

*Permanent address: Istituto di Fisica Teorica, Università di Napoli, Mostra d'Oltremare Pad. 19, 80125 Napoli, Italy.

¹W. A. Bardeen, Phys. Rev. **184**, 1848 (1969).

²D. Gross and R. Jackiw, Phys. Rev. D **6**, 477 (1972); H. Georgi and S. L. Glashow, *ibid.* **6**, 429 (1972).

³S. Coleman and B. Grossman (unpublished).

⁴S. N. Vergeles, quoted in A. A. Migdal, Phys. Lett.

81B, 37 (1979); K. Fujikawa, Phys. Rev. Lett. **44**, 1733 (1980); Phys. Rev. D **21**, 2848 (1980).

⁵N. K. Nielsen, NORDITA Report No. 78/24, 1978 (unpublished); N. K. Nielsen, M. T. Grisaru, H. Römer, and P. van Nieuwenhuizen, Nucl. Phys. **B140**, 477 (1978) and references therein.

⁶J. Wess and B. Zumino, Phys. Lett. **37B**, 95 (1971).