

City University of New York (CUNY)

CUNY Academic Works

Computer Science Technical Reports

CUNY Academic Works

2014

TR-2014009: Estimating the Norms of Random Circulant and Toeplitz Matrices and Their Inverses II

Victor Y. Pan

Guoliang Qian

John Svadlenka

Liang Zhao

[How does access to this work benefit you? Let us know!](#)

More information about this work at: https://academicworks.cuny.edu/gc_cs_tr/400

Discover additional works at: <https://academicworks.cuny.edu>

This work is made publicly available by the City University of New York (CUNY).
Contact: AcademicWorks@cuny.edu

Estimating the Norms of Random Circulant and Toeplitz Matrices and Their Inverses II *

Victor Y. Pan^{[1],[2],[a]}, Guoliang Qian^{[2],[b]}, John Svadlenka^{[2],[c]}, and Liang Zhao^{[2],[d]}

^[1] Department of Mathematics and Computer Science
Lehman College of the City University of New York
Bronx, NY 10468 USA

^[2] Ph.D. Programs in Mathematics and Computer Science
The Graduate Center of the City University of New York
New York, NY 10036 USA

^[a] victor.pan@lehman.cuny.edu

<http://comet.lehman.cuny.edu/vpan/>

^[b] gqian@gc.cuny.edu

^[c] svajoth@verizon.net

^[d] lzhaol@gc.cuny.edu

Abstract

We combine some basic techniques of linear algebra with some expressions for Toeplitz and circulant matrices and the properties of Gaussian random matrices to estimate the norms of Gaussian Toeplitz and circulant random matrices and their inverses. In the case of circulant matrices we obtain sharp probabilistic estimates, which show that the matrices are expected to be very well conditioned. Our probabilistic estimates for the norms of standard Gaussian Toeplitz random matrices are within a factor of $\sqrt{2}$ from those in the circulant case. We also achieve partial progress in estimating the norm of the Toeplitz inverse. Namely we yield reasonable probabilistic upper estimates assuming certain bounds on the absolute values of two corner entries of the inverse. Empirically we observe that the condition numbers of Toeplitz and general random matrices tend to be of the same order. As the matrix size grows, these numbers grow equally slowly, although faster than in the case of circulant random matrices.

2000 Math. Subject Classification: 15A52, 15A12, 65F22, 65F35

Key Words: Matrix norms, Condition numbers, Toeplitz matrices, Circulant matrices, Random matrices

1 Introduction

Estimating the condition numbers $\kappa(A) = \|A\| \|A^{-1}\|$ of structured random matrices A is a well known challenge (cf. [SST06]), linked to the design of efficient randomized matrix algorithms, e.g., in the papers [HMT11], [XXG12], [PQZ13]. We seek such estimates for Gaussian random Toeplitz

*Some results of this paper have been presented at the ACM-SIGSAM International Symposium on Symbolic and Algebraic Computation (ISSAC '2011), San Jose, CA, 2011, the 7th International Congress on Industrial and Applied Mathematics (ICIAM 2011), in Vancouver, British Columbia, Canada, July 18-22, 2011, the SIAM International Conference on Linear Algebra, in Valencia, Spain, June 18-22, 2012, and the 18th Conference of the International Linear Algebra Society (ILAS'2013), Providence, RI, 2013

and circulant matrices, mostly by employing and combining some basic techniques of linear algebra and the Toeplitz and circulant matrix structure. In the case of circulant matrices A our sharp probabilistic estimates for the norms $\|A\|$ and $\|A^{-1}\|$ show that the matrices are expected to be very well conditioned. In the case of $n \times n$ Toeplitz matrices A our estimates for the norm $\|A\|$ are within a factor of $\sqrt{2}$ from the bounds in the circulant case. Estimating the norms $\|A^{-1}\|$ turns out to be a harder problem. We obtain reasonable probabilistic upper bounds on that norm, and consequently on the condition number $\kappa(A) = \|A\| \|A^{-1}\|$, under the assumption that the norms of the first row and the first column of the inverse do not exceed dramatically the absolute values of their two corner entries, $(A^{-1})_{1,n}$ and $(A^{-1})_{n,1}$, respectively. For some large and important special classes of $n \times n$ Toeplitz matrices the condition numbers grow exponentially in n as $n \rightarrow \infty$ [BG05], but in our tests with both random general and Toeplitz random matrices their condition numbers consistently grew with the same reasonably slow rate as n grew large, although the growth was still significantly faster than in the case of circulant matrices. Our study can be immediately extended to the cases of factor circulant and Hankel matrices and partly to the case of other than Gaussian probability distributions. In particular we estimated empirically the norms of random circulant and Toeplitz matrices and of their inverses defined under the standard Gaussian and uniform probability distributions, and observed similar results under both distributions (see Section 8). Finally we note that random unitary circulant $n \times n$ matrices can be defined by n random real parameters, chosen under any probability distribution in any real range (cf. [PQYa, Section 5.4]).

We organize our paper as follows. We recall some definitions and basic results on general matrix computations in the next section and on Toeplitz and circulant matrices in Section 3. In Section 4 we deduce sharp estimates on the norms of circulant and Toeplitz matrices in terms of their generating vectors. In Section 5 we similarly estimate the norms of the inverses of these matrices. We recall the definition of Gaussian random matrices and some basic facts about them in Section 6. In Section 7 we extend our estimates of Sections 4 and 5 to standard Gaussian Toeplitz and circulant random matrices and their inverses. In Section 8 we cover numerical tests, which are the contribution of the last four co-authors. In the Appendix we discuss nondegeneration of random matrices.

2 Some definitions and basic results

In this section we recall some customary definitions and basic properties of matrix computations [GL96], [S98]. A^T is the transpose of a matrix A . A^H is its Hermitian transpose. $A^H = A^T$ for a real matrix A . $A^T = A$ if A is a real symmetric matrix, $A^H = A$ if A is a Hermitian matrix. $D = \text{diag}(d_i)_{i=0}^{n-1}$ is a diagonal matrix with diagonal entries d_0, \dots, d_{n-1} .

A square matrix A is unitary if $A^H A = A A^H = I$, and is orthogonal if $A^T A = A A^T = I$. For a vector $\mathbf{v} = (v_i)_{i=0}^{n-1}$ define the norms $\|\mathbf{v}\|_1 = \sum_i |v_i|$, $\|\mathbf{v}\|_2 = (\sum_i |v_i|^2)^{1/2}$, and $\|\mathbf{v}\|_\infty = \max_i |v_i|$. For a matrix $A = (a_{ij})_{i,j=0}^{n-1}$ define its s -norms $\|A\|_s = \inf_{\|\mathbf{v}\|_s=1} \|A\mathbf{v}\|_s$ for $s = 1, 2, \infty$ and its Frobenius norm $\|A\|_F = (\sum_{i,j=0}^{n-1} |a_{ij}|^2)^{1/2}$. We write $\|\mathbf{v}\| = \|\mathbf{v}\|_2$ and $\|A\| = \|A\|_2$. $\|A\|$ is called the spectral norm of a matrix A . Assume an $n \times n$ matrix A , a diagonal matrix $D = \text{diag}(d_i)_{i=0}^{n-1}$, and two unitary $n \times n$ matrices U and V . Then

$$\|A\|_1 = \|A^T\|_\infty = \|A^H\|_\infty = \max_j \sum_{i,j} |a_{ij}|, \quad (2.1)$$

$$\frac{1}{\sqrt{n}} \|A\|_s \leq \|A\| \leq \sqrt{n} \|A\|_s \text{ for } s = 1 \text{ and } s = \infty, \quad \|A\|^2 \leq \|A\|_1 \|A\|_\infty, \quad (2.2)$$

$$\|D\| = \|D\|_1 = \|D\|_\infty = \max_{i=1}^n \{ |d_i| \}, \quad \|D\|_F^2 = \sum_{i=1}^n |d_i|^2, \quad (2.3)$$

$$\|UAV\| = \|A\| \text{ and } \|UAV\|_F = \|A\|_F, \quad (2.4)$$

$$\|A\| \leq \|A\|_F \leq \sqrt{\rho} \|A\| \text{ where } \rho = \text{rank}(A), \quad (2.5)$$

$$\|A + B\|_s \leq \|A\|_s + \|B\|_s, \quad \|FG\|_s \leq \|F\|_s \|G\|_s \text{ for } s = 1, 2, \infty. \quad (2.6)$$

For the latter bounds the matrix sizes must match, to define the matrices $A + B$ and FG .

$\kappa(A) = \|A\| \|A^{-1}\|$ is the *condition number* of a nonsingular matrix A . A matrix is *ill conditioned* if its condition number is large, and it is *well conditioned* if this number is reasonably bounded.

3 Toeplitz and f -circulant matrices and their inverses

\mathbf{e}_i is the i th coordinate vector of dimension n for $i = 1, \dots, n$. Two vectors

$$\mathbf{t}_+ = (t_i)_{i=1-n}^{n-1} \text{ and } \mathbf{t} = (t_i)_{i=0}^{n-1} \quad (3.1)$$

define a *Toeplitz* $n \times n$ matrix $T_n = T(\mathbf{t}_+) = (t_{i-j})_{i,j=0}^{n-1}$, a lower *triangular Toeplitz* $n \times n$ matrix $Z(\mathbf{t}) = (t_{i-j})_{i,j=0}^{n-1}$ (where $t_k = 0$ for $k < 0$), and its transpose $(Z(\mathbf{t}))^T$. Now consider the $n \times n$ matrices $Z = Z_0 = Z(\mathbf{e}_2)$ of downshift and $Z_f = Z + f\mathbf{e}_1^T \mathbf{e}_n$ of f -*circulant shift* for $f \neq 0$ (see equation (3.2) below). Then $Z\mathbf{t} = (t_{i-1})_{i=0}^{n-1}$ and $Z(\mathbf{t}) = Z_0(\mathbf{t}) = \sum_{i=0}^{n-1} t_i Z^i$. An f -*circulant* (or factor circulant) matrix $Z_f(\mathbf{t}) = \sum_{i=0}^{n-1} t_i Z_f^i$ is a special Toeplitz $n \times n$ matrix defined by its first column vector $\mathbf{t} = (t_i)_{i=0}^{n-1}$ and a scalar f . f -circulant matrix is called *circulant* where $f = 1$.

$$T_n = \begin{pmatrix} t_0 & t_{-1} & \cdots & t_{1-n} \\ t_1 & t_0 & \ddots & \vdots \\ \vdots & \ddots & \ddots & t_{-1} \\ t_{n-1} & \cdots & t_1 & t_0 \end{pmatrix}, \quad Z = \begin{pmatrix} 0 & \cdots & 0 \\ 1 & \ddots & & \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 \end{pmatrix}, \quad Z_f = \begin{pmatrix} 0 & \cdots & f \\ 1 & \ddots & & \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 \end{pmatrix}. \quad (3.2)$$

$J = J_n = (\mathbf{e}_n \mid \dots \mid \mathbf{e}_1)$ is the reflection matrix of the size $n \times n$ where $J = J^T = J^{-1}$. A Toeplitz matrix T_n and its inverse (if defined) are *persymmetric*, that is,

$$JT_n J = T_n \text{ and } JT_n^{-1} J = T_n^{-1}. \quad (3.3)$$

$\omega = \exp(2\pi\sqrt{-1}/n)$ denotes a primitive n th root of unity. $\Omega = (\omega^{ij})_{i,j=0}^{n-1}$ is the matrix of the discrete Fourier transform at n points.

Theorem 3.1. $\Omega^H \Omega = nI$, that is, $\frac{1}{\sqrt{n}}\Omega$ is a unitary matrix.

The following theorem implies that the inverses (wherever they are defined) and pairwise products of f -circulant matrices are f -circulant.

Theorem 3.2. (See [CPW74].) We have $Z_1(\mathbf{t}) = \Omega^{-1}D(\Omega\mathbf{t})\Omega$. More generally, for any $f \neq 0$, we have $Z_f^n(\mathbf{t}) = U_f^{-1}D(U_f\mathbf{t})U_f$ where $U_f = \Omega D(\mathbf{f})$, $\mathbf{f} = (f^i)_{i=0}^{n-1}$, $D(\mathbf{u}) = \text{diag}(u_i)_{i=0}^{n-1}$ for a vector $\mathbf{u} = (u_i)_{i=0}^{n-1}$.

Theorem 3.3. Write $T_k = (t_{i-j})_{i,j=0}^{k-1}$ for $k = n, n+1$.

(a) Let the matrix T_n be nonsingular and write $\mathbf{p} = T_n^{-1}\mathbf{e}_1$ and $\mathbf{q} = T_n^{-1}\mathbf{e}_n$. If $p_1 = \mathbf{e}_1^T \mathbf{p} \neq 0$, then $p_1 T_n^{-1} = Z(\mathbf{p})(Z(J\mathbf{q}))^T - Z(Z\mathbf{q})(Z(Z\mathbf{J}\mathbf{p}))^T$.

In parts (b) and (c) below let a Toeplitz $(n+1) \times (n+1)$ matrix T_{n+1} be nonsingular and write $\hat{\mathbf{v}} = (v_i)_{i=0}^n = T_{n+1}^{-1}\mathbf{e}_1$, $\mathbf{v} = (v_i)_{i=0}^{n-1}$, $\mathbf{v}' = (v_i)_{i=1}^n$, $\hat{\mathbf{w}} = (w_i)_{i=0}^n = T_{n+1}^{-1}\mathbf{e}_{n+1}$, $\mathbf{w} = (w_i)_{i=0}^{n-1}$, and $\mathbf{w}' = (w_i)_{i=1}^n$.

(b) If $v_0 \neq 0$, then the matrix T_n is nonsingular and $v_0 T_n^{-1} = Z(\mathbf{v})(Z(J\mathbf{w}'))^T - Z(\mathbf{w})(Z(J\mathbf{v}'))^T$.

(c) If $v_n \neq 0$, then the matrix $T_{1,0} = (t_{i-j})_{i=1,j=0}^{n,n-1}$ is nonsingular and $v_n T_{1,0}^{-1} = Z(\mathbf{w})(Z(J\mathbf{v}'))^T - Z(\mathbf{v})(Z(J\mathbf{w}'))^T$.

Proof. See [GS72] on parts (a) and (b); see [GK72] on part (c). □

4 The norm bounds for circulant and Toeplitz matrices

Theorem 4.1. *Assume a pair of vectors \mathbf{t} and \mathbf{t}_+ of (3.1), defining the circulant and Toeplitz $n \times n$ matrices $Z_1(\mathbf{t})$ and $T_n = T(\mathbf{t}_+)$. Then*

$$\|Z(\mathbf{t})\|_s \leq \|Z_1(\mathbf{t})\|_s \text{ for } s = F, 1, 2, \infty, \quad (4.1)$$

$$\|Z_1(\mathbf{t})\| \leq \|Z_1(\mathbf{t})\|_1 = \|Z_1(\mathbf{t})\|_\infty = \|\mathbf{t}\|_1 \leq \sqrt{n} \|\mathbf{t}\|, \quad (4.2)$$

$$\|Z_1(\mathbf{t})\| = \|\Omega\mathbf{t}\|_\infty, \quad \|Z_1(\mathbf{t})\|_F = \sqrt{n} \|\mathbf{t}\|, \quad (4.3)$$

$$\|T_n\| \leq \|T_n\|_1 = \|T_n\|_\infty \leq \|\mathbf{t}_+\|_1 \leq \sqrt{2n-1} \|\mathbf{t}_+\| \text{ and } \|T_n\|_F \leq \sqrt{2n-1} \|\mathbf{t}_+\|. \quad (4.4)$$

Proof. Readily verify equation (4.1). Note that $\frac{1}{\sqrt{n}}\Omega$ and $\sqrt{n}\Omega^{-1}$ are unitary matrices. Combine equations (2.1) and (2.2) and obtain that $\|Z_1(\mathbf{t})\| = \|D(\Omega\mathbf{t})\| = \|\Omega\mathbf{t}\|_\infty$, whereas $\|Z_1(\mathbf{t})\|_F^2 = \|D(\Omega\mathbf{t})\|_F^2 = \|\Omega\mathbf{t}\|^2 = n\|\mathbf{t}\|^2$. This proves relationships (4.2). Combine Theorems 3.1 and 3.2 with relationships (2.3) and (2.4) and obtain that $\|Z_1(\mathbf{t})\|_F^2 = \|D(\Omega\mathbf{t})\|_F^2 = \|(\Omega\mathbf{t})\|^2 = n\|\mathbf{t}\|^2$, yielding equation (4.3). Embed the $n \times n$ matrix T_n into the circulant $(2n-1) \times (2n-1)$ matrix $Z_1(\mathbf{t}_+)$ and obtain that $\|T_n\|_s \leq \|Z_1(\mathbf{t}_+)\|_s$ where s can stand for $F, 1, 2$, or ∞ . Together with relationships (4.2) and (4.3), this implies relationships (4.4). \square

Remark 4.1. Extension to the case of f -circulant matrices. *Theorem 3.2 implies that*

$$\frac{1}{g(f)} \|Z_1(\mathbf{v})\| \leq \|Z_f(\mathbf{v})\| \leq g(f) \|Z_1(\mathbf{v})\|$$

and if the matrices $Z_1(\mathbf{v})$ and $Z_f(\mathbf{v})$ are nonsingular, then also

$$\frac{1}{g(f)} \|Z_1(\mathbf{v})^{-1}\| \leq \|Z_f(\mathbf{v})^{-1}\| \leq g(f) \|Z_1(\mathbf{v})^{-1}\|$$

for all vectors \mathbf{v} , scalars $f \neq 0$, $g(f) = \max\{|f|, 1/|f|\}$, and $j = 1, \dots, n$. Therefore we can readily extend our norm estimates from circulant to f -circulant matrices for $f \neq 0$. In particular $\|Z_f(\mathbf{v})\| = \|Z_1(\mathbf{v})\|$ and $\|Z_f(\mathbf{v})^{-1}\| = \|Z_1(\mathbf{v})^{-1}\|$ where $|f| = 1$.

Remark 4.2. Extension to the case of Hankel matrices. *All our estimates for Toeplitz matrices are immediately extended to the case of Hankel matrices $H_n = (h_{i+j})_{i,j=0}^{n-1}$ because the products $H_n J = T_n$ and $J H_n = T_n$ are $n \times n$ Toeplitz matrices.*

5 The norm bounds for the inverses of circulant and Toeplitz matrices

Hereafter we write $((v_i)_{i=0}^{n-1})^{-1} = (1/v_i)_{i=0}^{n-1}$. Similarly to equation (4.3) deduce that

$$\|(Z_1(\mathbf{t}))^{-1}\|_F = \|(D(\Omega\mathbf{t}))^{-1}\|_F = \|(\Omega\mathbf{t})^{-1}\|, \quad \|(Z_1(\mathbf{t}))^{-1}\| = \|(D(\Omega\mathbf{t}))^{-1}\| = \|(\Omega\mathbf{t})^{-1}\|_\infty. \quad (5.1)$$

Combine the latter bound with (2.2) and obtain that

$$\|(Z_1(\mathbf{t}))^{-1}\|_s \leq \sqrt{n} \|(\Omega\mathbf{t})^{-1}\|_\infty \text{ for } s = 1 \text{ and } s = \infty.$$

Theorem 5.1. *Under the assumptions of part (a) of Theorem 3.2 it holds that $\|p_1 T_n^{-1}\|_s \leq 2\|\mathbf{p}\|_1 \|\mathbf{q}\|_1 \leq 2n\|\mathbf{p}\| \|\mathbf{q}\|$ for $s = 1, 2, \infty$.*

Proof. Recall from part (a) of Theorem 3.3 that $p_1 T_n^{-1} = Z(\mathbf{p})(Z(J\mathbf{q}))^T - Z(Z\mathbf{q})(Z(J\mathbf{p}))^T$. Therefore (cf. (2.6)) $\|p_1 T_n^{-1}\|_s \leq \|Z(\mathbf{p})\|_s \|Z(J\mathbf{q})^T\|_s + \|Z(Z\mathbf{q})\|_s \|Z(ZJ\mathbf{p})^T\|_s$ for $s = 1, 2, \infty$. Combine this bound with relationships (4.1) and (4.2) and deduce that $\|p_1 T_n^{-1}\|_s \leq \|\mathbf{p}\|_1 \|J\mathbf{q}\|_1 + \|Z\mathbf{q}\|_1 \|ZJ\mathbf{p}\|_1$. Note that $\|J\mathbf{w}\|_1 = \|\mathbf{w}\|_1$ and $\|Z\mathbf{w}\|_1 \leq \|\mathbf{w}\|_1$ for every vector \mathbf{w} and obtain Theorem 5.1. \square

6 Gaussian random matrices

By extending the norm bounds of the two previous sections we can estimate the norms of circulant and Toeplitz matrices with random entries selected under various probability distributions. We are going to do this just under the Gaussian distribution, which enables simpler and stronger estimates.

Definition 6.1. $F_\gamma(y) = \text{Probability}\{\gamma \leq y\}$ is the cumulative distribution function (cdf) of a real random variable γ evaluated at a real point y . $F_{g(\mu,\sigma)}(y) = \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^y \exp(-\frac{(x-\mu)^2}{2\sigma^2})dx$ for a Gaussian random variable $g(\mu,\sigma)$ with a mean μ , a positive variance σ^2 , and a positive standard deviation σ .

It follows that

$$|y - \mu|/\sigma \geq t \text{ with the probability } \sqrt{\frac{2}{\pi}} \int_t^{+\infty} \exp(-\frac{x^2}{2})dx, \quad (6.1)$$

that is, the probability decays very fast as t grows large.

Definition 6.2. A matrix (or a vector) is a Gaussian random matrix (or vector) with a mean μ and a positive variance σ^2 if it is filled with independent identically distributed (hereafter we write i.i.d.) Gaussian random variables, all having the mean μ and variance σ^2 . $\mathcal{G}_{\mu,\sigma}^{m \times n}$ is the set of such Gaussian $m \times n$ random matrices, which are standard for $\mu = 0$ and $\sigma^2 = 1$. By restricting this set to Toeplitz or f -circulant matrices we obtain the sets $\mathcal{T}_{\mu,\sigma}^{m \times n}$ and $\mathcal{Z}_{f,\mu,\sigma}^{n \times n}$ of Gaussian random Toeplitz and Gaussian random f -circulant matrices, respectively, which are standard for $\mu = 0$ and $\sigma^2 = 1$.

Definition 6.3. Assume a Gaussian random vector $\mathbf{v} = (v_i)_{i=0}^{n-1} \in \mathcal{G}_{\mu,\sigma}^{n \times 1}$. Write $\alpha_{\mu,\sigma,n}(y)$, $\beta_{\mu,\sigma,n}(y)$ and $\chi_{\mu,\sigma,n}(y)$ to denote the cdfs of the norms $\|\mathbf{v}\|_\infty = \max_{i=0}^{n-1} |v_i|$, $\|\mathbf{v}\|_1 = \sum_{i=0}^{n-1} |v_i|$, and $\|\mathbf{v}\| = (\sum_{i=0}^{n-1} v_i^2)^{1/2}$, respectively. Then (cf. (2.2)) $\alpha_{\mu,\sigma,n}(y) \leq \chi_{\mu,\sigma,n}(y) \leq \alpha_{\mu,\sigma,n}(y/\sqrt{n})$, $\beta_{\mu,\sigma,n}(y) \leq \chi_{\mu,\sigma,n}(y) \leq \beta_{\mu,\sigma,n}(y/\sqrt{n})$, and $\chi_{0,1,n}(y) = \frac{2}{2^{n/2}\Gamma(n/2)} \int_0^y x^{n-1} \exp(-x^2/2)dx$ for $y \geq 0$ where $\Gamma(h) = \int_0^\infty x^{h-1} \exp(-x)dx$ is the Gamma function, $\Gamma(n+1) = n!$ for nonnegative integers n .

We recall the following basic results.

Lemma 6.1. Suppose G is a Gaussian matrix, S and T are square orthogonal matrices, and the products SG and GT are well defined. Then SG and GT are Gaussian matrices.

Lemma 6.2. [SST06, Lemma A.2]. For a nonnegative scalar y , a unit vector $\mathbf{t} \in \mathbb{R}^{n \times 1}$, and a vector $\mathbf{b} \in \mathcal{G}_{\mu,\sigma}^{n \times 1}$, we have $F_{|\mathbf{t}^T \mathbf{b}|}(y) \leq \sqrt{\frac{2}{\pi}} \frac{y}{\sigma}$.

Remark 6.1. The latter bound is independent of μ and n and holds for any μ even if all coordinates of the vector \mathbf{b} are fixed except for a single coordinate in $\mathcal{G}_{\mu,\sigma}$.

Gaussian random general, Toeplitz and circulant matrices have full rank with probability 1 (see Appendix A). Hereafter dealing with them we always assume that they have full rank.

7 Norm bounds for Gaussian random circulant and Toeplitz matrices and their inverses

7.1 The norms of Gaussian random Toeplitz and circulant matrices

Theorem 7.1. Assume a Gaussian random vector $\mathbf{t} \in \mathcal{G}_{\mu,\sigma}^{n \times n}$ and n i.i.d. Gaussian random variables $g_i \in \mathcal{G}_{\mu,\sigma}^{1 \times 1}$, $i = 0, \dots, n-1$. Then $\|Z_1(\mathbf{t})\| = \|\Omega \mathbf{t}\|_\infty = \sqrt{n} \max_{i=0}^{n-1} |g_i|$.

Proof. Combine (2.4) and Lemma 6.1. □

Combine our estimates of Theorems 4.1 and 7.1 with Definition 6.2 and obtain the following upper bounds on the norms of Gaussian random Toeplitz and circulant matrices.

Corollary 7.1. For a Gaussian random Toeplitz $n \times n$ matrix T_n and a Gaussian random circulant $n \times n$ matrix $Z_1(\mathbf{t})$ it holds that $F_{\|T_n\|_s}(y) \geq \chi_{\mu,\sigma,2n-1}(y/\sqrt{2n-1})$, $F_{\|Z_1(\mathbf{t})\|}(y) \geq \alpha_{\mu,\sigma,n}(y/\sqrt{n})$, $F_{\|Z_1(\mathbf{t})\|_F}(y) \geq \beta_{\mu,\sigma,n}(y/\sqrt{n})$, and $F_{\|Z_1(\mathbf{t})\|_h}(y) \geq \chi_{\mu,\sigma,n}(y/\sqrt{n})$ where h can stand for 1 or ∞ .

7.2 The expected norms of the inverses of Gaussian random matrices have order at least $1/\sigma$ where the mean μ greatly exceeds the standard deviation σ

Bounds (6.1) imply that the matrix $M \in \mathcal{G}_{\mu,\sigma}^{n \times n}$ is expected to be approximated within the norm bound of order σ by the matrix $\mu \mathbf{e} \mathbf{e}^T$ of rank 1, for the vector $\mathbf{e} = (1, \dots, 1)^T$ provided that $|\mu| \gg \sigma$. If indeed so, then the norm $\|M^{-1}\|$ has order of at least $1/\sigma$. A similar argument applies to Gaussian random Toeplitz and circulant matrices. In the next section we derive our estimates for any pair of (μ, σ) , but one can avoid the latter undesired growth of the norm of the inverse by restricting the study to standard Gaussian random Toeplitz and circulant matrices.

7.3 The norm of the inverse of a Gaussian random circulant matrix

Theorem 7.2. Assume a vector $\mathbf{t} \in \mathcal{G}_{\mu,\sigma}^{n \times 1}$ and n i.i.d. Gaussian random variables $g_i \in G_{\mu,\sigma}^{1 \times 1}$, $i = 0, \dots, n-1$. Then $\|(Z_1(\mathbf{t}))^{-1}\| = \|\mathbf{t}^{-1}\|_\infty / \sqrt{n} = 1/(\sqrt{n} \min_{i=0}^{n-1} |g_i|)$, whereas $\|(Z_1(\mathbf{t}))^{-1}\|_F^2 = \|\mathbf{t}^{-1}\|^2/n = (1/n) \sum_{i=0}^{n-1} (1/g_i^2)$.

Proof. Equation (5.1) implies that $\|(Z_1(\mathbf{t}))^{-1}\| = 1/\min_{i=1}^n |u_i|$ where $\mathbf{u} = (u_i)_{i=1}^n = \Omega \mathbf{t}$. Apply Lemma 6.1 and obtain that \mathbf{u} is a Gaussian random vector in $\mathcal{G}_{\mu,\sigma}$. \square

Corollary 7.2. Keep the assumptions of Theorem 7.2. Then $\kappa(Z_1(\mathbf{t})) = \max_{i,j=0}^{n-1} |g_i/g_j|$,

Proof. Combine the bounds of Theorems 7.1 and 7.2 on the spectral norms of the matrix $Z_1(\mathbf{t})$ and its inverse. \square

Theorem 7.3. (Cf. Table 8.8.) Assume $y \geq 0$ and a vector $\mathbf{t} \in \mathcal{G}_{\mu,\sigma}^{n \times 1}$. Then $F_{\|(Z_1(\mathbf{t}))^{-1}\|}(z/\sqrt{n}) = 1 - (1-p)^n$, for $p = \frac{1}{\sigma\sqrt{2\pi}} \int_{-1/z}^{1/z} \exp(-\frac{(x-\mu)^2}{2\sigma^2}) dx$, and so $F_{\|(Z_1(\mathbf{t}))^{-1}\|}(z/\sqrt{n}) \approx np$ where the value p is small. (This is the case if the value $|z|/\sqrt{n}$ is large or the distance between μ and the range $[-\sqrt{n}/z, \sqrt{n}/z]$ is large.)

Proof. By virtue of Theorem 7.2 $\|(Z_1(\mathbf{t}))^{-1}\| = 1/(\sqrt{n} \min_{i=0}^{n-1} |g_i|)$ for i.i.d. variables $g_i \in G_{\mu,\sigma}^{1 \times 1}$, $i = 0, \dots, n-1$. For any i , $i = 1, \dots, n$, it holds that $|g_i| \leq y$ with probability $1-p$ where $p = \frac{1}{\sigma\sqrt{2\pi}} \int_{-y}^y \exp(-\frac{(x-\mu)^2}{2\sigma^2}) dx$ (cf. Definition 6.1), and so $\min_{i=1}^n |g_i| \leq y$ with probability $(1-p)^n$ because g_i are independent random variables. Equivalently $1/(\sqrt{n} \min_{i=1}^n |g_i|) \geq 1/(y\sqrt{n})$ with probability $(1-p)^n$. Therefore $\|(Z_1(\mathbf{t}))^{-1}\| = 1/(\sqrt{n} \min_{i=1}^n |g_i|) \leq 1/(\sqrt{n} y)$ with probability $1 - (1-p)^n \geq np$. Substitute $z = 1/y$ and obtain the theorem. \square

Remark 7.1. By applying bounds (2.2) and (2.5) we can extend the theorem to estimate the cdfs $F_{\|Z_1(\mathbf{t})^{-1}\|_s}(z)$ where s can stand for $F, 1$, or ∞ .

7.4 Norm bounds for the inverse of a Gaussian random Toeplitz matrix

Our next subject is the estimates for the norm $\|T_n^{-1}\|_s$ for $s = 1, 2, \infty$ and a Gaussian random Toeplitz matrix $T_n \in \mathcal{T}_{\mu,0}^{n \times n}$, which is known to be nonsingular with probability 1. We can extend these estimates to the norm $\|T_n^{-1}\|_F$ by using (2.5).

Theorem 7.4. Given a matrix $T_n = (t_{i-j})_{i,j=1}^n \in \mathcal{T}_{0,1}^{n \times n}$, assumed to be nonsingular, write $\mathbf{p} = (p_i)_{i=1}^n = T_n^{-1} \mathbf{e}_1$, $\mathbf{q} = (q_i)_{i=1}^n = T_n^{-1} \mathbf{e}_n$, $g_n = p_n/|\mathbf{p}|$, $h_1 = q_1/|\mathbf{q}|$, and $p_1 = \mathbf{e}_n^T \mathbf{p} = \mathbf{e}_1^T \mathbf{q} = q_n$ (cf. (3.3)). Then $\|p_1 T_n^{-1}\|_s \leq 2n/(\alpha\beta)$ for $s = 1, 2, \infty$ and two random variables α and β such that

$$F_\alpha(y) \leq \sqrt{\frac{2n}{\pi}} \frac{y}{\sigma|g_n|} \text{ and } F_\beta(y) \leq \sqrt{\frac{2n}{\pi}} \frac{y}{\sigma|h_1|} \text{ for } y \geq 0. \quad (7.1)$$

Proof. By virtue of Theorem 5.1 we just need to estimate the two random variables $\|\mathbf{p}\|$ and $\|\mathbf{q}\|$. By virtue of its definition the vector \mathbf{p} is orthogonal to the vectors $T_n \mathbf{e}_2, \dots, T_n \mathbf{e}_n$, whereas $\mathbf{p}^T T_n \mathbf{e}_1 = 1$ (cf. [SST06]). Consequently the vectors $T_n \mathbf{e}_2, \dots, T_n \mathbf{e}_n$ uniquely define the vector $\mathbf{g} = (g_i)_{i=1}^n = \mathbf{p}/\|\mathbf{p}\|$, whereas $|\mathbf{g}^T T_n \mathbf{e}_1| = 1/\|\mathbf{p}\|$. The last coordinate t_{n-1} of the vector $T_n \mathbf{e}_1$ is independent of the vectors $T_n \mathbf{e}_2, \dots, T_n \mathbf{e}_n$ and consequently of the vector \mathbf{g} . Apply Remark 6.1 to estimate the cdf of the random variable $\alpha/|g_n| = 1/(\|\mathbf{p}\| |g_n|) = |\mathbf{g}^T T_n \mathbf{e}_1|/|g_n|$ and obtain that $F_\alpha(y) \leq \sqrt{\frac{2n}{\pi}} \frac{y}{\sigma|g_n|}$ for $y \geq 0$ (cf. (7.1)). Likewise we deduce that $F_\beta(y) \leq \sqrt{\frac{2n}{\pi}} \frac{y}{\sigma|h_1|}$ for $y \geq 0$ (cf. (7.1)). Combine these bounds with Theorem 5.1. \square

7.5 Bounding the leading entry of the inverse

Theorem 7.4 bounds the norm $\|T_n^{-1}\|$ in terms of the random variables $|u_n|$, $|v_1|$, and $|p_1| = |q_n|$. By applying parts (b) and (c) of Theorem 3.3 instead of its part (a), we similarly deduce the bounds $\|v_0 T_{n+1}^{-1}\| \leq 2/(\alpha\beta)$ and $\|v_n T_{n+1}^{-1}\| \leq 2/(\alpha\beta)$ for two pairs of random variables α and β that satisfy (7.1) for $n+1$ replacing n . Note that $p_1 = \frac{\det T_{n-1}}{\det T_n}$, $v_0 = \frac{\det T_n}{\det T_{n+1}}$, and $v_n = \frac{\det T_{0,1}}{\det T_{n+1}}$ for $T_{0,1} = (t_{i-j})_{i=0,j=1}^{n-1,n}$. Next we bound the geometric means of the ratios $|\frac{\det T_{h+1}}{\det T_h}|$ for $h = 1, \dots, k-1$. $1/|p_1|$ and $1/|v_0|$ are such ratios for $k = n-1$ and $k = n$, respectively, whereas the ratio $1/|v_n|$ is similar to $1/|v_0|$, under slightly distinct notation.

Theorem 7.5. *Let $T_h \neq O$ denote $h \times h$ matrices for $h = 1, \dots, k$ whose entries have absolute values at most t for a random variable t , e.g., for $t = \|T\|$. Furthermore let $T_1 = (t)$. Then the geometric mean $(\prod_{h=1}^{k-1} |\frac{\det T_{h+1}}{\det T_h}|)^{1/(k-1)} = \frac{1}{t} |\det T_k|^{1/(k-1)}$ is at most $k^{\frac{1}{2}(1+\frac{1}{k-1})} t$.*

Proof. The theorem follows from Hadamard's upper bound $|\det M| \leq k^{k/2} t^k$, which holds for any $k \times k$ matrix $M = (m_{i,j})_{i,j=1}^k$ with $\max_{i,j=1}^k |m_{i,j}| \leq t$. \square

The theorem shows that the geometric mean of the ratios $|\frac{\det T_{h+1}}{\det T_h}|$ for $h = 1, \dots, k-1$ is not greater than $k^{0.5+\epsilon(k)} t$ where $\epsilon(k) \rightarrow 0$ as $k \rightarrow \infty$. Furthermore if $T_n \in \mathcal{T}_{\mu,\sigma}^{n \times n}$ we can write t and recall Definition 6.1 to define the cdf of the Gaussian random variable t . This implies a reasonable lower bound on the expected value $|p_1| = |q_n|$.

7.6 The generic corner property, empirical results, and a link of Toeplitz inversion to polynomial computations

Our study in the two previous subsections implies that the norms of the inverses of Gaussian random Toeplitz matrices and consequently their condition numbers are expected to be reasonably bounded provided that the values $|g_n|$ and $|h_1|$ are not small. We call the latter provision the *generic property of two corners of the inverse*. Empirically standard Gaussian random Toeplitz matrices of reasonable sizes tend to be reasonably well conditioned. Our tests in the next section show this directly, whereas the numerical tests in [XXG12] and [PQZ13], provide additional indirect support. Namely the latter tests succeeded by employing random Toeplitz multipliers, whereas they would have been expected to fail numerically if the multipliers were ill conditioned. Thus the generic property tends to hold empirically in the case of standard Gaussian random Toeplitz matrices. Formal support of these empirical data should be specialized to exclude the case where the absolute value of the mean value μ greatly exceeds the standard deviation σ . Indeed in this case the norm $\|T_n^{-1}\|$ is likely to be large (see Section 7.2), which contradicts to the generic property by virtue of Theorems 7.4 and 7.5.

We conclude this section by recalling a link of Toeplitz matrix inversion to some polynomial computations. Namely it is well known (see [P01, Section 2.11]) that the equation $T_n \mathbf{p} = \mathbf{e}_1$ is equivalent to the polynomial equation $t(x)p(x) \bmod x^{2n+1} = r(x)$ where $r(x)$ is a monic polynomial of degree n , whereas $p(x)$ and $t(x)$ are two polynomials of degrees n and $2n-1$, respectively, with the coefficient vectors \mathbf{p} and \mathbf{t}_+ , respectively. For a given vector \mathbf{t}_+ , the Euclidean algorithm computes the coefficients of the polynomials $p(x)$ and $r(x)$ (apart from the cases of degeneracy, occurring with probability 0 for Gaussian random input) but provides no explicit expressions for these coefficients.

8 Numerical Experiments

Our numerical experiments with random general, Toeplitz and circulant matrices have been performed in the Graduate Center of the City University of New York on a Dell server with a dual core 1.86 GHz Xeon processor and 2GB memory running Windows Server 2003 R2. The tests were implemented with MATLAB R2012b (for Table 8.1 through Table 8.5) and Fortran (for Table 8.6 through Table 8.8). Fortran code was compiled with the GNU gfortran compiler within the Cygwin environment. In MATLAB, Gaussian (resp. uniform) random numbers were generated with the intrinsic function *randn()* (resp. *rand()*), while in Fortran random numbers were generated with the *random_number()* intrinsic Fortran function, assuming the uniform probability distribution. The tests were designed by the first author and performed by his coauthors.

We have computed the condition numbers of standard Gaussian random $n \times n$ matrices for $n = 2^k$, $k = 5, 6, \dots$, as well as complex general, Toeplitz, and circulant matrices whose i.i.d. entries had standard Gaussian real and imaginary random parts. We performed 1000 tests for each class of inputs, each dimension n , and each nullity r . Tables 8.2–8.4 display the test results. The last four columns of each table display the average (mean), minimum, maximum, and standard deviation of the computed condition numbers of the input matrices, respectively. Namely we computed the values $\kappa(A) = \|A\| \|A^{-1}\|$ for general, Toeplitz, and circulant matrices A . The estimates in Section 7.3 are sharp, and so we were most interested in testing the condition numbers of random Toeplitz matrices. The displayed data illustrates that these condition numbers grow substantially faster with the growth of the matrix size than in the case of random circulant matrices but not faster than in the case of random general matrices, expected to be well-conditioned according to the results of the intensive formal and experimental studies in [D88], [E88], [ES05], [CD05], [SST06]. For comparison we also performed similar tests using random variables generated under the uniform probability distribution in the range $[-1, 1]$ and displayed the test results in Tables 8.5–8.8. In all cases the tests showed similar behavior of the norms and condition numbers of general, Toeplitz and circulant matrices generated under the uniform and Gaussian probability distributions.

Table 8.1: The norms of standard Gaussian random general, Toeplitz and circulant $n \times n$ matrices A and their inverses

matrix A	n	$\ A\ _1$	$\ A\ _2$	$\frac{\ A\ _1}{\ A\ _2}$	$\ A^{-1}\ _1$	$\ A^{-1}\ _2$	$\frac{\ A^{-1}\ _1}{\ A^{-1}\ _2}$
General	32	4.83×10^1	1.51×10^1	3.20	1.29×10^1	7.12×10^0	1.81
General	64	9.28×10^1	2.18×10^1	4.25	1.88×10^1	9.61×10^0	1.96
General	128	1.80×10^2	3.13×10^1	5.76	3.36×10^1	1.56×10^1	2.15
General	256	3.51×10^2	4.47×10^1	7.85	5.67×10^1	2.51×10^1	2.26
General	512	6.87×10^2	6.34×10^1	10.8	5.81×10^1	2.46×10^1	2.36
General	1024	1.35×10^3	9.00×10^1	15.0	7.60×10^1	3.08×10^1	2.46
Toeplitz	32	4.38×10^1	1.57×10^1	2.79	5.89×10^0	3.20×10^0	1.84
Toeplitz	64	8.58×10^1	2.32×10^1	3.69	8.18×10^0	4.15×10^0	1.97
Toeplitz	128	1.68×10^2	3.57×10^1	4.72	9.10×10^0	4.29×10^0	2.12
Toeplitz	256	3.32×10^2	5.22×10^1	6.36	1.53×10^1	6.68×10^0	2.30
Toeplitz	512	6.58×10^2	7.75×10^1	8.48	1.31×10^1	5.16×10^0	2.54
Toeplitz	1024	1.30×10^3	1.15×10^2	11.3	2.04×10^1	7.62×10^0	2.29
Circulant	32	4.04×10^1	1.62×10^1	2.50	1.40×10^0	8.97×10^{-1}	1.56
Circulant	64	8.04×10^1	2.24×10^1	3.29	1.52×10^0	9.55×10^{-1}	1.59
Circulant	128	1.61×10^2	3.64×10^1	4.42	1.79×10^0	1.14×10^0	1.57
Circulant	256	3.21×10^2	5.71×10^1	5.62	1.88×10^0	1.14×10^0	1.65
Circulant	512	6.44×10^2	8.21×10^1	7.84	2.27×10^0	1.45×10^0	1.57
Circulant	1024	1.28×10^3	1.24×10^2	10.4	2.03×10^0	1.17×10^0	1.74

Table 8.2: The condition numbers of standard Gaussian random real $n \times n$ matrices

n	min	max	mean	std
32	2.43×10^1	3.71×10^5	3.93×10^2	1.75×10^3
64	4.14×10^1	3.10×10^6	1.46×10^3	1.21×10^4
128	8.15×10^1	7.48×10^5	1.63×10^3	4.93×10^3
256	1.71×10^2	3.37×10^7	8.71×10^3	1.12×10^5
512	4.04×10^2	3.40×10^7	1.54×10^4	1.52×10^5
1024	8.10×10^2	4.44×10^7	1.86×10^4	1.49×10^5
2048	1.53×10^3	6.92×10^7	3.01×10^4	1.96×10^5

Table 8.3: The condition numbers of standard Gaussian random Toeplitz real $n \times n$ matrices

n	min	max	mean	std
32	6.24×10^0	1.27×10^4	1.44×10^2	6.51×10^2
64	1.04×10^1	5.81×10^4	2.70×10^2	1.93×10^3
128	2.41×10^1	1.29×10^5	6.05×10^2	4.54×10^3
256	3.41×10^1	1.25×10^5	8.00×10^2	4.77×10^3
512	6.42×10^1	6.33×10^5	2.09×10^3	2.35×10^4
1024	1.11×10^2	2.44×10^5	6.72×10^3	8.32×10^4
2048	3.00×10^2	3.51×10^5	9.61×10^3	1.83×10^4
4096	4.81×10^2	2.54×10^5	7.44×10^3	2.92×10^4

Table 8.4: The condition numbers of standard Gaussian random circulant $n \times n$ matrices

n	min	max	mean	std
32	4.76×10^0	6.29×10^1	1.53×10^1	9.39×10^0
64	9.17×10^0	3.67×10^2	3.11×10^1	3.81×10^1
128	1.20×10^1	3.88×10^2	4.23×10^1	4.66×10^1
256	1.61×10^1	3.89×10^2	6.65×10^1	5.63×10^1
512	2.56×10^1	5.05×10^2	1.05×10^2	8.72×10^1
1024	3.64×10^1	2.33×10^3	1.70×10^2	2.51×10^2
2048	5.86×10^1	9.52×10^2	1.93×10^2	1.39×10^2
4096	8.17×10^2	3.92×10^3	3.92×10^2	5.34×10^2
8192	1.32×10^2	2.38×10^3	4.39×10^2	3.28×10^2
16384	2.08×10^2	1.61×10^4	9.73×10^2	1.98×10^2
32768	1.97×10^2	8.26×10^3	8.75×10^2	9.47×10^2
65536	3.52×10^2	5.91×10^3	1.37×10^3	9.95×10^2
131072	5.79×10^2	1.35×10^4	2.00×10^3	1.98×10^3
262144	8.26×10^2	1.10×10^4	2.57×10^3	1.79×10^3
524288	1.03×10^3	3.91×10^4	4.34×10^3	4.68×10^3
1048576	1.54×10^3	6.56×10^4	8.22×10^3	9.93×10^3

Table 8.5: The norms of uniformly random general, Toeplitz and circulant $n \times n$ matrices A and their inverses

matrix	n	$\ A\ _1$	$\ A\ _2$	$\frac{\ A\ _1}{\ A\ _2}$	$\ A^{-1}\ _1$	$\ A^{-1}\ _2$	$\frac{\ A^{-1}\ _1}{\ A^{-1}\ _2}$
General	32	2.76×10^1	8.61×10^0	3.21	2.23×10^1	1.26×10^1	1.77
General	64	5.41×10^1	1.25×10^1	4.32	3.13×10^1	1.61×10^1	1.94
General	128	1.06×10^2	1.80×10^1	5.89	4.58×10^1	2.23×10^1	2.06
General	256	2.09×10^2	2.57×10^1	8.10	7.61×10^1	3.44×10^1	2.21
General	512	4.11×10^2	3.66×10^1	11.2	1.01×10^2	4.39×10^1	2.30
General	1024	8.14×10^2	5.19×10^1	15.7	1.57×10^2	6.39×10^1	2.45
Toeplitz	32	2.60×10^1	8.90×10^0	2.92	8.29×10^0	4.50×10^0	1.84
Toeplitz	64	5.14×10^1	1.38×10^1	3.73	1.19×10^1	6.03×10^0	1.97
Toeplitz	128	1.01×10^2	2.06×10^1	4.91	1.56×10^1	7.23×10^0	2.15
Toeplitz	256	2.01×10^2	3.01×10^1	6.66	1.99×10^1	8.71×10^0	2.29
Toeplitz	512	3.99×10^2	4.45×10^1	8.96	2.43×10^1	9.46×10^0	2.57
Toeplitz	1024	7.95×10^2	6.61×10^1	12.0	3.45×10^1	1.24×10^1	2.79
Circulant	32	2.44×10^1	9.19×10^0	2.66	3.50×10^0	2.64×10^0	1.32
Circulant	64	4.87×10^1	1.41×10^1	3.45	3.36×10^0	2.38×10^0	1.41
Circulant	128	9.74×10^1	2.12×10^1	4.60	3.54×10^0	2.45×10^0	1.45
Circulant	256	1.96×10^2	3.22×10^1	6.08	3.11×10^0	1.83×10^0	1.69
Circulant	512	3.91×10^2	4.74×10^1	8.26	3.36×10^0	2.01×10^0	1.67
Circulant	1024	7.85×10^2	7.10×10^1	11.1	3.81×10^0	2.29×10^0	1.66

Table 8.6: The condition numbers of uniformly random real $n \times n$ matrices

n	min	max	mean	std
32	2.4×10^1	1.8×10^3	2.4×10^2	3.3×10^2
64	4.6×10^1	1.1×10^4	5.0×10^2	1.1×10^3
128	1.0×10^2	2.7×10^4	1.1×10^3	3.0×10^3
256	2.4×10^2	8.4×10^4	3.7×10^3	9.7×10^3
512	3.9×10^2	7.4×10^5	1.8×10^4	8.5×10^4
1024	8.8×10^2	2.3×10^5	8.8×10^3	2.4×10^4
2048	2.1×10^3	2.0×10^5	1.8×10^4	3.2×10^4

Table 8.7: The condition numbers of uniformly random real Toeplitz $n \times n$ matrices

n	min	mean	max	std
256	9.1×10^2	9.2×10^3	1.3×10^5	1.8×10^4
512	2.3×10^3	3.0×10^4	2.4×10^5	4.9×10^4
1024	5.6×10^3	7.0×10^4	1.8×10^6	2.0×10^5
2048	1.7×10^4	1.8×10^5	4.2×10^6	5.4×10^5
4096	4.3×10^4	2.7×10^5	1.9×10^6	3.4×10^5
8192	8.8×10^4	1.2×10^6	1.3×10^7	2.2×10^6

Table 8.8: The condition numbers of uniformly random circulant $n \times n$ matrices

n	min	mean	max	std
256	9.6×10^0	1.1×10^2	3.5×10^3	4.0×10^2
512	1.4×10^1	8.5×10^1	1.1×10^3	1.3×10^2
1024	1.9×10^1	1.0×10^2	5.9×10^2	8.6×10^1
2048	4.2×10^1	1.4×10^2	5.7×10^2	1.0×10^2
4096	6.0×10^1	2.6×10^2	3.5×10^3	4.2×10^2
8192	9.5×10^1	3.0×10^2	1.5×10^3	2.5×10^2
16384	1.2×10^2	4.2×10^2	3.6×10^3	4.5×10^2
32768	2.3×10^2	7.5×10^2	5.6×10^3	7.1×10^2
65536	2.4×10^2	1.0×10^3	1.2×10^4	1.3×10^3
131072	3.9×10^2	1.4×10^3	5.5×10^3	9.0×10^2
262144	6.3×10^2	3.7×10^3	1.1×10^5	1.1×10^4
524288	8.0×10^2	3.2×10^3	3.1×10^4	3.7×10^3
1048576	1.2×10^3	4.8×10^3	3.1×10^4	5.1×10^3

Appendix

A Randomness and nonsingularity

The total degree of a multivariate monomial is the sum of its degrees in all its variables. The total degree of a polynomial is the maximal total degree of its monomials.

Lemma A.1. [DL78], [S80], [Z79]. For a set Δ of a cardinality $|\Delta|$ in any fixed ring let a polynomial in m variables have a total degree d and let it not vanish identically on this set. Then the polynomial vanishes in at most $d|\Delta|^{m-1}$ points.

We assume that Gaussian random variables range over infinite sets Δ , usually over the real line or its interval. Then the lemma implies that a nonzero polynomial vanishes with probability 0. Consequently a square Gaussian general, Toeplitz or circulant random matrix is nonsingular with probability 1 because its determinant is a polynomial in the entries. Likewise rectangular Gaussian general, Toeplitz and circulant random matrices have full rank with probability 1, and similarly under the other probability distributions whose measures are absolutely continuous relatively to Lebesgue's measure. These results can be also adapted to the case of the probability distribution over finite sets [DL78], [S80], [Z79].

Acknowledgements: Our research has been supported by NSF Grant CCF-1116736 and PSC CUNY Awards 64512-0042 and 65792-0043. We are grateful to the reviewer for helpful comments.

References

- [BG05] A. Böttcher, S. M. Grudsky, *Spectral Properties of Banded Toeplitz Matrices*, SIAM Publications, Philadelphia, 2005.
- [CD05] Z. Chen, J. J. Dongarra, Condition Numbers of Gaussian Random Matrices, *SIAM J. on Matrix Analysis and Applications*, **27**, 603–620, 2005.
- [CPW74] R. E. Cline, R. J. Plemmons, and G. Worm, Generalized Inverses of Certain Toeplitz Matrices, *Linear Algebra and Its Applications*, **8**, 25–33, 1974.
- [D88] J. Demmel, The Probability That a Numerical Analysis Problem Is Difficult, *Math. of Computation*, **50**, 449–480, 1988.
- [DL78] R. A. Demillo, R. J. Lipton, A Probabilistic Remark on Algebraic Program Testing, *Information Processing Letters*, **7**, **4**, 193–195, 1978.
- [E88] A. Edelman, Eigenvalues and Condition Numbers of Random Matrices, *SIAM J. on Matrix Analysis and Applications*, **9**, **4**, 543–560, 1988.
- [ES05] A. Edelman, B. D. Sutton, Tails of Condition Number Distributions, *SIAM J. on Matrix Analysis and Applications*, **27**, **2**, 547–560, 2005.
- [GK72] I. Gohberg, N. Y. Krupnick, A Formula for the Inversion of Finite Toeplitz Matrices, *Matematicheskie Issledovaniia* (in Russian), **7**, **2**, 272–283, 1972.
- [GL96] G. H. Golub, C. F. Van Loan, *Matrix Computations*, Johns Hopkins University Press, Baltimore, Maryland, 1996 (third addition).
- [GS72] I. Gohberg, A. Sementsul, On the Inversion of Finite Toeplitz Matrices and Their Continuous Analogs, *Matematicheskie Issledovaniia* (in Russian), **7**, **2**, 187–224, 1972.
- [HMT11] N. Halko, P. G. Martinsson, J. A. Tropp, Finding Structure with Randomness: Probabilistic Algorithms for Constructing Approximate Matrix Decompositions, *SIAM Review*, **53**, **2**, 217–288, 2011.

- [P01] V. Y. Pan, *Structured Matrices and Polynomials: Unified Superfast Algorithms*, Birkhäuser/Springer, Boston/New York, 2001.
- [PQYa] V. Y. Pan, G. Qian, A. Zheng, Supporting GENP and Low-rank Approximation with Random Multipliers, Tech. Report TR 2014008, *PhD Program in Comp. Sci., Graduate Center, CUNY*, 2014,
available at <http://www.cs.gc.cuny.edu/tr/techreport.php?id=472>
- [PQZ13] V. Y. Pan, G. Qian, A. Zheng, Randomized Preprocessing versus Pivoting, *Linear Algebra and Its Applications*, **438**, 4, 1883–1899, 2013.
- [S80] J. T. Schwartz, Fast Probabilistic Algorithms for Verification of Polynomial Identities, *Journal of ACM*, **27**, 4, 701–717, 1980.
- [S98] G. W. Stewart, *Matrix Algorithms, Vol I: Basic Decompositions*, SIAM, 1998.
- [SST06] A. Sankar, D. Spielman, S.-H. Teng, Smoothed Analysis of the Condition Numbers and Growth Factors of Matrices, *SIAM J. on Matrix Analysis*, **28**, 2, 446–476, 2006.
- [XXG12] J. Xia, Y. Xi, M. Gu, A Superfast Structured Solver for Toeplitz Linear Systems via Randomized Sampling, *SIAM J. Matrix Anal. Appl.*, **33**, 837–858, 2012.
- [Z79] R. E. Zippel, Probabilistic Algorithms for Sparse Polynomials, *Proceedings of EU-ROSAM'79, Lecture Notes in Computer Science*, **72**, 216–226, Springer, Berlin, 1979.