

3-16-1999

# Novel Carbon-Carbon Bond Oxidative Cleavage of Hexabenzyl-hexaazaisowurtzitane by n-BuONO and $(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$

Xiao-Pei Guan  
*CUNY City College*

Hong Yan  
*Beijing Polytechnic University*

Jian-Guang Sun  
*CUNY City College*

Yong-Zhong Yu

## How does access to this work benefit you? Let us know!

Follow this and additional works at: [https://academicworks.cuny.edu/cc\\_pubs](https://academicworks.cuny.edu/cc_pubs)

 Part of the [Chemistry Commons](#), [Civil and Environmental Engineering Commons](#), and the [Environmental Sciences Commons](#)

---

### Recommended Citation

Guan, Xiao-Pei; Yan, Hong; Sun, Jian-Guang; and Yu, Yong-Zhong, "Novel Carbon-Carbon Bond Oxidative Cleavage of Hexabenzyl-hexaazaisowurtzitane by n-BuONO and  $(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$ " (1999). *CUNY Academic Works*.  
[https://academicworks.cuny.edu/cc\\_pubs/481](https://academicworks.cuny.edu/cc_pubs/481)

This Article is brought to you for free and open access by the City College of New York at CUNY Academic Works. It has been accepted for inclusion in Publications and Research by an authorized administrator of CUNY Academic Works. For more information, please contact [AcademicWorks@cuny.edu](mailto:AcademicWorks@cuny.edu).

## Novel Carbon-Carbon Bond Oxidative Cleavage of Hexabenzyl-hexaazaisowurtzitane by *n*-BuONO and $(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$

Xiao-Pei Guan<sup>1,\*</sup> Hong Yan<sup>2</sup>, Jian-Guang Sun<sup>1</sup> and Yong-Zhong Yu

School of Chemical Engineering and Materials Science, Beijing Institute of Technology, 100081 Beijing, P. R. China

<sup>1</sup>Current address: Department of Chemistry, The City College of CUNY, Convent Avenue at 138th Street, New York, NY 10031, USA, e-mail: [xpguan@hotmail.com](mailto:xpguan@hotmail.com)

<sup>2</sup>Current address: Department of Chemistry and Environmental Engineering, Beijing Polytechnic University, Beijing 100022, P. R. China, e-mail: [hongyan@bj.col.com.cn](mailto:hongyan@bj.col.com.cn)

\*Author to whom correspondence should be addressed.

Received: 12 May 1998 / Accepted: 5 March 1999 / Published: 16 March 1999

---

**Abstract:** Treatment of 2,4,6,8,10,12-hexabenzyl-2,4,6,8,10,12-hexaazatetracyclo[5.5.0.0<sup>5,9</sup>.0<sup>3,11</sup>]dodecane (**1**) with *n*-butyl nitrite or ammonium cerium nitrate results in novel C(1)-C(7) oxidative cleavage to give 1,3,4,5,7,8-hexabenzyl-*cis-cisoid-cis*-hexahydro-1*H*,5*H*-diimidazolium[4,5-*b*:4',5'-*e*]pyrazine dinitrate (**2**).

**Keywords:** Oxidative cleavage, hexabenzylhexaazaisowurtzitane, iminium cation.

---

### Introduction


2,4,6,8,10,12-Hexabenzyl-2,4,6,8,10,12-hexaazatetracyclo[5.5.0.0<sup>5,9</sup>.0<sup>3,11</sup>]dodecane (hexabenzyl-hexaazaisowurtzitane, **1**) is an interesting polyaza “cage” compound synthesized recently [1]. In an effort to study the chemical behaviour of **1**, we have explored a series of debenylation method to remove the *N*-benzyl groups in **1** [2]. Reductive acetyldebenzylation of **1** ( $\text{H}_2$ , 20%Pd(OH)<sub>2</sub>/C, Ac<sub>2</sub>O) gives 2,6,8,12-tetraacetyl-4,10-dibenzyl-2,4,6,8,10,12-hexaazatetracyclo[5.5.0.0<sup>5,9</sup>.0<sup>3,11</sup>]dodecane[2,3], which could be further transformed to hexanitrohexaazaisowurtzitane [2]. In this communication we

wish to report an unexpected carbon-carbon oxidative cleavage of **1** caused by *n*-BuONO and  $(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$ , respectively (Scheme 1).

## Results and Discussion

*n*-BuONO is a convenient reagent for *N*-dealkylnitrosation of *N,N*-dialkyl aromatic amines with preferential debenylation [4]. However, under the same condition, compound **1** did not go through *N*-dealkylnitrosation. Instead, diimidazolium dinitrate **2** was obtained in 42% yield, whose structure was established through the analyses of IR,  $^1\text{H}$  and  $^{13}\text{C}$  NMR, MS and elemental microanalysis. In particular, the IR spectrum shows a strong peak at  $1640\text{ cm}^{-1}$ , indicating the existence of iminium cation, which was further confirmed by  $^1\text{H}$  NMR spectrum ( $\delta$  9.04 ppm, 2H) and  $^{13}\text{C}$  NMR spectrum ( $\delta$  158.4 ppm, d), and a very intense band of nitrate anion at  $1382\text{ cm}^{-1}$ . The elemental analysis and the field desorption mass spectrum ( $m/z$  770,  $\text{M}-\text{NO}_3^-$ , 100%), is consistent with formula  $\text{C}_{48}\text{H}_{48}\text{N}_8\text{O}_6$ . In addition, the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra also indicated that the molecule was mirror symmetric. These results demonstrate that the product **2** is 1,3,4,5,7,8-hexabenzyl-*cis-cisoid-cis*-hexahydro-1*H*,5*H*-diimidazolium[4,5-*b*:4',5'-*e*]pyrazine.

Under mild conditions,  $(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$  can remove the *N*-(4-methoxybenzyl) group on 2,5-piperazinediones [5]. However, when **1** was exposed to  $(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$ , instead of debenylation, **2** was isolated in 59% yield.




**Scheme 1.**

The easy cleavage of *C*(1)-*C*(7) bond reflects that the cage skeleton of **1** is somewhat strained. Crystal structure analysis of hexa(4'-chlorobenzyl)hexaazaisowurtzitane shows that the framework *C*-*C* bonds are longer than usual value for unstrained  $C_{sp^3}$ - $C_{sp^3}$  bond and *C*(1)-*C*(7) bond [ $1.579(3)\text{Å}$ ] is the longest [1b]. This implies that *C*(1)-*C*(7) bond is probably the weakest *C*-*C* bond in this cage skeleton.

A mechanism that accounts for the observed oxidative cleavage of **1** to **2** is depicted in Scheme 2. The initially formed aminium radical cation **3** rapidly undergoes *C*(1)-*C*(7) cleavage followed by a second electron loss to give diimidazolium dinitrate **2**. It has been reported that 1,2-diamine radical cations undergo rapid *C*-*C* bond fragmentation to form an  $\alpha$ -amino radical and an iminium ion [6], and iminium salts are generally believed to be the key intermediates during the oxidation of tertiary amines

[7]. The readily oxidative cleavage of **1** must arise from the through-bond interaction between  $N(2)$ ,  $N(6)$ ,  $N(8)$  and  $N(12)$  atoms, which stabilizes the intermediate-like transition state relative to the ground state. Relief of strain energy contained within the hexaazaisowurtzitane framework and formation of the resonance-stabilized imidazolium ion provides the driving force for the oxidative cleavage of  $C(1)$ - $C(7)$  bond. The similar type of Grob fragmentation was also observed in the reaction of triethylenediamine with either  $\text{ClO}_2$  or  $\text{HOCl}$  [8].



**Scheme 2.**

In oxidative dealkylation of tertiary amine, the intermediate iminium cation, which are generally believed to be the key intermediates, is hydrolyzed more or less rapidly, in aqueous solution, with the formation of the corresponding secondary amine [7]. Usually, imidazolium salts are also too labile to resist hydrolysis [9]. The high stability of **2** towards hydrolysis may be partially attributed to being capped of the cage skeleton by the six sterically hindered and hydrophobic benzyl groups.

## Experimental

### Oxidation of Hexaazahexbenzylisowurtzitane (**1**) with *n*-BuONO

A mixture of **1** (501 mg, 0.707 mmol), *n*-BuONO (5.0 mL, 55 mmol),  $\text{NH}_4\text{Cl}$  (50 mg, 0.99 mmol) and  $\text{H}_2\text{O}$  (150  $\mu\text{L}$ , 8.3 mmol) was refluxed under  $\text{N}_2$  for 1h. Volatile material was then evaporated in vacua and the residue was washed with a small amount of water. Recrystallization of the crude solid from  $\text{CH}_3\text{CN}/\text{CHCl}_3/\text{hexane}$  afforded pure **2** as colorless crystals (249 mg, 42.3%), mp 224.2~225.0  $^\circ\text{C}$ ; IR (KBr,  $\text{cm}^{-1}$ ): 1640( $\text{C}=\text{N}^+$ ), 1382( $\text{NO}_3^-$ ), 692;  $^1\text{H}$  NMR (DMSO- $d_6$ , 60MHz):  $\delta$  3.93(s, 4H,  $\text{CH}_2$ ), 4.59(d, 4H,  $J_{\text{AB}}=15.0\text{Hz}$ ,  $\text{CH}_2$ ), 4.99(d, 4H,  $J_{\text{AB}}=15.0\text{Hz}$ ,  $\text{CH}_2$ ), 5.33(s, 4H, CH), 6.9~7.6 (m, 30H, Ph), 9.04(s, 2H,  $\text{HC}=\text{N}^+$ ) ;  $^{13}\text{C}$  NMR (DMSO- $d_6$ , 75Hz):  $\delta$  48.2(t), 58.4(t), 68.9(d), 127.9(d), 128.0(d),

128.3(d), 128.5(d), 128.7(d), 129.0(d), 133.4(s), 135.9(s), 158.4(d); MS (FD, m/z): 770(M<sup>+</sup>-NO<sub>3</sub>, 100%). Anal. Calcd. for C<sub>48</sub>H<sub>48</sub>N<sub>8</sub>O<sub>6</sub>: C, 69.21; H, 5.81; N, 13.45; O, 11.52; Found: C, 69.13; H, 6.00; N, 13.67; O, 11.37.

*Oxidation of Hexaazahexbenzylisowurtzitane (1) with (NH<sub>4</sub>)<sub>2</sub>Ce(NO<sub>3</sub>)<sub>6</sub>*

A solution of **1** (710 mg, 1.00 mmol) and (NH<sub>4</sub>)<sub>2</sub>Ce(NO<sub>3</sub>)<sub>6</sub> (12.995 g, 23.70 mmol) in acetonitrile (75 mL) and water (25 mL) was stirred at r.t. for 24h. EtOAc (50 mL) was then added. The organic layer was separated, washed with water, dried over MgSO<sub>4</sub> and evaporated in vacua. The residual solid was recrystallized from CH<sub>3</sub>CN/CHCl<sub>3</sub>/ hexane to give pure **2** as colorless crystals (490 mg, 58.8%).

## References and Notes

1. (a) Nielsen, A.T.; Nissan, R.A.; Vanderah, D.J.; Coon, C.L.; Gilardi, R.D.; George, C.F.; Flippen-Anderson, J. *J. Org. Chem.* **1990**, *55*, 1459; (b) Crampton, M.R.; Hamid, J.; Millar, R.; Ferguson, G. *J. Chem. Soc., Perkin Trans. 2*, **1993**, 923.
2. Guan, X.-P. *Ph. D. Dissertation*, Beijing Institute of Technology, **1993**.
3. Bellamy, A.J., *Tetrahedron* **1995**, *51*, 4711.
4. Verando, G; Giumanini, A.G.; Strazzolini, P. *Tetrahedron* **1991**, *47*, 7845.
5. Yamaura, M.; Suzuki, T.; Hashimoto, H.; Yoshimura, J.; Okamoto, T.; Shin, C.-G. *Bull. Chem. Soc. Jpn.* **1985**, *58*, 1413.
6. (a) Leon, J.W.; Whitten, D.G. *J. Am. Chem. Soc.* **1993**, *115*, 8038; (b) Wang, Y.; Lucia, L.A.; Schanze, K.S. *J. Phys. Chem.* **1995**, *99*, 1961.
7. Rosenblatt, D.H.; Burrows, E.P. In "The Chemistry of Amino, Nitro and Nitroso Compounds and Their Derivatives". Patai, S. Ed.; Wiley: New York. **1982**, *Supplement F*, pp1085-1149.
8. Dennis, W.H., Jr.; Hull, L.A.; Rosenblatt, D.H. *J. Org. Chem.* **1967**, *32*, 3783.
9. (a) May M.; Bardos, T.J.; Barger, F.L.; Lansford, M.; Ravel, J.M.; Sutherland, G.L.; William, S. *J. Am. Chem. Soc.* **1951**, *73*, 3067; (b) Jaenicke, L.; Brode, E. *Liebigs Ann. Chem.* **1959**, *624*, 120; (c) Hafferl, W.; Lundin, R.; Ingraham, L. L. *Biochemistry* **1963**, *2*, 1298.

*Samples Availability:* Available from the authors.