

City University of New York (CUNY)

CUNY Academic Works

Publications and Research

Queens College

2021

First Report of *Mummenhoffia alliacea* (Brassicaceae) for New York

Daniel Atha

New York Botanical Garden

Eve Levine

John F. Gaskin

USDA, Agricultural Research Service

César Castillo

CUNY Queens College

[How does access to this work benefit you? Let us know!](#)

More information about this work at: https://academicworks.cuny.edu/qc_pubs/494

Discover additional works at: <https://academicworks.cuny.edu>

This work is made publicly available by the City University of New York (CUNY).

Contact: AcademicWorks@cuny.edu

**FIRST REPORT OF MUMMENHOFFIA ALLIACEA (BRASSICACEAE)
FOR NEW YORK**

DANIEL ATHA

Center for Conservation Strategy
New York Botanical Garden
Bronx, New York 10458
datha@nybg.org

EVE LEVINE

212 W 17th St, #3b
New York, New York 10011
elevine800@gmail.com

JOHN F. GASKIN

USDA Agricultural Research Service
1500 North Central Avenue
Sidney, Montana 59270
john.gaskin@usda.gov

CÉSAR CASTILLO

Queens College
City University of New York
65-30 Kissena Blvd
Flushing, New York 11367
cesar.castillo@qc.cuny.edu

ABSTRACT

Mummenhoffia alliacea is naturalized in New York. The first documented occurrences are reported from Dutchess, Kings, New York (Manhattan), Onondaga, Queens, and Suffolk counties. Verification is provided by herbarium specimens at NY and photographs uploaded to iNaturalist and cited at the Global Biodiversity Information Facility.

Spontaneous occurrences of *Mummenhoffia alliacea* (L.) Esmailbegi & Al-Shehbaz (Garlic Pennycress) were found in 2017, 2018, 2019, 2020 and 2021 during routine observation of the flora of New York State. Observations were uploaded to iNaturalist and identifications confirmed by the third author (JG). These are the first reports of this species (and genus) growing spontaneously in New York state (Al-Shehbaz 2009; Werier 2017; USDA, NRCS 2021).

Mummenhoffia alliacea was transferred from *Thlaspi* to the recently named genus *Mummenhoffia* Esmailbegi & Al-Shehbaz (Esmailbegi et al. 2018). The species is native to central and southern Europe and was first reported as introduced to North America in Pennsylvania in 1947 (Cusick 2015). Plants are annual and in North America are typically found in disturbed areas such as agricultural fields, gardens and highway edges.

Vouchers. New York. Queens Co.: New York City, I-495 and Cross Island Parkway, east-bound, raised median, 21 Apr 2021, *Atha 16190* (NY, to be distributed to BH, F, MO, MU, TEX, US). **New York Co.:** New York City, Riverside Park South between 65th and 66th Sts, ca 500 m E of the Hudson River, 26 Apr 2021, *Atha 16195* (NY, to be distributed to BH, F, MO, MU, TEX, US).

Additional populations in New York State are documented in iNaturalist as follows (iNaturalist observation numbers). **Dutchess Co.:** 65451510. **Kings Co.:** 74673073. **New York Co.:** 73145620, 73145621, 73145622, 73145624, 73145626, 73145627, 73145628, 73145630, 73145633, 73235274,

73235294, 73235303, 73235321, 73235344, 73235381, 73354025, 73354891, 73355589, 73358048, 73734913, 73986078, 73986091, 73986137, 73986167, 73986191, 73986211, 73986258, 74013946, 74014591. Onondaga Co.: 22711360, 22816069. Queens Co.: 11278343, 21744939, 23772101, 24973369, 74730008, 74730009. Suffolk Co.: 73045649. Voucher specimens and iNaturalist observations indexed by the Global Biodiversity Information Facility (GBIF 2021).



Figure 1. *Mummenhoffia alliacea*. A. Whole plant (Queens Co., *Atha* 16190, NY and iNaturalist observation 74730008), ruler is 30 cm long. B. Cauline leaves with claspings, auriculate bases (46771830). C. Glaucous stem with spreading hairs at the base (74014591). D. Fruits (73145633), smallest increments are millimeters. E. Population (74730009).

Vegetatively and in flower (before fruiting) *Mummenhoffia alliacea* is similar in appearance to *Capsella bursa-pastoris*, *Lepidium campestre*, *Lepidium virginicum*, *Microthlaspi perfoliatum*, and especially *Thlaspi arvense*. Distinguishing these species when mature is easier: *Capsella bursa-pastoris* and *Microthlaspi perfoliatum* have fruit that are notched at the tip (vs. blunt in *Mummenhoffia alliacea*), and the *Lepidium* species mentioned here have flattened fruit (vs somewhat inflated). Finally, *Mummenhoffia alliacea* can be distinguished from *Thlaspi arvense* by the following key.

Key to distinguish *Mummenhoffia alliacea* from *Thlaspi arvense*

1. Bruised plant with garlic-like odor; stems terete, glaucous, pubescent basally with spreading hairs; upper stem leaves clasping with auricles wrapping around the stem; fruit scarcely winged, the wings < 1mm wide; seeds alveolar..... **Mummenhoffia alliacea**
1. Bruised plant without garlic odor; stems ribbed and sometimes winged, green, glabrous basally; upper stem leaves clasping but auricles not wrapping around the stem; fruits broadly winged, the wings 3.5–5.0 mm wide; seeds concentrically striate..... **Thlaspi arvense**

Al-Shehbaz (2009), in his treatment of the Brassicaceae for the Flora of North America, reported this species (as *Thlaspi alliaceum*) from Delaware, Indiana, Kentucky, Louisiana, Maryland, North Carolina, Ohio, Pennsylvania, Tennessee, Virginia, and West Virginia. The USDA Plants Database also reports the species from Georgia, Missouri, and New Jersey (USDA-NCRS 2021). Verified observations in iNaturalist are reported from Massachusetts (iNaturalist observation number 41052744).

The species was first reported for New Jersey in 2006 (Lamont & Young 2006) and will undoubtedly be found in many more New York counties as it moves north and east. The small seeds are probably most often distributed by air currents and machinery along transportation corridors such as roads and railroads (Thompson et al. 2013). The Queens County population in Alley Pond Park at Restoration Pond, along the Cross Island Parkway, tripled in size from 2019 to 2021.

ACKNOWLEDGEMENTS

The authors are grateful to David Werier for review of the manuscript and helpful comments. The New York City Department of Parks and Recreation is acknowledged for their ongoing efforts and collaboration to document and conserve the indigenous flora of New York City. We thank naturalists in the region such as Sara Rall, Susan Hewitt, Matthew Wills, Sandy Wolkenberg, and the iNaturalist users pduhon, agentcolson, and other keen observers of the local flora who helpfully share their observations and comments via iNaturalist. Funding for the New York City EcoFlora was provided in part by the Institute of Museum and Library Services grant to the New York Botanical Garden (MG-70-19-0057).

LITERATURE CITED

- Al-Shehbaz, I. 2009. *Thlaspi*. Pp. 745–746, in Flora of North America Editorial Committee (eds.) Flora of North America North of Mexico, Vol. 7: Magnoliophyta: Salicaceae to Brassicaceae. Oxford Univ. Press, New York and Oxford.
- Cusick, A.W. 2015. *Thlaspi alliaceum* L. (Brassicaceae): An exotic, invasive annual rapidly spreading in Pennsylvania, together with the first collections of this species in North America. *Castanea* 80: 43–44.
- Esmailbegi, S., I.A. Al-Shehbaz, M. Pouch, T. Mandáková, K. Mummenhoff, M.R. Rahiminejad, M. Mirtadzadini, and M.A. Lysak. 2018. Phylogeny and systematics of the tribe Thlaspideae (Brassicaceae) and the recognition of two new genera. *Taxon* 67: 324–340.
- GBIF. 2021. Global Biodiversity Information Facility Occurrence Download < <https://doi.org/10.15468/dl.cqpc4x>> Accessed 27 May 2021.

- Lamont, E.E. and S.M. Young. 2006. Noteworthy plants reported from the Torrey Range — 2004 and 2005. *J. Torr. Bot. Soc.* 133: 648–659.
- Thompson, R.L., D.B. Poindexter, K. Rivers Thompson and P.F. Threadgill. 2013. *Thlaspi alliaceum* (Brassicaceae) naturalized in Georgia, Missouri, and North Carolina. *Phytoneuron* 2013–86: 1–13.
- USDA, NRCS. 2021. The PLANTS Database. National Plant Data Team, Greensboro, North Carolina. <<http://plants.usda.gov>> Accessed 21 April 2021.
- Werier, D. 2017. Catalogue of the vascular plants of New York State. *Mem. Torrey Bot. Soc.* 27: 1–542.