City University of New York (CUNY)

CUNY Academic Works

Publications and Research

CUNY Graduate Center

2018

Is a Meta-analytic Approach to Burnout's Prevalence Timely?

Eric Laurent

Université Bourgogne - Franche-Comté

Irvin Sam Schonfeld

CUNY Graduate Center

Renzo Bianchi

Université de Neuchâtel

[How does access to this work benefit you? Let us know!](#)

More information about this work at: https://academicworks.cuny.edu/gc_pubs/477

Discover additional works at: <https://academicworks.cuny.edu>

This work is made publicly available by the City University of New York (CUNY).

Contact: AcademicWorks@cuny.edu

Is a meta-analytic approach to burnout's prevalence timely?

This letter to the editor concerns the following article:

Cañadas-De la Fuente GA, Gómez-Urquiza JL, Ortega-Campos EM, et al. Prevalence of burnout syndrome in oncology nursing: a meta-analytic study. *Psychooncology* in press. doi:10.1002/pon.4632

Cañadas-De la Fuente et al¹ performed a meta-analysis of the prevalence of emotional exhaustion, depersonalization, and diminished personal accomplishment—the three definitional components of the burnout syndrome—among nursing professionals working in oncology units. They found that 30% of these workers presented with emotional exhaustion; 15%, with depersonalization; and 35%, with diminished personal accomplishment. The authors concluded that burnout is a problem of considerable importance among nursing professionals employed in oncology units. Although we agree that oncology nursing can be a stressful occupation, we have concerns regarding the relevance and interpretability of the findings of Cañadas-De la Fuente et al.¹

The estimates of burnout's prevalence that the authors included in their meta-analysis have been established with (different versions of) the Maslach Burnout Inventory, using the tercile-based cutoff scores presented in the manual of the questionnaire.² As pointed out in several recent publications,^{3–5} these categorization criteria are devoid of any clinical or theoretical underpinning and are not diagnostic criteria. Therefore, relying on such categorization criteria for estimating the prevalence of burnout is unwarranted. In order to be valid, meta-estimates need to be based on primary estimates that are themselves valid. Unfortunately, this condition cannot be met in research on burnout's prevalence.

We note that some investigators have begun the groundwork required for defining burnout in a clinically meaningful way. Lundgren-Nilsson et al,⁶ for instance, attempted to establish a cutoff score for identifying burnout based on an 18-item version of the Shirom-Melamed Burnout Questionnaire—an alternative measure of burnout symptoms. In order to accomplish their goal, these authors relied on a sample of patients seeking medical care at a specialized outpatient stress clinic and fulfilling the ICD-10 criteria for “other reactions to severe stress” (F.43.8A; Swedish amended version). The work of Lundgren-Nilsson et al⁶ was only preliminary. Moreover, it did not reflect a consensual view of burnout. Indeed, the definition of burnout endorsed by these authors differs from the definition of burnout associated with the Maslach Burnout Inventory. However, the work of Lundgren-Nilsson et al⁶ at least provides investigators with a cutoff score having explicit external reference points rather than reflecting a clinically groundless sample split.

All in all, the meta-analysis of Cañadas-De la Fuente et al¹ is inconclusive because of the very state of burnout research. Because there is

mounting evidence that burnout is a depressive condition,⁷ we recommend that investigators focus on depression, rather than burnout, in occupational health research and practice.

CONFLICTS OF INTEREST

None.

ORCID

Renzo Bianchi <http://orcid.org/0000-0003-2336-0407>

Eric Laurent¹

Irvin Sam Schonfeld²

Renzo Bianchi³

¹Laboratory of Psychology (EA 3188), Bourgogne Franche-Comté University, Besançon, France

²Department of Psychology, The City College of the City University of New York, New York, NY, USA

³Institute of Work and Organizational Psychology, University of Neuchâtel, Neuchâtel, Switzerland

Correspondence

Renzo Bianchi, Institute of Work and Organizational Psychology, University of Neuchâtel, Émile-Argand 11, 2000 Neuchâtel, Switzerland.
Email: renzo.bianchi@unine.ch

REFERENCES

1. Cañadas-De la Fuente GA, Gómez-Urquiza JL, Ortega-Campos EM, et al. Prevalence of burnout syndrome in oncology nursing: a meta-analytic study. *Psychooncology* in press. doi:https://doi.org/10.1002/pon.4632
2. Maslach C, Jackson SE, Leiter MP. *Maslach Burnout Inventory Manual*. 4th ed. Palo Alto, CA: Consulting Psychologists Press; 2016.
3. Bianchi R, Schonfeld IS, Laurent E. The dead end of current research on burnout prevalence. *J Am Coll Surg*. 2016;223(2):424–425. https://doi.org/10.1016/j.jamcollsurg.2016.05.012
4. Bianchi R, Schonfeld IS, Laurent E. Can we trust burnout research? *Ann Oncol*. 2017;28(9):2320–2321. https://doi.org/10.1093/annonc/mdx267
5. Schears RM. Defining physician burnout, and differentiating between burnout and depression—II. *Mayo Clin Proc*. 2017;92(9):1455–1456. https://doi.org/10.1016/j.mayocp.2017.07.006
6. Lundgren-Nilsson Å, Jonsdottir IH, Pallant J, et al. Internal construct validity of the Shirom-Melamed Burnout Questionnaire (SMBQ). *BMC Public Health*. 2012;12(1). https://doi.org/10.1186/1471-2458-12-1
7. Bianchi R, Schonfeld IS, Laurent E. Burnout: moving beyond the status quo. *Int J Stress Manag*. in press. https://doi.org/10.1037/str0000088