

City University of New York (CUNY)

CUNY Academic Works

Publications and Research

New York City College of Technology

2022

Risk Levels and Adverse Clinical Outcomes Among Patients With Nonvalvular Atrial Fibrillation Receiving Oral Anticoagulants

Gregory Lip
University of Liverpool

Richard R. Murphy
Federal Aviation Administration

Farhad Sahiar
Federal Aviation Administration

Timothy J. Ingall
Mayo Clinic

Amol Dhamane
Bristol-Myers Squibb

See next page for additional authors

[How does access to this work benefit you? Let us know!](#)

More information about this work at: https://academicworks.cuny.edu/ny_pubs/936

Discover additional works at: <https://academicworks.cuny.edu>

This work is made publicly available by the City University of New York (CUNY).
Contact: AcademicWorks@cuny.edu

Authors

Gregory Lip, Richard R. Murphy, Farhad Sahiar, Timothy J. Ingall, Amol Dhamane, Mauricio Ferri, Patrick Hlavacek, Madison T. Preib, Allison V. Keshishian, Cristina Russ, Lisa Rosenblatt, Huseyin Yuce, and Steve Deitelzweig



Risk Levels and Adverse Clinical Outcomes Among Patients With Nonvalvular Atrial Fibrillation Receiving Oral Anticoagulants

Gregory Y. H. Lip, MD; Richard R. Murphy, MD; Farhad Sahiar, MD; Timothy J. Ingall, MD, PhD; Amol D. Dhamane, MS; Mauricio Ferri, MD; Patrick Hlavacek, MPH; Madison T. Preib, MPH; Allison Keshishian, MPH; Cristina Russ, MD; Lisa Rosenblatt, MD; Huseyin Yuces, PhD; Steven Deitelzweig, MD

Abstract

IMPORTANCE The CHA₂DS₂-VASc score (calculated as congestive heart failure, hypertension, age 75 years and older, diabetes, stroke or TIA, vascular disease, age 65 to 74 years, and sex category) is the standard for assessing risk of stroke and systemic embolism and includes age and thromboembolic history. To our knowledge, no studies have comprehensively evaluated safety and effectiveness outcomes among patients with nonvalvular atrial fibrillation receiving oral anticoagulants according to independent, categorical risk strata.

OBJECTIVE To evaluate the incidence of key adverse outcomes among patients with nonvalvular atrial fibrillation receiving oral anticoagulants by CHA₂DS₂-VASc risk score range, thromboembolic event history, and age group.

DESIGN, SETTING, AND PARTICIPANTS This cohort study was a retrospective claims data analysis using combined data sets from 5 large health claims databases. Eligible participants were adult patients with nonvalvular atrial fibrillation who initiated oral anticoagulants. Data were analyzed between January 2012 and June 2019.

EXPOSURE Initiation of oral anticoagulants.

MAIN OUTCOMES AND MEASURES We observed clinical outcomes (including stroke or systemic embolism, major bleeding, and a composite outcome) on treatment through study end, censoring for discontinuation of oral anticoagulants, death, and insurance disenrollment. The population was stratified by CHA₂DS₂-VASc risk score; history of stroke, systemic embolism, or transient ischemic attack; and age groups. We calculated time to event, incidence rates, and cumulative incidence for outcomes.

RESULTS We identified 1 141 097 patients with nonvalvular atrial fibrillation; the mean (SD) age was 75.0 (10.5) years, 608 127 patients (53.3%) were men, and over 1 million were placed in the 2 highest risk categories (high risk 1, 327 766 participants; high risk 2, 688 449 participants). Deyo-Charlson Comorbidity Index scores ranged progressively alongside CHA₂DS₂-VASc risk score strata (mean [SD] scores: low risk, 0.4 [1.0]; high risk 2, 4.1 [2.9]). The crude incidence of stroke and systemic embolism generally progressed alongside risk score strata (low risk, 0.25 events per 100 person-years [95% CI, 0.18-0.34 events]; high risk 2, 3.43 events per 100 person-years [95% CI, 3.06-4.20 events]); patients at the second-highest risk strata with thromboembolic event history had higher stroke incidence vs patients at the highest risk score strata without event history (2.06 events per 100 person-years [95% CI, 2.00-3.12 events] vs 1.18 events per 100 person-years [95% CI, 1.14-1.30 events]). Major bleeding and composite incidence also increased progressively alongside risk score strata (major bleeding: low risk, 0.68 events per 100 person-years [95% CI, 0.56-0.82 events]; high risk 2, 6.29 events per 100 person-years [95% CI, 6.21-6.62 events]; composite incidence: 1.22

(continued)

Key Points

Question What are the relative safety and effectiveness outcomes among patients with nonvalvular atrial fibrillation receiving anticoagulants by CHA₂DS₂-VASc risk score range, thromboembolic event history, and age group?

Findings In this cohort study including data from more than 1 million patients, the observed incidence of stroke, systemic embolism, and major bleeding events generally conformed to an expected increasing incidence by risk score.

Meaning These results suggest that the assessment of individual baseline risk profiles can better inform clinical decision-making, research, and policy for this patient population.

+ [Invited Commentary](#)

+ [Supplemental content](#)

Author affiliations and article information are listed at the end of this article.

Open Access. This is an open access article distributed under the terms of the CC-BY-NC-ND License.

Abstract (continued)

events per 100 person-years [95% CI, 1.06-1.41 events]; high risk 2, 10.67 events per 100 person-years [95% CI, 10.26-11.48 events]). The 12-month cumulative incidence proportions for stroke and systemic embolism, major bleeding, and composite outcomes progressed alongside risk score strata (stroke or systemic embolism, 0.30%-1.85%; major bleeding, 0.55%-5.55%; composite, 1.05%-8.23%). Age subgroup analysis followed similar trends.

CONCLUSIONS AND RELEVANCE The observed incidence of stroke or systemic embolism and major bleeding events generally conformed to an expected increasing incidence by risk score, adding insight into the importance of specific risk score range, thromboembolic event history, and age group strata. These results can help inform clinical decision-making, research, and policy.

JAMA Network Open. 2022;5(8):e2229333. doi:10.1001/jamanetworkopen.2022.29333

Introduction

Atrial fibrillation (AF) is the most common cardiac arrhythmia in the US, currently estimated to affect up to 6 million people and a projected 12 million by 2030.^{1,2} The risk of AF rises with age, with a sharp increase at ages 60 to 69 years and progressive increases for ages 70 to 79 and 80 to 89 years.¹⁻³

Patients with AF are nearly 5 times more likely to experience stroke or systemic embolism (SE) as compared with the general population, and stroke or SE is associated with considerable morbidity, mortality, and health care costs among people living with AF—especially those aged 65 years or older.^{4,5} Other potentially debilitating conditions associated with AF include transient ischemic attack (TIA), myocardial infarction (MI), and retinal vascular occlusion (RVO).^{6,7} As with AF, risk for these conditions increases with age.⁸⁻¹⁰ TIA and MI are also independent risk factors for stroke among patients with AF.¹¹ The CHA₂DS₂-VASc score is currently the most common thromboembolism risk categorization method used for patients with AF.^{12,13} CHA₂DS₂-VASc scoring includes congestive heart failure or left ventricular dysfunction, hypertension, age 75 years or older (double-weighted), diabetes, stroke or TIA (double-weighted), vascular disease (including MI), age 65 to 74 years, and sex category (women).¹¹

Oral anticoagulants (OACs) are used for stroke prevention among patients with AF; warfarin was the standard of care until circa 2010, but newer direct OACs (DOACs) have similar effectiveness and safety profiles ranging from noninferior to superior as compared with warfarin and are now recommended for most patients.¹² All OAC therapy is complicated by some risk of major bleeding (MB), which increases at advanced ages.¹⁴ However, some clinicians may overestimate MB risk; subsequently, they may be reluctant to prescribe OACs to eligible older patients who could be exposed to unnecessary stroke risk as a result.¹⁵ Thus, it is important to monitor changes in bleeding risk as the OAC landscape continues to evolve; this is particularly true of older patients as well as those at greater risk of stroke per other CHA₂DS₂-VASc criteria.¹⁴

Several network meta-analyses and studies have evaluated safety and effectiveness among subgroups of patients with clinically relevant comorbidities. The ARISTOPHANES study also stratified results by age category and CHA₂DS₂-VASc score.¹⁶ However, CHA₂DS₂-VASc treatment guidelines for male vs female patients have since changed,¹² and there remains a dearth of data that comprehensively evaluate risk across stroke risk score ranges, thromboembolic event history, or categorical age ranges. It is unclear if the presence of multiple factors (in CHA₂DS₂-VASc risk score), other than prior stroke, would present the same estimate of future stroke risk as a prior stroke—especially for younger individuals. An estimate of differences in risk of future stroke risk by such risk stratifications may aid clinicians and policy makers in identifying more specific at-risk subgroups.

Thus, we undertook this retrospective analysis of a large data set from combined commercial and federal data sources to evaluate the incidence of key adverse events among patients with NVA

who initiated OACs. We categorized all included patients by CHA₂DS₂-VASc risk score ranges and further stratified results by prior stroke, SE, or TIA and age group.

Methods

Data Sources

We conducted this retrospective analysis with combined data sets from the Optum Insight Research Database (January 2012 through March 2019), Humana (January 2012 through March 2019), IQVIA LifeLink Health Plan Claims Data set (January 2012 through March 2019), IBM Watson MarketScan (January 2012 through June 2019), and US Centers for Medicare & Medicaid Services (CMS) Medicare (January 2012 through December 2017). Of note, patients with Medicare supplemental plans in MarketScan and IQVIA data were not included in the study to avoid potential duplicates with Medicare Part A and Part B. More details on the data sets have been published in previous articles.¹⁶

Both the data sets and the security of the offices where analysis was completed (and where the data sets are kept) met the requirements of the Health Insurance Portability and Accountability Act of 1996. Solutions IRB determined this study to be exempt from the Office for Human Research Protections Regulations for the Protection of Human Subjects (45 CFR 46) under exemption 4: research involving the collection or study of existing data, documents, records, pathological specimens, or diagnostic specimens, if these sources are publicly available or if the information is recorded by the investigator in such a manner that subjects cannot be identified, directly or through identifiers linked to the subjects. The HIPAA Authorization Waiver was granted in accordance with the specifications of 45 CFR 164.512(i). This project was conducted in full accordance with all applicable laws and regulations, and adhered to the project plan that was reviewed by Solutions IRB. This report follows the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline for observational studies.

We identified all medical claims with *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)*, *International Statistical Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM)*, and *ICD-10 Procedural Coding System* diagnosis and procedure codes as well as Health Care Common Procedure Coding System and *Current Procedural Terminology* codes.

Patient Selection

We included adult patients (as of the index date) with 1 or more claims for an OAC prescription (warfarin, apixaban, dabigatran, rivaroxaban, or edoxaban) during the identification period (January 2013 through June 2019). The first OAC prescription during the identification period was defined as the index date. We further required patients who had 1 or more claims with an AF diagnosis (*ICD-9* code, 427.31; *ICD-10*, I480-I482, I4891) prior to or on the index date. To evaluate baseline demographic and clinical characteristics, we also required patients to have 12 or more months of health plan enrollment with medical and pharmacy benefits prior to the index date (baseline period). We excluded patients with evidence of rheumatic mitral valvular heart disease, heart valve replacement or transplantation, venous thromboembolism, or transient AF (within 12 months prior to or on the index date); hip or knee replacement or transplantation (within 6 weeks prior to the index date); pregnancy during the study period; or 1 or more previous OAC prescriptions during the 12 months prior to the index date. Patients with a follow-up time of zero days and those with more than 1 OAC prescription claim on the index date were also excluded. To isolate time on treatment, we observed patients from the day after the index date through the earliest of OAC discontinuation, death, disenrollment, or study end (eTable 1 in the [Supplement](#)).

Baseline Characteristics

We evaluated baseline demographic and clinical characteristics for the overall sample and for risk strata defined by mean CHA₂DS₂-VASc score on the index date as follows: low risk (men, 0; women,

1); moderate risk (men, 1; women, 2); high risk 1 (men, 2 to 3; women, 3); and high risk 2 (men and women, 4 or more). The CHA₂DS₂-VASc risk scores were calculated based on the presence (or absence) of individual components during the 12 months prior to and on the index date. Within risk groups, we further stratified results by age group (18 to 54, 55 to 64, 65 to 74, and 75 years or older); baseline stroke, SE, or TIA history (those with an event history vs no event history, which may also be viewed as secondary vs primary prevention strategy); sex (men, women); and baseline stroke, SE, or TIA history. We captured age, sex, and Deyo-Charlson comorbidity index (CCI) score on the index date as well as key comorbidities and baseline medications of interest during the 12 months prior to and on the index date (eTable 2 in the [Supplement](#)).

Outcomes

For the overall sample and within each strata of interest, we reported crude incidence rates for stroke or SE, MB, TIA, RVO, MI, and a composite of all outcomes. We also reported cumulative incidence for stroke or SE, MB, and composite outcomes. The outcomes were defined by hospitalization with a primary diagnosis of the respective outcomes (eTable 3 in the [Supplement](#)).

Statistical Analysis

We described baseline demographic and clinical characteristics with standard summary statistics, with numbers and percentages reported for categorical variables and means, medians, standard deviations (SDs) and with interquartile ranges reported for continuous variables. Because not all of the databases have race included as a variable, race and ethnicity were not a part of the study. We evaluated cohort differences in categorical and binary variables using the χ^2 test or Fisher exact test and compared continuous variables using the Student *t* test or Wilcoxon rank sum test. We presented clinical outcome incidence rates per 100 person-years (PY) overall and within risk score, age, and event history strata—calculated as the number of patients with events of interest divided by time at risk for developing the event (multiplied by 100). Unadjusted Kaplan-Meier survival curves were generated to evaluate the cumulative incidence of the outcomes. The a priori significance level of 2-sided *P* < .05 was used to perform the analysis. All analysis was conducted with statistical software SAS version 9.4 (SAS Institute Inc).

Results

After applying the inclusion and exclusion criteria, we included 1 141 097 patients with NVAf who initiated an OAC. Among the overall study population, we categorized 29 298 (2.6%) as low risk, 95 584 (8.4%) as moderate risk, 327 766 (28.7%) as high risk 1, and 688 449 (60.3%) as high risk 2 ([Figure 1](#)).

Baseline Patient Characteristics


The mean (SD) age in the overall sample was 75.0 (10.5) years, with a notable divergence between the low risk (53.9 [8.8] years) and moderate risk (53.9 [8.5] years) vs high risk 1 (71.9 [9.1] years) and high risk 2 (79.0 [8.3] years) cohorts. The majority (608 127 [53.3%]) of patients in the overall sample were men; trends generally aligned among risk cohorts, with the exception of the high risk 2 cohort, which was majority women (408 895 [59.4%]). The overall mean (SD) Deyo-CCI score was 1.6 (1.9); cohort scores ranged progressively from low risk (0.4 [1.0]) to high risk 2 (4.1 [2.9]). The most common comorbidity overall was hypertension (964 137 [84.5%]), followed by coronary artery disease (487 047 [42.7%]), diabetes (410 617 [36.0%]), anemia and coagulation defects (330 848 [29.0%]), and congestive heart failure (328 219 [28.8%]); proportions increased progressively with stroke risk ranges across all comorbidities ([Table 1](#)). DOAC treatment was more common than warfarin (794 670 [69.7%] vs 346 427 [30.3%]), but warfarin proportions increased progressively with risk score range (low risk, 4088 of 29 298 [14.0%] to high risk 2, 235 145 of 688 449 [34.2%]). Trends among patients with and without prior stroke or TIA within the high risk 1 and high risk 2

cohorts followed generally similar trends, but the high risk 2 cohort with prior stroke, SE, or TIA had a higher comorbidity burden (eTable 4 in the Supplement).

Event Incident Rates

The incidence rates for stroke and SE among the overall population were 1.31 events per 100 PY (95% CI, 1.29-1.34 events) and ranged from 0.25 events per 100 PY (95% CI, 0.18-0.34) for individuals with low risk and no stroke, SE, or TIA event history to 3.43 per 100 PY (95% CI, 3.06-4.20) for individuals in the high risk 2 group with thromboembolic event history. MB incidence rates among the overall population were 4.08 events per 100 PY (95% CI, 4.04-4.12) and ranged from 0.68 per 100 PY (95% CI, 0.56-0.82) for individuals in the low risk group with no stroke, SE, or TIA event history to 6.29 per 100 PY (95% CI, 6.21-6.62) for individuals in the high risk 2 group with event history. Composite incidence rates among the overall population were 6.15 events per 100 PY (95% CI, 6.10-6.20) and ranged from 1.22 per 100 PY (95% CI, 1.06-1.41) for individuals with low risk and no stroke, SE, or TIA event history to 10.67 per 100 PY (95% CI, 10.26-11.48) for individuals in the high risk 2 group with event history. Rates for TIA, RVO, and MI followed similar predominantly progressive trends (Table 2). Incidence rates among age groups followed similar, predominantly progressive increasing trends by risk score, but incidence rates tended to peak with patients aged 55 to 64 years for the majority of end points. Notable exceptions were stroke incidence among patients in the high risk 1 group with event history, which peaked with patients aged 18 to 54 years (4.19 per 100 PY; 95% CI, 2.89-6.06 per 100 PY), and MB incidence among patients in the high risk 2 group with event history, which peaked with patients aged 75 years or older (6.89 per 100 PY; 95% CI, 6.70-7.09 per 100 PY) (eTable 5 in the Supplement). Stroke incidence rates stratified by gender also followed progressively increasing trends by risk score. Men with low and high risk 1 scores had higher incidence rates compared with women. Among patients with high risk 2 scores, women had higher incidence rates compared with men (eTable 6 in the Supplement).

Figure 1. Patient Selection Criteria



AF indicates atrial fibrillation; OAC, oral anticoagulant; VTE, venous thromboembolism.

Cumulative Incidence

From OAC initiation, the 12-month cumulative incidence of stroke or SE ranged from 0.30% (low risk) to 1.85% (high risk 2). The 12-month cumulative incidence of MB ranged from 0.55% (low risk) to 5.55% (high risk 2). The 12-month cumulative incidence composite outcomes ranged from 1.05% (low risk) to 8.23% (high risk 2) (Figure 2). The cumulative incidence of TIA, MI, and the composite outcome for the high risk 1 and high risk 2 cohorts stratified by prior stroke, SE, and TIA is illustrated in the eFigure in the Supplement.

MB incidence rates among the patients in the high risk 1 group without stroke, SE, or TIA event history were higher than patients in that group with event history (2.17 per 100 PY vs 1.63 per 100 PY), and MI incidence followed directionally with a more pronounced relative difference (0.51 per 100 PY vs 0.27 per 100 PY). Given that mean Deyo-CCI scores were similar between the subgroups, these findings may be attributable to difference in mean ages and age group distribution (no history: mean [SD] age, 72.1 [9.0] years with 275 161 patients [85.4%] aged 65 years and older vs with history: mean [SD] 59.8 [8.4] years with 1299 patients [22.9%] aged 65 to 74 years and no patients older

Table 1. Demographics and Clinical Characteristics of OAC-Treated Patients With NVAf, by Risk Cohort

Baseline characteristics	Patients, No. (%)				
	Overall (N = 1 141 097)	Low (n = 29 298) ^a	Moderate (n = 95 584) ^a	High risk 1 (n = 327 766) ^a	High risk 2 (n = 688 449) ^a
Age, y					
Mean (SD)	75.0 (10.5)	53.9 (8.8)	53.9 (8.5)	71.9 (9.1)	79.0 (8.3)
18-54	43 326 (3.8)	12 338 (42.1)	15 451 (16.2)	12 507 (3.8)	3030 (0.4)
55-64	107469 (9.4)	16 960 (57.9)	38 313 (40.1)	38 799 (11.8)	13 397 (2.0)
65-74	379 288 (33.2)	0	41 820 (43.8)	164 998 (50.3)	172 470 (25.1)
≥75	611 014 (53.5)	0	0	111 462 (34.0)	499 552 (72.6)
Sex					
Men	608 127 (53.3)	21 873 (74.7)	61 421 (64.3)	245 279 (74.8)	279 554 (40.6)
Women	532 970 (46.7)	7425 (25.3)	34 163 (35.7)	82 487(25.2)	408 895 (59.4)
Deyo-Charlson Comorbidity Index, mean (SD)	1.6 (1.9)	0.4 (1.0)	0.6 (1.3)	1.6 (1.9)	4.1 (2.9)
CHA ₂ DS ₂ -VAsC score, mean (SD)	4.1 (1.9)	0.3 (0.4)	1.4 (0.5)	2.7 (0.5)	5.3 (1.26)
HAS-BLED score, mean (SD)	3.0 (1.4)	0.4 (0.6)	1.5 (0.8)	2.5 (1.0)	3.6 (1.2)
Baseline comorbidities^b					
Bleeding history	216 640 (19.0)	2020 (6.9)	8078 (8.5)	44 424 (13.6)	162 118 (23.6)
Stroke history	143 792 (12.6)	NA	NA	3859 (1.2)	139 933 (20.3)
Congestive heart failure	328 219 (28.8)	NA	2632 (2.8)	32 607 (10.0)	292 980 (42.6)
Diabetes	410 617 (36.0)	NA	2789 (2.9)	68 109 (20.8)	339 719 (49.4)
Hypertension	964 137 (84.5)	NA	46 623 (48.8)	252 619 (77.1)	664 895 (96.6)
Renal disease	268 867 (23.6)	434 (1.5)	3555 (3.7)	40 409 (12.3)	224 469 (32.6)
Liver disease	62 643 (5.5)	760 (2.6)	3404 (3.6)	15 506 (4.7)	42 973 (6.2)
Myocardial infarction	149 478 (13.1)	NA	729 (0.8)	15 607 (4.8)	133 142 (19.3)
Dyspepsia or stomach discomfort	216 485 (19.0)	2904 (9.9)	10 371 (10.9)	44 582 (13.6)	158 628 (23.0)
Nonstroke or SE peripheral vascular disease	266 533 (23.4)	NA	896 (0.9)	22 838 (7.0)	242 799 (35.3)
TIA	147 586 (12.9)	NA	NA	3762 (1.2)	143 824 (20.9)
Anemia and coagulation defects	330 848 (29.0)	1646 (5.6)	8001 (8.4)	57 358 (17.5)	263 843 (38.3)
Alcoholism	20 920 (1.8)	754 (2.6)	2500 (2.6)	7154 (2.2)	10 512 (1.5)
Peripheral artery disease	268 438 (23.5)	NA	896 (0.9)	22 885 (7.0)	244 657 (35.5)
Coronary artery disease	487 047 (42.7)	2227 (7.6)	13 324 (13.9)	102 552 (31.3)	368 944 (53.6)
Index treatment					
Warfarin	346 427 (30.3)	4088 (14.0)	18 505 (19.4)	88 689 (27.1)	235 145 (34.2)
DOAC ^c	794 670 (69.7)	25 210 (86.0)	77 079 (80.6)	239 077 (72.9)	453 304 (65.8)

Abbreviations: DOAC, direct oral anticoagulant; NA, not applicable; NVAf, nonvalvular atrial fibrillation; OAC, oral anticoagulant; SE, systemic embolism; TIA, transient ischemic attack.

^a Patients were stratified by CHA₂DS₂-VAsC score on index: low (0 for men and 1 for women), moderate (1 for men and 2 for women), high risk 1 (2-3 for men and 3 for women), and high risk 2 (≥4 for men and women).

^b Comorbidities were defined using diagnosis codes in any claim position.

^c Including apixaban, dabigatran, rivaroxaban, or edoxaban.

than 75 years). Age is a well-established independent risk factor for MB as well as MI,^{9,17-21} and the average age for first MI is 65.6 years old for men and 72.0 years old for women.²²

Discussion

Our study used large, nationally representative commercial and federal data sets to comprehensively evaluate stroke and SE, MB, and other key adverse events among highly specific groups of patients with NVAf who initiated anticoagulation. Results revealed an older, mostly male study population with considerable comorbidity among those at the highest risk of stroke, which is consistent with reported general AF population characteristics and correlations between increasing age, CHA₂DS₂-VASc scores, and overall comorbidity.^{23,24} Overall crude incidence ranges across key outcomes (stroke and SE, MB) were generally consistent with unadjusted results in the preponderance of similar studies.^{16,25,26} The progressive increase of proportions of patients treated with OAC, in line with general CHA₂DS₂-VASc score ranges, was also consistent with the ARISTOPHANES study.¹⁶ Novel findings included incidence and cumulative incidence of stroke and SE, MB, and composite outcomes that predominantly increased in line with CHA₂DS₂-VASc risk score ranges, with pronounced increases in cumulative incidence across outcomes observed among the high risk 2 cohort.

Notable results included stroke and SE incidence among high risk 1 patients with stroke, SE, or TIA event history (2.06 per 100 PY), which was nearly twice that of patients in the high risk 2 group event history (1.18 per 100 PY) despite lower overall CHA₂DS₂-VASc and Deyo-CCI scores (1.9 vs 3.8) among the former group. These data suggest that prior stroke, SE, or TIA is a significant factor in adverse event outcomes independent of the overall CHA₂DS₂-VASc score and overall comorbidity. Although our combination of descriptive end points and highly specific subgroups preclude direct comparison at this time, these findings align directionally with data on independent risk for prior stroke and SE, which has been reported to increase the relative risk of subsequent stroke by 2.5 times as compared with no prior stroke, SE, or TIA and has remained stable over time (between 1994 and 2019).²⁷⁻²⁹

Among age group strata, overall incidence of adverse events generally tended to peak with the 55-to-64-year-old subgroup and progressively decrease at older strata. Notably, MB incidence peaked with this age group in all risk score event history strata except patients in the low risk group without event history (who had higher incidence in the age 18 to 54 years subgroup) and patients in the high risk 2 group with event history (who had the highest incidence among the age 75 years and older subgroup). This finding appears to be contrary to the abovementioned hypothesis of age distribution in MB incidence among history vs no history subgroups and is inconsistent with age cutoffs for bleeding risk in most risk assessment tools.¹⁷⁻²¹ To the best of our knowledge, there is currently no comparable clinical data that stratifies risk across the same range of age groups;

Table 2. Incidence Rates per 100 Person-Years by Risk Group and Prior Stroke, SE, or TIA

Event	Incidence rate per 100 PY (95% CI)								
	All	Low		Moderate		High 1		High 2	
		No prior	Prior	No prior	Prior	No prior	Prior	No prior	Prior
Stroke	1.31 (1.29-1.34)	0.25 (0.18-0.34)	NA	0.31 (0.27-0.35)	NA	0.56 (0.54-0.60)	2.06 (2.00-3.12)	1.18 (1.14-1.30)	3.43 (3.06-4.20)
MB	4.08 (4.04-4.12)	0.68 (0.56-0.82)		0.94 (0.87-1.02)		2.17 (2.11-2.22)	1.63 (1.34-2.24)	5.26 (5.20-5.40)	6.29 (6.21-6.62)
TIA	0.36 (0.35-0.37)	0.10 (0.07-0.17)		0.11 (0.08-0.13)		0.15 (0.14-0.17)	0.60 (0.41-0.96)	0.30 (0.26-0.38)	1.07 (1.01-1.14)
RVO ^a	0.01 (0.00-0.01)	0.01 (0.00-0.04)		0.00 (0.00-0.01)		0.00 (0.00-0.01)	0.00 (0) ^b	0.01 (0-0.01)	0.01 (0-0.01)
MI	0.92 (0.90-0.94)	0.25 (0.18-0.34)		0.26 (0.22-0.30)		0.51 (0.49-0.54)	0.27 (0.10-0.44)	1.19 (1.16-1.24)	1.34 (1.25-1.38)
Composite ^c	6.15 (6.10-6.20)	1.22 (1.06-1.41)		1.48 (1.39-1.57)		3.14 (3.07-3.20)	3.97 (3.75-5.19)	7.41 (7.31-7.72)	10.67 (10.26-11.48)


Abbreviations: MB, major bleeding; MI, myocardial infarction; PY, person-years; RVO, retinal vascular occlusion; SE, systemic embolism; TIA, transient ischemic attack.

^c The composite score was defined as having any of the events of interest (Stroke/SE, MB, TIA, RVO, or MI).


^b Could not be estimated as there were not enough events to estimate a standard error.

^c P values were significant (P < .05) for all events except RVO.


Figure 2. Cumulative Incidence for Stroke and Systemic Embolism, Major Bleeding, and Composite Events



No. at risk	0	3	6	9	12	15	18	21	24
Low risk	15663	8510	5429	3825	2784	2093	1595	1248	
Moderate risk	61877	40881	29152	21604	16271	12486	9727	7625	
High risk 1	229110	159590	117952	90087	69622	55134	43469	34307	
High risk 2	466398	318504	232474	175576	134183	104994	81963	63863	



No. at risk	0	3	6	9	12	15	18	21	24
Low risk	15654	8506	5428	3826	2782	2088	1593	1246	
Moderate risk	61805	40794	29093	21545	16213	12436	9685	7604	
High risk 1	228339	158866	117350	89558	69148	54715	43105	33986	
High risk 2	462260	315503	230196	173614	132605	103634	80833	62950	



No. at risk	0	3	6	9	12	15	18	21	24
Low risk	15629	8482	5408	3811	2771	2082	1588	1241	
Moderate risk	61698	40679	28984	21453	16149	12386	9635	7562	
High risk 1	227640	158219	116727	88991	68642	54261	42715	33634	
High risk 2	458213	311710	226825	170635	130087	101454	78937	61418	

continued research that compares risk across similarly broad age strata may be useful to clinicians considering OAC treatment for patients at advanced ages. Other unexpected results included stroke incidence among high risk 1 with event history and high risk 2 without event history, which peaked with the 18-to-54-year-old subgroup. This may be attributable to relatively low sample size among the 18-to-54-year-old high risk 1 and 2 subgroups. However, it may also suggest, apart from risk score, age, and event history, influencing factors such as dyslipidemia and tobacco use, or different weighting of factors included in the risk score, which warrants continued investigation.³⁰

In concert, these data add validation to the use of the CHA₂DS₂-VASc score for stroke risk assessment in routine practice as well as more granular data on specific score ranges. Moreover, the data suggest value in considering age and stroke, SE, and TIA history independent of—and in tandem with—stroke and bleeding risk assessment tools, which may aid more precise overall risk assessment for individual patients.

Strengths and Limitations

In addition to this novel granular approach to risk assessment, a key strength of this study is the large and comprehensive sample of patients pooled from diverse and complimentary national commercial and federal data sets. However, the data should be interpreted in the context of several limitations. As with all retrospective designs, the present study was limited to the observation of associations as opposed to inference of causality. While claims data are extremely valuable to augment randomized controlled trial results with observation of real clinical practice, all claims databases also have certain inherent limitations because claims are collected for administrative purposes and not research. For example, the presence of a claim for a filled prescription does not indicate that the medication was taken as prescribed or at all, and the presence of a diagnosis code on a medical claim does not indicate the positive presence of disease, as the diagnosis code may be incorrectly coded or included as rule-out criteria rather than actual disease.

Limitations specific to this study include the differences between databases regarding information on concomitant use of over-the-counter medications such as aspirin or medication provided as samples. In addition, limited laboratory data availability precluded longitudinal evaluation of specific international normalized ratio values among patients treated with warfarin as well as other laboratory values such as creatinine clearance. Results should be interpreted with respect to these limitations.

Conclusions

The findings of this study for incidence of stroke or SE and MB by CHA₂DS₂-VASc risk score categories generally confirm expected increasing incidence by risk score, with more comprehensive insight into the importance of specific clinical risk factors among patients with NVAF receiving anticoagulants. These data can aid in the assessment of individual baseline risk profiles and thereby better inform clinical decision-making, research, and policy for this patient population.

ARTICLE INFORMATION

Accepted for Publication: June 29, 2022.

Published: August 31, 2022. doi:10.1001/jamanetworkopen.2022.29333

Open Access: This is an open access article distributed under the terms of the [CC-BY-NC-ND License](#). © 2022 Lip GYH et al. *JAMA Network Open*.

Corresponding Author: Gregory Y. H. Lip, MD, Price-Evans Professor of Cardiovascular Medicine, Liverpool Centre for Cardiovascular Science, University of Liverpool and Liverpool Heart & Chest Hospital, Thomas Drive, Liverpool L14 3PE, United Kingdom (gregory.lip@liverpool.ac.uk).

Author Affiliations: Liverpool Centre for Cardiovascular Science, University of Liverpool and Liverpool Heart &

Chest Hospital, Liverpool, United Kingdom (Lip); Department of Clinical Medicine, Aalborg Thrombosis Research Unit, Aalborg University, Aalborg, Denmark (Lip); Federal Aviation Administration, Oklahoma City, Oklahoma (Murphy, Sahiar); Mayo Clinic, Scottsdale, Arizona (Ingall); Bristol Myers Squibb, Lawrenceville, New Jersey (Dhamane, Ferri, Rosenblatt); Pfizer, New York, New York (Hlavacek, Russ, Yuce); STATinMED Research, Ann Arbor, Michigan (Preib, Keshishian); New York City College of Technology, City University of New York, New York (Yuce); Department of Hospital Medicine, Ochsner Clinic Foundation, New Orleans, Louisiana (Deitelzweig); Ochsner Clinical School, University of Queensland School of Medicine, Herston, Queensland, Australia (Deitelzweig).

Author Contributions: Dr Lip had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Lip, Sahiar, Dhamane, Ferri, Hlavacek, Preib, Keshishian, Russ.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Lip, Sahiar, Dhamane, Keshishian.

Critical revision of the manuscript for important intellectual content: Lip, Murphy, Sahiar, Ingall, Dhamane, Ferri, Hlavacek, Preib, Russ, Rosenblatt, Yuce, Deitelzweig.

Statistical analysis: Murphy, Sahiar, Dhamane, Preib, Rosenblatt, Yuce.

Obtained funding: Dhamane.

Administrative, technical, or material support: Murphy, Sahiar, Ingall, Dhamane, Hlavacek, Preib, Keshishian, Deitelzweig.

Supervision: Lip, Dhamane, Ferri, Preib, Keshishian, Rosenblatt.

Conflict of Interest Disclosures: Dr Lip reported receiving consultant or speaker fees from Bristol Myers Squibb–Pfizer, Boehringer Ingelheim, and Daiichi-Sankyo outside the submitted work. Dr Hlavacek reported receiving personal fees from Pfizer during the conduct of the study. Dr Keshishian reported employment with STATinMED Research, a paid consultant to Pfizer and Bristol Myers Squibb during the conduct of the study. Dr Russ reported holding stock from Pfizer Inc during the conduct of the study. Dr Rosenblatt reported owning stock from Bristol Myers Squibb during the conduct of the study. Dr Deitelzweig reported research fees from Bristol Myers Squibb and Pfizer during the conduct of the study; he reported research fees from Bristol Myers Squibb, Pfizer, and Alexion outside the submitted work. No other disclosures were reported.

Funding/Support: This study was funded by Bristol Myers Squibb and Pfizer.

Role of the Funder/Sponsor: Bristol Myers Squibb and Pfizer were involved in the design, analysis, and interpretation of the data in this study, as well as the drafting of the manuscript, revising it critically for intellectual content, and final approval of the manuscript. They were not involved in the conduct of the study and the collection and management of data.

REFERENCES

1. Colilla S, Crow A, Petkun W, Singer DE, Simon T, Liu X. Estimates of current and future incidence and prevalence of atrial fibrillation in the U.S. adult population. *Am J Cardiol*. 2013;112(8):1142-1147. doi:10.1016/j.amjcard.2013.05.063
2. Kornej J, Börschel CS, Benjamin EJ, Schnabel RB. Epidemiology of atrial fibrillation in the 21st century: novel methods and new insights. *Circ Res*. 2020;127(1):4-20. doi:10.1161/CIRCRESAHA.120.316340
3. Schnabel RB, Yin X, Gona P, et al. 50 year trends in atrial fibrillation prevalence, incidence, risk factors, and mortality in the Framingham Heart Study: a cohort study. *Lancet*. 2015;386(9989):154-162. doi:10.1016/S0140-6736(14)61774-8
4. Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke*. 1991;22(8):983-988. doi:10.1161/01.STR.22.8.983
5. Yousufuddin M, Young N. Aging and ischemic stroke. *Aging (Albany NY)*. 2019;11(9):2542-2544. doi:10.18632/aging.101931
6. LaMori JC, Mody SH, Gross HJ, et al. Burden of comorbidities among patients with atrial fibrillation. *Ther Adv Cardiovasc Dis*. 2013;7(2):53-62. doi:10.1177/1753944712464101
7. Callizo J, Feltgen N, Ammermann A, et al. Atrial fibrillation in retinal vascular occlusion disease and non-arteritic anterior ischemic optic neuropathy. *PLoS One*. 2017;12(8):e0181766. doi:10.1371/journal.pone.0181766
8. Kleindorfer D, Panagos P, Pancioli A, et al. Incidence and short-term prognosis of transient ischemic attack in a population-based study. *Stroke*. 2005;36(4):720-723. doi:10.1161/01.STR.0000158917.59233.b7
9. Rodgers JL, Jones J, Bolleddu SI, et al. Cardiovascular risks associated with gender and aging. *J Cardiovasc Dev Dis*. 2019;6(2):E19. doi:10.3390/jcdd6020019

10. Scott IU, Campochiaro PA, Newman NJ, Biousse V. Retinal vascular occlusions. *Lancet*. 2020;396(10266):1927-1940. doi:10.1016/S0140-6736(20)31559-2
11. Lip GY, Nieuwlaat R, Pisters R, Lane DA, Crijns HJ. Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach: the Euro Heart Survey on Atrial Fibrillation. *Chest*. 2010;137(2):263-272. doi:10.1378/chest.09-1584
12. January CT, Wann LS, Calkins H, et al. 2019 AHA/ACC/HRS focused update of the 2014 AHA/ACC/HRS Guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. 2019;74(1):104-132. doi:10.1016/j.jacc.2019.01.011
13. Lip GYH, Banerjee A, Boriani G, et al. Antithrombotic therapy for atrial fibrillation: CHEST guideline and expert panel report. *Chest*. 2018;154(5):1121-1201. doi:10.1016/j.chest.2018.07.040
14. Mitrousi K, Lip GYH, Apostolakis S. Age as a risk factor for stroke in atrial fibrillation patients: implications in thromboprophylaxis in the era of novel oral anticoagulants. *J Atr Fibrillation*. 2013;6(1):783.
15. Zathar Z, Karunatileke A, Fawzy AM, Lip GYH. Atrial fibrillation in older people: concepts and controversies. *Front Med (Lausanne)*. 2019;6:175. doi:10.3389/fmed.2019.00175
16. Lip GYH, Keshishian A, Li X, et al. Effectiveness and safety of oral anticoagulants among nonvalvular atrial fibrillation patients. *Stroke*. 2018;49(12):2933-2944. doi:10.1161/STROKEAHA.118.020232
17. Pisters R, Lane DA, Nieuwlaat R, de Vos CB, Crijns HJ, Lip GY. A novel user-friendly score (HAS-BLED) to assess 1-year risk of major bleeding in patients with atrial fibrillation: the Euro Heart Survey. *Chest*. 2010;138(5):1093-1100. doi:10.1378/chest.10-0134
18. Singer DE, Chang Y, Borowsky LH, et al. A new risk scheme to predict ischemic stroke and other thromboembolism in atrial fibrillation: the ATRIA study stroke risk score. *J Am Heart Assoc*. 2013;2(3):e000250. doi:10.1161/JAHA.113.000250
19. Hijazi Z, Lindbäck J, Alexander JH, et al; ARISTOTLE and STABILITY Investigators. The ABC (age, biomarkers, clinical history) stroke risk score: a biomarker-based risk score for predicting stroke in atrial fibrillation. *Eur Heart J*. 2016;37(20):1582-1590. doi:10.1093/eurheartj/ehw054
20. O'Brien EC, Simon DN, Thomas LE, et al. The ORBIT bleeding score: a simple bedside score to assess bleeding risk in atrial fibrillation. *Eur Heart J*. 2015;36(46):3258-3264. doi:10.1093/eurheartj/ehv476
21. Gage BF, Yan Y, Milligan PE, et al. Clinical classification schemes for predicting hemorrhage: results from the National Registry of Atrial Fibrillation (NRAF). *Am Heart J*. 2006;151(3):713-719. doi:10.1016/j.ahj.2005.04.017
22. Benjamin EJ, Muntner P, Alonso A, et al; American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart Disease and Stroke Statistics-2019 update: a report from the American Heart Association. *Circulation*. 2019;139(10):e56-e528. doi:10.1161/CIR.0000000000000659
23. Lakshminarayan K, Anderson DC, Herzog CA, Qureshi AI. Clinical epidemiology of atrial fibrillation and related cerebrovascular events in the United States. *Neurologist*. 2008;14(3):143-150. doi:10.1097/NRL.0b013e31815cfae
24. Chao TF, Lip GYH, Liu CJ, et al. Relationship of aging and incident comorbidities to stroke risk in patients with atrial fibrillation. *J Am Coll Cardiol*. 2018;71(2):122-132. doi:10.1016/j.jacc.2017.10.085
25. Claxton JS, Lutsey PL, MacLehose RF, Chen LY, Lewis TT, Alonso A. Geographic disparities in the incidence of stroke among patients with atrial fibrillation in the United States. *J Stroke Cerebrovasc Dis*. 2019;28(4):890-899. doi:10.1016/j.jstrokecerebrovasdis.2018.12.005
26. Alberts M, Chen YW, Lin JH, Kogan E, Twyman K, Milentijevic D. Risks of stroke and mortality in atrial fibrillation patients treated with rivaroxaban and warfarin. *Stroke*. 2020;51(2):549-555. doi:10.1161/STROKEAHA.119.025554
27. Stroke Risk in Atrial Fibrillation Working Group. Independent predictors of stroke in patients with atrial fibrillation: a systematic review. *Neurology*. 2007;69(6):546-554. doi:10.1212/01.wnl.0000267275.68538.8d
28. Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation: analysis of pooled data from five randomized controlled trials. *Arch Intern Med*. 1994;154(13):1449-1457. doi:10.1001/archinte.1994.00420130036007
29. Oladiran O, Nwosu I. Stroke risk stratification in atrial fibrillation: a review of common risk factors. *J Community Hosp Intern Med Perspect*. 2019;9(2):113-120. doi:10.1080/20009666.2019.1593781
30. Smajlović D. Strokes in young adults: epidemiology and prevention. *Vasc Health Risk Manag*. 2015;11:157-164. doi:10.2147/VHRM.S53203

SUPPLEMENT.**eTable 1.** Inclusion and Exclusion Criteria Codes**eTable 2.** CHA₂DS₂-VASc Score Codes**eTable 3.** Clinical Outcome Diagnostic Codes**eTable 4.** Baseline Characteristics for High Risk 1 and 2 Cohorts**eTable 5.** Incidence Rates of Clinical Outcomes per 100 Person-years Stratified by Age and Risk Level**eTable 6.** Incidence Rates of Clinical Outcomes per 100 Person-years Stratified by Gender and Risk Level**eFigure.** Kaplan-Meier Plot for Primary vs. Secondary Prevention for High Risk 1 and High Risk 2 Groups